

Back to Nature: Combating *Candida albicans* Biofilm, Phospholipase and Hemolysin Using Plant Essential Oils

Ahmed M. El-Baz ¹, Rasha A. Mosbah ², Reham M. Goda ¹, Basem Mansour ³, Taranum Sultana ⁴, Tanya E. S. Dahms ⁴ and Amira M. El-Ganiny ^{5,*}

¹ Microbiology and Biotechnology Department, Faculty of Pharmacy, Delta University for Science and Technology, International coastal road, Gamasa City, Mansoura, 11152 Dakhaliya, Egypt; elbaz_pharmacy@yahoo.com (A.M.E.-B.); Dr_reham_magdy@yahoo.com (R.M.G.)

² Infection control unit, Zagazig University hospitals, 44519 Zagazig, Egypt; rashamosbah@hotmail.com

³ Pharmaceutical Chemistry Department, Faculty of Pharmacy, Delta University for Science and Technology, International Coastal Road, Gamasa City, Mansoura, 11152 Dakhaliya, Egypt; basem2412@yahoo.com

⁴ Department of Chemistry and Biochemistry, University of Regina, Regina, SK S4S 1P4, Canada; taranum.sultana0@gmail.com (T.S.); tanya.dahms@uregina.ca (T.E.S.D.)

⁵ Microbiology and Immunology Department, Faculty of Pharmacy, Zagazig University, 44519 Zagazig, Egypt

* Correspondence: amiraganiny@yahoo.com or AMElganiny@pharmacy.zu.edu.eg; Tel.: 002-010-22799736, Fax: 002-055-2303266




Figure S1. Effect of essential oils on *C. albicans* growth. The growth was assessed by measuring the OD600 of fungal suspensions after overnight incubation with $\frac{1}{4}$ MIC of A- Clove oil, B- Cinnamon oil, C- Jasmine oil, D- Rosemary oil. Comparison was done for EO-treated cultures and untreated ones, the test was done in duplicate. EOS had no statistically

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

significant inhibition of fungal growth. The data shown are the means \pm standard errors. Unpaired t test (two-tailed) was used for statistical analysis.

22
23




Figure S2. Effect of essential oils on *C. albicans* viability. The growth was assessed by measuring viable count after overnight incubation with $\frac{1}{4}$ MIC of **A-** Clove oil, **B-** Cinnamon oil, **C-** Jasmine oil, **D-** Rosemary oil. Comparison was done for EO-treated cultures and untreated ones, the test was done in duplicate. EOS had no statistically significant effect on viability of *Candida* cells. The data shown are the means \pm standard errors. Unpaired t test (two-tailed) was used for statistical analysis.

24
25
26
27
28
29