

Article

## Rat Lung Response to PM<sub>2.5</sub> Exposure under Different Cold Stresses

Bin Luo <sup>1,†,\*</sup>, Hongxia Shi <sup>2,†</sup>, Lina Wang <sup>1</sup>, Yanrong Shi <sup>1</sup>, Cheng Wang <sup>1</sup>, Jingli Yang <sup>1</sup>, Yaxiong Wan <sup>1</sup> and Jingping Niu <sup>1</sup>

<sup>1</sup> Institute of Occupational Health and Environmental Health, School of Public Health, Lanzhou University, Lanzhou 730000, China; E-Mails: wangln14@lzu.edu.cn (L.W.); shiyr13@lzu.edu.cn (Y.S.); wangchen2013@lzu.edu.cn (C.W.); yangjingli2013@lzu.edu.cn (J.Y.); wanyx@lzu.edu.cn (Y.W.); niujingp@lzu.edu.cn (J.N.)

<sup>2</sup> Lanzhou university Second Hospital, Lanzhou 730030, China; E-Mail: shihongxia0@126.com

† These authors contributed equally to this work.

\* Author to whom correspondence should be addressed; E-Mail: luob@lzu.edu.cn; Tel./Fax: +86-0931-8915-008.

External Editor: Paul B. Tchounwou

*Received: 12 October 2014; in revised form: 3 December 2014 / Accepted: 8 December 2014 / Published: 12 December 2014*

---

**Abstract:** Ambient particulate matters and temperature were reported to have additive effects over the respiratory disease hospital admissions and deaths. The purpose of this study is to discuss the interactive pulmonary toxicities of cold stress and fine particulate matter (PM<sub>2.5</sub>) exposure by estimating inflammation and oxidative stress responses. 48 Wistar male rats, matched by weight and age, were randomly assigned to six groups, which were treated with cold stress alone (0 °C, 10 °C, and 20 °C (Normal control)) and cold stresses plus PM<sub>2.5</sub> exposures respectively. Cold stress alone groups were intratracheal instillation of 0.25 mL normal saline, while cold stress plus PM<sub>2.5</sub> exposure groups were intratracheal instillation of 8 mg/0.25 mL PM<sub>2.5</sub>. These procedures were carried out for three times with an interval of 48 hours for each treatment. All rats were sacrificed after 48 hours of the third treatment. The bronchoalveolar lavage fluid (BALF) was collected for analyzing inflammatory cells and cytokines, and lung homogenate MDA was determined for oxidative stress estimation. Results showed higher level of total cell and neutrophil in

the BALF of PM<sub>2.5</sub> exposed groups ( $p < 0.05$ ). Negative relationships between cold stress intensity and the level of tumor necrosis factor alpha (TNF- $\alpha$ ), C-reactive protein (CRP) interleukin-6 (IL-6) and interleukin-8 (IL-8) in BALF were indicated in PM<sub>2.5</sub> exposure groups. Exposure to cold stress alone caused significant increase of inflammatory cytokines and methane dicarboxylic aldehyde (MDA) and decline of superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) activity only in 0 °C exposure group ( $p < 0.05$ ). The two-way ANOVA found significant interactive effects between PM<sub>2.5</sub> exposure and cold stress in the level of neutrophil, IL-6 and IL-8 and SOD activity ( $p < 0.05$ ). These data demonstrated that inflammation and oxidative stress involved in the additive effect of PM<sub>2.5</sub> exposure and cold stress on pulmonary toxicity, providing explanation for epidemiological studies on the health effect of ambient PM<sub>2.5</sub> and cold stress.

**Keywords:** fine particulate matter (PM<sub>2.5</sub>); cold stress; inflammatory response; oxidative stress; rat lung

---

## 1. Introduction

Temperature is one of the most common physical properties in the natural world. Suitable air temperature is critical for the body to sustain health, while abnormal ones, cold or hot, are harmful. Increasing evidences from epidemiological studies have revealed that lower temperature linked to the increase of adverse health effect, including increased hospital admission and death, exasperation of chronic lung diseases (e.g., chronic obstructive pulmonary disease, COPD). Research reported higher number of inflammatory cells in the lungs of healthy subjects after cold air inhalation [1]. Moreover, animal studies have shown that there were pulmonary cilium destruction, lung inflammation and even airway hyper-responsiveness in rats frequently exposed to cold stress [2,3].

As a critical pollutant of air pollution fine particulate matter (PM<sub>2.5</sub>) has been reported closely related to many adverse effects on respiratory system, including the increase of hospital admission and death and exacerbation of chronic pulmonary diseases [4–7]. With the increase of ambient PM<sub>2.5</sub> by 10 mg/m<sup>3</sup>, the exasperation rate of COPD for hospital doubled among 204 US urban counties [8]. Compared with coarse particulate matter (PM<sub>10</sub>), PM<sub>2.5</sub> may cause more adverse effects because of its smaller aerodynamic diameter and bigger particle amount under similar particle concentration [9]. PM<sub>2.5</sub> can traverse the blood-air barrier of alveolar, then deposit at lung [10]. It causes toxicities on alveolar macrophages and bronchial epithelial cells to induce inflammation in lung, the mechanism in which mainly include producing inflammatory factors and ROS [11–13].

Until now, few studies have studied the interactive effect of particulate matter and temperature on pulmonary health [14–18]. The few previous studies have mainly revealed the additive effect of PM<sub>10</sub> and low temperature on the emergency hospital admission and death of pulmonary diseases [15,16,18]. Experimental studies have unveiled that cold stress could inhibit airway cilium movement and phagocytes migration and phagocytosis [19–21]. We could possibly hypothesize that the inhalation of PM<sub>2.5</sub> during cold stress might induce more serious damage to the lung. Besides, cold stress often comes with high concentration of PM<sub>2.5</sub>, such as during the appearance of cold air and in the winter.


Therefore, we conducted a study aiming to evaluate the interactive effect of PM<sub>2.5</sub> and cold stress on lung toxicity by assessing several pulmonary biomarkers.

## 2. Methods

### 2.1. Animals

The experiments were carried out over eight-week-old Wistar male rats (190–225 g), which were purchased from Gansu University of traditional medicine. Forty-eight rats were assigned to six groups matched with age and weight. Rats were maintained in the animal house in standard cages at 20 °C ± 2 °C, RH 40%–60% and 12:12 h light-dark cycle with free access to laboratory chow and tap water throughout the study. All the animal studies were approved according to the Ethics Committee of Animal Care and Experimentation of the National Institute for Environmental Studies, China. After seven days' thermal acclimation, all groups were treated according to study design (Figure 1.).

**Figure 1.** Study design.



### 2.2. PM<sub>2.5</sub> Sampling and Processing

Samples of PM<sub>2.5</sub> were collected between April and May 2014 in Lanzhou, China, were used for the present study. The PM<sub>2.5</sub> samples were collected onto pre-weighed and pre-dried glass fiber filter, using particulate sampler (TH-150CIII, Wuhan Tianhong Instruments Co., Ltd., China) at a flow rate of 100 L/min. Filters were weighed and cut into squares of 1–2 cm<sup>2</sup>, from which particulate matters were removed by agitation for 20 min × 3 times in ultrapure water with ultrasonic shaker. After filtered by six layers of gauze, the solution was centrifuged at 12,000 rpm for 20 min to collect sediment for drying by a vacuum freeze drier (Christ/ALPHA2-4 LD, Germany). The dried PM<sub>2.5</sub> was kept at –20 °C before diluted for experiment.

### 2.3. Cold Stress and PM<sub>2.5</sub> Exposure Protocol

Before experiment, PM<sub>2.5</sub> was diluted in sterile saline in a concentration of 32 mg/mL. Six groups of rats were randomly assigned to different exposure protocols. Protocols were: (1) 20 °C with intratracheal instillation of 0.25 mL sterile saline or PM<sub>2.5</sub> (8 mg/rat) in animal lab (normal temperature control or PM<sub>2.5</sub> exposure group); (2) 4 h of 10 °C exposure with intratracheal instillation of 0.25 mL

sterile saline or PM<sub>2.5</sub> (8 mg/rat) in climate simulator; (3) 4 h of 0 °C exposure with intratracheal instillation of 0.25 mL sterile saline or PM<sub>2.5</sub> (8 mg/rat) in climate simulator. Each protocol was repeated for three times with an interval of 48 h. Therefore, the cumulative dose of PM<sub>2.5</sub> for each rat is 24 mg. Before instillation of PM<sub>2.5</sub> or sterile saline, rats were anesthetized through inhalation of diethyl ether.

#### 2.4. Bronchoalveolar Lavage

Forty-eight hours after the last intratracheal instillation of sterile saline or PM<sub>2.5</sub>, rats were anesthetized with chloral hydrate (7%, 0.6 mL/100g) and sacrificed through abdominal aorta. Immediately after death, the right lungs were temporarily closed with a haemostatic clamp, and the trachea to the right lung was cannulated. The right lungs were lavaged with 4 × 5 mL volumes of sterile saline. The first lavage fluid was placed in a separate tube for tumor necrosis factor alpha (TNF- $\alpha$ ), C-reactive protein (CRP) interleukin-6 (IL-6) and interleukin-8(IL-8) estimations. Tubes were centrifuged at 1500 rpm for 10 min at 4 °C, the supernatant removed and the cell pellet from the first lavage combined with the cells from the same lavage was re-suspended in 1 ml PBS. Total cells were counted directly and neutrophils were counted after stained with Wright-Giemsa dye under Microscope (BX53, Olympus, Japan). The supernatant of the first lavage fluid was kept at –80 °C for estimation of inflammatory cytokines.

#### 2.5. Lung Homogenate

Lung tissue was homogenized in ice-cold saline with glass-homogenizer. The final homogenate concentration was 10%, then centrifuged at 3000 rpm for 10 min at 4 °C to collect supernatant. All supernatant samples were stored at –80 °C for future biochemical analysis.

#### 2.6. Biomarker Estimations

Before detection, the supernatants of BALF and lung homogenate stored at –80 °C were recovered at 37 °C in an incubator. The TNF- $\alpha$ , CRP, IL-6 and IL-8 were detected with ELISA kit (RD Biosciences, USA). The contents of total protein and methane dicarboxylic aldehyde (MDA) and the activity of superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px) were determined with assay kit (Nanjing Jiancheng Bioengineering Institute, Jiangsu, China) .

#### 2.7. Statistics


All data, expressed as mean  $\pm$  SD, were analyzed using SPSS20.0 IBM statistical software for Windows with a two-way ANOVA (PM<sub>2.5</sub>, cold stress as factors). The differences between paired groups were determined using a paired t-test. The level of  $p < 0.05$  was defined as statistical significance.

### 3. Results

#### 3.1. Physiological Conditions

Among the whole exposure process, no rat died because of exposure factor or any other factors. The Figure 2 showed the body weight changes of all animals through the whole exposure process. Compared with the normal temperature group, the body weight changes of all exposure groups showed a stunted grow trend, especially after the first exposure.

**Figure 2.** Body weight changes of all groups. The data are presented as mean of each group, n = 8/group.



**Table 1.** The total cell numbers and neutrophils in bronchoalveolar lavage fluid (BALF) in all experimental groups.

Cells	0 °C + PM <sub>2.5</sub>	0 °C + Saline	10 °C + PM <sub>2.5</sub>	10 °C + Saline	20 °C + PM <sub>2.5</sub>	20 °C + Saline
Total cell (10 <sup>5</sup> )	5.08 ± 1.25 <sup>a,b</sup>	3.14 ± 0.69 <sup>b</sup>	3.82 ± 1.09 <sup>a</sup>	2.79 ± 0.61	3.73 ± 0.93 <sup>a</sup>	2.33 ± 0.95
Neutrophil (10 <sup>5</sup> )	1.76 ± 0.34 <sup>a,b</sup>	0.85 ± 0.40	1.28 ± 0.40	0.91 ± 0.43	0.82 ± 0.37	1.01 ± 0.28

Notes: n = 8/group, instillation of PM<sub>2.5</sub> (8 mg/rat). Data represent as the mean ± SD. Significance: <sup>a</sup> p < 0.05 compared to the corresponding cold stress alone group; <sup>b</sup> p < 0.05 compared to the normal temperature group with saline or PM<sub>2.5</sub>.

#### 3.2. Cell Counting in Bronchoalveolar Lavage Fluid (BALF)


The Table 1 showed total cells and neutrophils in BALF of different exposure groups. The total cell of all PM<sub>2.5</sub> exposure groups and the neutrophils of 0 °C plus PM<sub>2.5</sub> exposure group were higher than their corresponding cold stress alone exposure groups (p < 0.05). Compared with normal temperature groups, both 0 °C plus PM<sub>2.5</sub> exposure group and 0 °C exposure alone group had significantly more total cells, while only 0 °C plus PM<sub>2.5</sub> exposure group had more neutrophils (p < 0.05). Test of between-subjects effects found both cold stress and PM<sub>2.5</sub> exposure had significant effects over

neutrophils ( $p < 0.05$ ), while only PM<sub>2.5</sub> exposure had significant effects over total cell counts ( $p < 0.05$ ). Cold stress and PM<sub>2.5</sub> exposure had additive effects on the neutrophils ( $p < 0.05$ ).

### 3.3. Inflammatory Factors in BALF

In this study, four inflammatory factors (TNF- $\alpha$ , CRP, IL-6, and IL-8) were determined in the BALF, the results of which were depicted in Figure 3. All PM<sub>2.5</sub> exposure groups showed significantly higher level of four factors than their corresponding cold stress alone groups ( $p < 0.05$ ). A negative relationship was found between temperature and BALF level of four factors. There were higher levels of IL-8 and TNF- $\alpha$  in the 0 °C exposure alone group compared with the normal temperature group with saline instillation ( $p < 0.05$ ). However, no significant difference was found in other factors among cold stress alone groups ( $p > 0.05$ ). The two-way ANOVAs verified that cold stress and PM<sub>2.5</sub> had main effects on the contents of TNF- $\alpha$ , CRP, IL-6 and IL-8 ( $p < 0.05$ ), and there were significant interactions only in IL-6 and IL-8 between these two factors ( $p < 0.05$ ).


**Figure 3.** Inflammatory factors level in BALF. (A) TNF- $\alpha$ , (B) CRP, (C) IL-6, (D) IL-8.  $n = 8$ /group, instillation of PM<sub>2.5</sub> (8 mg/rat). Data represent as the mean  $\pm$  SD. Significance: <sup>a</sup>  $p < 0.05$  compared to the corresponding cold stress alone group; <sup>b</sup>  $p < 0.05$  compared to the normal temperature group with saline or PM<sub>2.5</sub>; <sup>c</sup>  $p < 0.05$  compared to 0 °C group with saline or PM<sub>2.5</sub>.




### 3.4. Pulmonary Oxidant and Antioxidants

In order to estimate the effect of cold stress and PM<sub>2.5</sub> on oxidative injury in rat lung, the levels of SOD, GSH-Px and MDA were detected and showed in Figure 4. The levels of SOD, GSH-Px and MDA in groups with PM<sub>2.5</sub> instillation were significantly different from their corresponding cold stress exposure alone groups ( $p < 0.05$ ). The activities of both SOD and GSH-Px were found to have a positive relationship with cold stress, which declined with the increase of cold stress intensity, especially among cold stress plus PM<sub>2.5</sub> exposure groups ( $p < 0.05$ ). The MDA content increased distinctly with the decline of temperature in cold stress plus PM<sub>2.5</sub> exposure groups. However, no obvious difference was found among the GSH-Px activity and MDA content in cold stress alone groups, except the significantly decline in SOD activity. Through the two-way ANOVAs statistic analysis, the main effects were found both in cold stress and PM<sub>2.5</sub> exposure in all those three factors ( $p < 0.05$ ), while their interaction only existed in the SOD activity ( $p < 0.05$ ).

**Figure 4.** The level of oxidant and antioxidants in rat lung. (A) SOD, (B) GSH-Px, (C) MDA.  $n = 8$ /group, instillation of PM<sub>2.5</sub> (8 mg/rat). Data represent as the mean  $\pm$  SD. Significance: <sup>a</sup>  $p < 0.05$  compared to the corresponding cold stress alone group; <sup>b</sup>  $p < 0.05$  compared to the normal temperature group with saline or PM<sub>2.5</sub>; <sup>c</sup>  $p < 0.05$  compared to 0 °C group with saline or PM<sub>2.5</sub>.




(a)



(b)

Figure 4. Cont.



#### 4. Discussions

Epidemiological studies have unveiled the interactive effects of low temperature and PM over respiratory diseases. The purpose of the present study is to estimate their interactions on lung injury through animal experiment for the first time. In a dust-storm of April 2014 in Lanzhou city, the PM<sub>2.5</sub> concentration was high as 752  $\mu\text{g}/\text{m}^3$  and the minimum temperature declined by about 5 °C to 2 °C. As the minute tidal volume of rat is about 0.16 L [22], the PM<sub>2.5</sub> exposure dose for rat in that day could be about 1.7 mg/d, which is about one fifth of the PM<sub>2.5</sub> exposure dose (8 mg/rat) used in our study. Concerning the living condition of rat, the cold stress in this study was 0 °C, 10 °C, 20 °C with a scale of 10 °C. Therefore, this study used 8 mg/d of PM<sub>2.5</sub> exposure and cold stress of 0 °C, 10 °C, and 20 °C as treating factors to estimate the toxicological interactions of PM<sub>2.5</sub> exposure and cold stress on lung.

Both cold stress and PM<sub>2.5</sub> exposure were reported to affect the body development by retarding the body weight growth [23,24]. This study showed similar results, each group with cold stress or PM<sub>2.5</sub> exposure experienced the retarded body weight growth. The total cell influx and TNF- $\alpha$ , CRP, IL-6, and IL-8 in BALF were often used to detect the early inflammatory response in lung [25]. In this study, cold stress alone caused significant increase of total cells, TNF- $\alpha$  and IL-8 in BALF especially in 0 °C group. Additionally, the two-way ANOVAs found its main effect over the four inflammatory factors. These results indicated that cold stress could cause pulmonary inflammation to some extent. Sabnis *et al.* found higher expressions of IL-6 and IL-8 in lung epithelial cells after cold stress of 18 °C for 0–4 h [26]. When combined with PM<sub>2.5</sub> exposure, a negative relationship was found between the neutrophil counts, TNF- $\alpha$ , CRP, IL-6 and IL-8 and cold stresses. This may suggest that PM<sub>2.5</sub> exposure added to the effect of cold stress on lung injury. PM<sub>2.5</sub> exposure alone could induce lung injury represented by the increased level of inflammatory cells and cytokines. The PM<sub>2.5</sub> exposure of three seasons from European cities induced higher level of TNF- $\alpha$  in BALF of Wistar rat [27]. Wang *et al.* revealed that PM<sub>2.5</sub> exposure (0.2 mg/rat, 0.8 mg/rat, 3.2 mg/rat) alone increased the total cells, neutrophils, TNF- $\alpha$  and IL-6 in BALF of Wistar rat [24], consistent with the results obtained in this study. When the interactive effect was considered, PM<sub>2.5</sub> exposure and cold stress had additive effect over the level of neutrophils, IL-6 and IL-8, indicating their additive effect over lung injury. In other

words, PM<sub>2.5</sub> exposure combined cold stress may cause more serious lung inflammation than each of them alone.

It has been suggested that the oxidative stress was one way for ambient particles to affect the lung health [28,29]. The mechanism mainly included attacking pulmonary antioxidant system and promoting the production of ROS in lung, which may be the initiation of inflammation and finally activate apoptosis to induce pulmonary disease [30]. The activity of SOD and GSH-Px and MDA level were often used to estimate the antioxidant system and ROS levels in lung [28,29]. In this study, the activity of SOD and GSH-Px reduced and the MDA level increased after PM<sub>2.5</sub> exposure, particularly in groups with cold stress. The positive relationship was found between the activity of SOD and GSH-Px and cold stress in PM<sub>2.5</sub> exposure groups, while a negative relationship existed between MDA level and cold stress. Besides, there were interactive effects between cold stress and PM<sub>2.5</sub> exposure on SOD activity. These results indicate that PM<sub>2.5</sub> exposure may increase pulmonary oxidative stress, and cold stress may add to that effect to lead more serious lung injury. However, when cold stress alone was considered, the significant effects over SOD and GSH-Px activity were only found in 0 °C group. Previous studies have not laid focus on the effect of cold stress on oxidative stress, but on the structure alteration of airway and inflammatory cytokines. The present results may suggest us that cold stress could have effect over pulmonary antioxidant system under lower temperature stress. Namely, cold stress may also induce lung injury by increasing oxidative stress in lung.

In this study, the obtained results suggested that leukocyte infiltration, pro-inflammatory cytokines and oxidative stress might involve in the adverse effect of PM<sub>2.5</sub> exposure on lung. Although the cold stress has affected the level of some pro-inflammatory cytokines and SOD activity in lung to some degree, it may probably play a supplementary role to enhance the pulmonary adverse effect caused by PM<sub>2.5</sub> exposure. Both PM<sub>2.5</sub> exposure and cold stress were reported to have adverse effects on lung in both *in vivo* and *in vitro* experimental studies [2,3,24,27,28,30]. Until now, no study has explored the interactive effect of PM<sub>2.5</sub> exposure and cold stress on respiratory system in animal experimental studies. Salman *et al.* reported that the phagocytic activity of peritoneal macrophage over latex particles was inhibited after cultured at 24 °C for 60 min [31]. A research over human unveiled that cold stress inhibited leukocyte migration and phagocytic activity of phagocytic cells [19,21]. There is possibility that cold stress may also inhibit the phagocytic activity of pulmonary macrophage over PM<sub>2.5</sub> and enhance the pulmonary toxicity of PM<sub>2.5</sub>. Additionally, experimental study found cilium injury by ciliated epithelial loss after frequently cold stress [19,21]. In a cold environment of 4 °C for 1 h, the constricted pulmonary vascular declined the pulmonary blood volume in rabbit, which inhibited the supply of immune cells in lung and cilium movement [32]. Cilium in respiratory system is the first defense line against PM<sub>2.5</sub>, the injured and retarded cilium would promote the inhalation of PM<sub>2.5</sub> to the lung. If these assumptions are possible, the pulmonary defense system against PM<sub>2.5</sub> will be seriously inhibited by cold stress, which may enhance the pulmonary toxicity caused by PM<sub>2.5</sub> exposure. However, these still need to be analyzed in future studies. Another limitation of our study is that we did not study specific chemical components of PM<sub>2.5</sub>, so we could not identify the causative constituents in the observed adverse effects. Therefore, the correlation between chemical elements and toxic effects should be discussed specifically in the future. Additionally, histological result of the lung was not discussed in this study because of some technique problems, so we could not provide any information about the structure change of lung issue, such as the spaces of alveolar ducts and sacs, the

thickening of interstitial connective tissues, inflammatory cell infiltration, *etc.*, in treated rats vs. control. However, these limitations would be covered in the next step of our project.

## 5. Conclusions

In summary, our results indicated that both cold stress and PM<sub>2.5</sub> exposure could cause adverse effects on lung and cold stress may enhance the toxic effect of PM<sub>2.5</sub> exposure. The interactive toxic mechanism for cold stress and PM<sub>2.5</sub> exposure to induce higher occurrence of respiratory disease events could be through provoking oxidative stress and higher production of pro-inflammatory cytokines.

## Acknowledgements

We thank Dongfang Han for his help in the biochemical analysis. This study was supported by grants from the National Natural Science Foundation of China (41405108) and the Elementary research foundation for Central University of Lanzhou University (861495).

## Author Contributions

Bin Luo was the primary writer and editor of the paper, together with Hongxia Shi who participated in the whole process of the literature review, preparing the paper and finishing the manuscript. Lina Wang, Yanrong Shi, Cheng Wang, Jingli Yang and Yaxiong Wan prepared experimental data and literature review in the paper. Jingping Niu participated in every step of preparing this manuscript.

## Conflicts of Interest

The authors declare no conflict of interest.

## References

1. Larsson, K.; Tornling, G.; Gavhed, D.; Muller-Suur, C.; Palmberg, L. Inhalation of cold air increases the number of inflammatory cells in the lungs in healthy subjects. *Eur. Respir. J.* **1998**, *12*, 825–830.
2. Davis, M.S.; Freed, A.N. Repeated hyperventilation causes peripheral airways inflammation, hyperreactivity, and impaired bronchodilation in dogs. *Am. J. Respir. Crit. Care Med.* **2001**, *164*, 785–789.
3. Davis, M.S.; Schofield, B.; Freed, A.N. Repeated peripheral airway hyperpnea causes inflammation and remodeling in dogs. *Med. Sci. Sports Exerc.* **2003**, *35*, 608–616.
4. Zanobetti, A.; Franklin, M.; Koutrakis, P.; Schwartz, J. Fine particulate air pollution and its components in association with cause-specific emergency admissions. *Environ. Health* **2009**, *8*, doi:10.1186/1476-069X-8-58.
5. Pope, C.R.; Dockery, D.W. Health effects of fine particulate air pollution: Lines that connect. *J. Air Waste Manag. Assoc.* **2006**, *56*, 709–742.
6. Dockery, D.W. Health effects of particulate air pollution. *Ann. Epidemiol.* **2009**, *19*, 257–263.

7. Zanobetti, A.; Schwartz, J. The effect of fine and coarse particulate air pollution on mortality: A national analysis. *Environ. Health Perspect.* **2009**, *117*, 898–903.
8. Dominici, F.; Peng, R.D.; Bell, M.L.; Pham, L.; McDermott, A.; Zeger, S.L.; Samet, J.M. Fine particulate air pollution and hospital admission for cardiovascular and respiratory diseases. *JAMA* **2006**, *295*, 1127–1134.
9. Gilmour, P.S.; Schladweiler, M.C.; Richards, J.H.; Ledbetter, A.D.; Kodavanti, U.P. Hypertensive rats are susceptible to TLR4-mediated signaling following exposure to combustion source particulate matter. *Inhal. Toxicol.* **2004**, *16*, 5–18.
10. Kendall, M. Fine airborne urban particles (PM<sub>2.5</sub>) sequester lung surfactant and amino acids from human lung lavage. *Am. J. Physiol. Lung Cell Mol. Physiol.* **2007**, *293*, 1053–1058.
11. Chen, B.Y.; Chan, C.C.; Lee, C.T.; Cheng, T.J.; Huang, W.C.; Jhou, J.C.; Han, Y.Y.; Chen, C.C.; Guo, Y.L. The association of ambient air pollution with airway inflammation in schoolchildren. *Am. J. Epidemiol.* **2012**, *175*, 764–774.
12. van Eeden, S.F.; Tan, W.C.; Suwa, T.; Mukae, H.; Terashima, T.; Fujii, T.; Qui, D.; Vincent, R.; Hogg, J.C. Cytokines involved in the systemic inflammatory response induced by exposure to particulate matter air pollutants (PM(10)). *Am. J. Respir. Crit. Care Med.* **2001**, *164*, 826–830.
13. Fujii, T.; Hayashi, S.; Hogg, J.C.; Mukae, H.; Suwa, T.; Goto, T.; Vincent, R.; van Eeden, S.F. Interaction of alveolar macrophages and airway epithelial cells following exposure to particulate matter produces mediators that stimulate the bone marrow. *Am. J. Respir. Cell Mol. Biol.* **2002**, *27*, 34–41.
14. Green, R.S.; Basu, R.; Malig, B.; Broadwin, R.; Kim, J.J.; Ostro, B. The effect of temperature on hospital admissions in nine California counties. *Int. J. Public Health* **2010**, *55*, 113–121.
15. Carder, M.; McNamee, R.; Beverland, I.; Elton, R.; Van Tongeren, M.; Cohen, G.R.; Boyd, J.; Macnee, W.; Agius, R.M. Interacting effects of particulate pollution and cold temperature on cardiorespiratory mortality in Scotland. *Occup. Environ. Med.* **2008**, *65*, 197–204.
16. Ren, C.; Tong, S. Temperature modifies the health effects of particulate matter in Brisbane, Australia. *Int. J. Biometeorol.* **2006**, *51*, 87–96.
17. Ren, C.; Williams, G.M.; Tong, S. Does particulate matter modify the association between temperature and cardiorespiratory diseases? *Environ. Health Perspect.* **2006**, *114*, 1690–1696.
18. Li, G.; Tao, H.; Guo, Y. Interaction between inhalable particulate matter and apparent temperature on respiratory emergency room visits of a hospital in Beijing. *J. Environ. Health* **2012**, *6*, 483–486.
19. Polderman, K.H. Mechanisms of action, physiological effects, and complications of hypothermia. *Crit. Care Med.* **2009**, *37*, S186–S202.
20. Shephard, R.J.; Shek, P.N. Cold exposure and immune function. *Can. J. Physiol. Pharmacol.* **1998**, *76*, 828–836.
21. Mourtzoukou, E.G.; Falagas, M.E. Exposure to cold and respiratory tract infections. *Int. J. Tuberc. Lung Dis.* **2007**, *11*, 938–943.
22. Crosfill, M.L.; Widdicombe, J.G. Physical characteristics of the chest and lungs and the work of breathing in different mammalian species. *J. Physiol.* **1961**, *158*, 1–14.
23. Luo, B.; Zhang, S.; Ma, S.; Zhou, J.; Wang, B. Effects of cold air on cardiovascular disease risk factors in rat. *Int. J. Environ. Res. Public Health* **2012**, *9*, 2312–2325.

24. Wang, G.; Zhao, J.; Jiang, S.; Song, W. Rat lung response to ozone and fine particulate matter (PM) exposures. *Environ. Toxicol.* **2013**, *10*, doi:10.1002/tox.21912.
25. Broekhuizen, R.; Vernooij, J.H.; Schols, A.M.; Dentener, M.A.; Wouters, E.F. Leptin as local inflammatory marker in COPD. *Respir. Med.* **2005**, *99*, 70–74.
26. Sabnis, A.S.; Shadid, M.; Yost, G.S.; Reilly, C.A. Human lung epithelial cells express a functional cold-sensing TRPM8 variant. *Am. J. Respir. Cell Mol. Biol.* **2008**, *39*, 466–474.
27. Halatek, T.; Stepnik, M.; Stetkiewicz, J.; Krajnow, A.; Kur, B.; Szymczak, W.; Rydzynski, K.; Dybing, E.; Cassee, F.R. The inflammatory response in lungs of rats exposed on the airborne particles collected during different seasons in four European cities. *J. Environ. Sci. Health A Tox. Hazard Subst. Environ. Eng.* **2011**, *46*, 1469–1481.
28. Li, N.; Kim, S.; Wang, M.; Froines, J.; Sioutas, C.; Nel, A. Use of a stratified oxidative stress model to study the biological effects of ambient concentrated and diesel exhaust particulate matter. *Inhal. Toxicol.* **2002**, *14*, 459–486.
29. Romieu, I.; Garcia-Esteban, R.; Sunyer, J.; Rios, C.; Alcaraz-Zubeldia, M.; Velasco, S.R.; Holguin, F. The effect of supplementation with omega-3 polyunsaturated fatty acids on markers of oxidative stress in elderly exposed to PM(2.5). *Environ. Health Perspect.* **2008**, *116*, 1237–1242.
30. Kooter, I.M.; Frederix, K.; Spronk, H.M.; Boere, A.J.; Leseman, D.L.; van Steeg, H.; ten Cate, H.; Cassee, F.R. Lung inflammation and thrombogenic responses in a time course study of Csb mice exposed to ozone. *J. Appl. Toxicol.* **2008**, *28*, 779–787.
31. Salman, H.; Bergman, M.; Bessler, H.; Alexandrova, S.; Beilin, B.; Djaldetti, M. Hypothermia affects the phagocytic activity of rat peritoneal macrophages. *Acta Physiol. Scand.* **2000**, *168*, 431–436.
32. Khadadah, M.; Mustafa, S.; Elgazzar, A. Effect of acute cold exposure on lung perfusion and tracheal smooth muscle contraction in rabbit. *Eur. J. Appl. Physiol.* **2011**, *111*, 77–81.

© 2014 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).