

Supplementary Materials

XRD Identification of Ore Minerals during Cruises: Refinement of Extraction Procedure with Sodium Acetate Buffer

Jelena Milinovic ^{1,*}, Ágata Alveirinho Dias ^{1,2}, Ana I. Janeiro ¹, Manuel F.C. Pereira ³, Sofia Martins ^{1,4}, Sven Petersen ⁴ and Fernando J.A.S. Barriga ¹

¹ Instituto Dom Luiz (IDL), Faculty of Sciences, University of Lisbon, Building C1, Campo Grande, 1749-016 Lisboa, Portugal, j.milinovic@fct.unl.pt (J.M.), agata.dias@usj.edu.mo (Á.A.D.), smartins@geomar.de (S.M.), aiferreira@fc.ul.pt (A.I.J.), fbarriga@fc.ul.pt (F.J.A.S.B.)

² Institute of Science and Environment, University of Saint Joseph, Rua de Londres 116, Macau 999-078, China

³ CERENA-Natural Resources and Environment Studies Center, Instituto Superior Técnico, University of Lisbon, Av. Rovisco Pais, 1049-001 Lisbon, Portugal, mfc@ist.utl.pt

⁴ GEOMAR-Helmholtz Centre for Ocean Research Kiel, Wischhofstraße 1-3, 24148 Kiel, Germany, spetersen@geomar.de

* Correspondence: j.milinovic@fct.unl.pt

Received: 3 November 2019; Accepted: 8 February 2020; Published: 12 February 2020




Figure S1. XRD-pattern (and identified mineral phases) of the sample GC-627/B-151.

