

Supplementary Materials

Synthesis of oligonucleotides containing 2'-*N*-alkylaminocarbonyl-2'-amino-LNA (2'-urea-LNA) moieties using post-synthetic modification strategy

Shoko Yamashita ^{1,†}, Kodai Nishida ^{1,†}, Takashi Osawa ^{1,2}, Ayumi Nakanishi ¹, Yuta Ito ¹ and Yoshiyuki Hari ^{1,*}

¹ Faculty of Pharmaceutical Sciences, Tokushima Bunri University, Nishihama, Yamashiro-cho, Tokushima 770-8514, Japan

² Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita 565-0871, Japan

* Correspondence: hari@ph.bunri-u.ac.jp

† These authors contributed equally to the work.

Table of contents

1. An example of post-synthetic modification of ON2 (Figure S1)	S2
2. ¹ H and ¹³ C NMR spectra of compound 2 (Figure S2).....	S3
3. ¹ H and ³¹ P NMR spectra of compound 3 (Figure S3).....	S4




Figure S1. An example of post-synthetic modification of **ON2**.





Figure S2. ^1H and ^{13}C NMR spectra of compound 2





Figure S3. ^1H and ^{31}P NMR spectra of compound **3**