

Correction

Correction: Gupta, P.M.; et al. Iron, Anemia, and Iron Deficiency Anemia among Young Children in the United States *Nutrients* 2016, 8, 330

Priya M. Gupta *, Cria G. Perrine, Zuguo Mei and Kelley S. Scanlon

Division of Nutrition, Physical Activity, and Obesity, Centers for Disease Control and Prevention, Atlanta, GA 30341, USA; hkg3@cdc.gov (C.G.P.); zam0@cdc.gov (Z.M.); kxs5@cdc.gov (K.S.S.)

* Correspondence: kso7@cdc.gov; Tel.: +1-908-418-0970

Received: 10 August 2017; Accepted: 14 August 2017; Published: 15 August 2017

We would like to submit the following correction to our recently published paper [1] due to the error in classification of children as anemic. The details are as follows:

- (1) The sentence in abstract “Prevalence of ID, anemia, and IDA among children 1–5 years was 7.1% (5.5, 8.7), 3.2% (2.0, 4.3), and 1.1% (0.6, 1.7), respectively.” has been changed as “Prevalence of ID, anemia, and IDA among children 1–5 years was 7.1% (5.5, 8.7), 3.9% (2.0, 4.3), and 1.1% (0.6, 1.7), respectively.”
- (2) In the last paragraph of Section 2, the sentence “Anemia was defined as hemoglobin concentration <11.0 g/dL” has been changed as “Anemia was defined as hemoglobin concentration <11.0 g/dL for children 6–59 months and <11.5 g/dL for children 60–72 months”.
- (3) In Section 3, the last two sentences have been changed as “The prevalence of anemia and IDA were 3.9% and 1.1%, respectively (Table 1). Approximately 28% of children who were anemic were iron deficient”.
- (4) Table 1 has been changed as:

Table 1. Prevalence of iron deficiency, anemia, and iron deficiency anemia among children 1–5 years in the United States, National Health and Nutrition Examination Survey (NHANES) 2007–2010.

	<i>n</i>	Iron Deficiency ¹		Anemia ²		Iron Deficiency Anemia ³	
		%	95% Confidence Interval	%	95% Confidence Interval	%	95% Confidence Interval
1–5 years (12–71.9 months)	1437	7.1	(5.5, 8.7)	3.9	(2.5, 5.3)	1.1	(0.6, 1.7)
1–2 years (12–35.9 months)	643	13.5*	(9.8, 17.2)	5.4	(3.5, 7.4)	2.7	(1.2, 4.2)
3–5 years (36–71.9 months)	794	3.7	(1.9, 5.5)	3.1	(1.4, 4.7)	-**	-

- (5) The first sentence of the last paragraph in Section 3 has been changed as “The prevalence of ID was higher among children aged 1–2 years ($p < 0.05$)”.
- (6) In Section 4, the sentence “Our results showing the prevalence of anemia and IDA, among children 1–5 years, as 3.2% and 1.1% respectively, suggests little change in these indicators over the past decade.” has been changed as “Our results showing the prevalence of anemia and IDA among children 1–5 years as 3.9% and 1.1%, respectively, suggests little change in these indicators over the past decade.”

We apologize for any inconvenience caused to our readers.

Reference

1. Gupta, P.M.; Perrine, C.G.; Mei, Z.; Scanlon, K.S. Iron, Anemia, and Iron Deficiency Anemia among Young Children in the United States. *Nutrients* **2016**, *8*, 330. [[CrossRef](#)] [[PubMed](#)]



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).