

Review

# Cyclooxygenase-1 (COX-1) and COX-1 Inhibitors in Cancer: A Review of Oncology and Medicinal Chemistry Literature

Alessandra Pannunzio and Mauro Coluccia \*

Department of Pharmacy-Drug Sciences, University of Bari "Aldo Moro", Via E. Orabona 4, I-70125 Bari, Italy; alessandra.pannunzio@uniba.it

\* Correspondence: mauro.coluccia@uniba.it; Tel.: +39-080-544-2788

Received: 4 September 2018; Accepted: 9 October 2018; Published: 11 October 2018



**Abstract:** Prostaglandins and thromboxane are lipid signaling molecules deriving from arachidonic acid by the action of the cyclooxygenase isoenzymes COX-1 and COX-2. The role of cyclooxygenases (particularly COX-2) and prostaglandins (particularly PGE<sub>2</sub>) in cancer-related inflammation has been extensively investigated. In contrast, COX-1 has received less attention, although its expression increases in several human cancers and a pathogenetic role emerges from experimental models. COX-1 and COX-2 isoforms seem to operate in a coordinate manner in cancer pathophysiology, especially in the tumorigenesis process. However, in some cases, exemplified by the serous ovarian carcinoma, COX-1 plays a pivotal role, suggesting that other histopathological and molecular subtypes of cancer disease could share this feature. Importantly, the analysis of functional implications of COX-1-signaling, as well as of pharmacological action of COX-1-selective inhibitors, should not be restricted to the COX pathway and to the effects of prostaglandins already known for their ability of affecting the tumor phenotype. A knowledge-based choice of the most appropriate tumor cell models, and a major effort in investigating the COX-1 issue in the more general context of arachidonic acid metabolic network by using the systems biology approaches, should be strongly encouraged.

**Keywords:** cyclooxygenase-1; cyclooxygenase-2; cancer; inflammation; tumorigenesis; COX-1 inhibitor

## 1. Introduction

Already in 1863, the German pathologist Rudolf C. Virchow had observed that some cancers were inherently associated with white blood cell infiltration [1]. However, it was only at the end of 20th century that an increasing body of research led to propose chronic inflammation as one of the enabling characteristics of cancer development [2,3], and the complex inflammatory network in premalignant and frankly malignant disease is now actively explored with the aim of translating epidemiological and experimental evidence into clinical practice [4].

Cancer may originate in the chronic inflammation setting associated with persistent infections, immune-mediated damage, or prolonged exposure to irritants. On the other hand, the genetic and epigenetic alterations underlying the cancerogenesis process inevitably modify the tissue homeostasis and may induce a chronic inflammatory response. Irrespective of the presumed primary or secondary nature of the process, inflammatory cells and mediators can be detected in most tumor tissues, where they act on both tumor and stromal cells and contribute to determine a tumor-promoting microenvironment [5].

In the general complexity of the inflammatory response (hundreds of chemical mediators have been identified, but how they function in a coordinated manner is still not fully understood), the arachidonic acid (AA) metabolites play a relevant role which appears intertwined with their

functions in cell and tissue homeostasis [6]. Prostaglandins (PGs), including PGD<sub>2</sub>, PGE<sub>2</sub>, PGF<sub>2α</sub>, PGI<sub>2</sub> and thromboxane (TX)A<sub>2</sub> (collectively known also as prostanoids) are produced from arachidonic acid by sequential actions of cyclooxygenases (COX-1 or COX-2) and specific synthases, and they exert their effects in autocrine and/or paracrine manner mainly through G protein-coupled receptors (GPCRs) at the cell surface [7].

The involvement of prostaglandins in cancer was first evidenced in human esophageal carcinoma cells, when their invasive and metastatic potential in nude mice was found to be related to PGE<sub>2</sub> and PGF<sub>2α</sub> production [8]. Elevated levels of PGE<sub>2</sub> have been found in numerous cancers, and its effects on multiple cell signaling pathways involved in tumor malignant phenotype induction and maintenance have been thereafter demonstrated [9–13]. However, PGE<sub>2</sub> is not the only PG involved in carcinogenesis. PGD<sub>2</sub> potentially contributes to the colon cancer risk in ulcerative colitis [14]; more recently, (TX)A<sub>2</sub> was found to be involved in colorectal cancer pathophysiology [15,16], as well as in multiple myeloma [17] and lung [18] cancer cell proliferation.

The committed step of PG biosynthesis is catalyzed by COX-1 and COX-2 isoforms. These enzymes display many similarities in structure and catalytic properties and yield the same product, PGH<sub>2</sub>. However, COX-1 and COX-2 are different in their regulation of expression, tissue distribution, and associated synthases, thus subserving distinct biological tasks. COX-1 is constitutively expressed in most tissues (e.g., platelets, lung, prostate, brain, gastrointestinal tract, kidney, liver and spleen), where the COX-1-derived prostanoids are involved in homeostatic functions. It is generally accepted that COX-1 activity maintains the prostanoid production at a basal rate, and allows a rapid increase when cell membrane remodeling produces a rise of free AA. In contrast, COX-2 is generally considered as the inducible isoform, responsible for enhanced prostanoid production in response to inflammatory stimuli and growth factors during inflammation and various pathological conditions, including cancer [19,20]. It should be noted, however, that the concept of “constitutive” and “inducible” isoforms has been challenged by growing evidence indicating that both isoforms are present in normal tissues and can be up-regulated in various pathological conditions [21]. Both the expression and regulation of COX isoforms have been intensively investigated, and reviews about transcriptional regulatory mechanisms [22], and the regulation of gene expression at the co- or post-transcriptional level [23] have been recently published.

The role of cyclooxygenases (in particular COX-2) and prostaglandin products (in particular PGE<sub>2</sub>) in cancer-related inflammation has been extensively investigated in many neoplastic diseases, including esophageal [24], gastrointestinal [25,26] and pancreatic [27] cancers, breast [28] and cervical [29] cancers, renal [30], prostate [31], and bladder [32] cancers, skin [33,34] and head and neck [35] cancers, hematological tumors [36], and mesothelioma [37]. Tumor cells are often characterized by COX-2 aberrant expression [38,39], resulting from transcriptional and/or post-transcriptional alterations and contributing to tumor diseases, such as in colorectal cancer [40]. Importantly, the tumor-associated aberrant expression of COX-2 is often related to epigenetic alterations affecting COX-2 gene (PTGS2) as well as other genes involved into biosynthesis and signaling of its main prostaglandin products [41].

Epidemiological data are convincingly supported by various *in vivo* experimental systems, including chemically induced or transgenic models of colorectal, breast, and other types of cancer [42]. Finally, the therapeutic potential of aspirin, traditional nonsteroidal anti-inflammatory drugs (NSAIDs) and COX-2-selective inhibitors (coxibs) in human cancer has also been widely explored. A very large body of research, including about 2000 relevant publications only for aspirin, proposes that the anticancer activity of these drugs depends upon their ability to interfere with multiple pro-tumorigenic signaling pathways, some of them COX-dependent [38] and others COX-independent [43].

Unlike COX-2, the role of COX-1 in cancer has generally received less attention. Being usually considered as the constitutively expressed isoform that is involved in homeostatic cell and tissue functions, COX-1 was thought not be involved in carcinogenesis. However, increased levels of COX-1 expression have been occasionally reported in several cancers [44], and it has been shown that the genetic disruption of COX-1 is as effective as COX-2 disruption in reducing intestinal [45] and skin

tumorigenesis [46] in mouse models, thus suggesting that both COX isoforms could cooperate in the cancerogenesis process. In the following, an overview of the available evidence about COX-1 expression and involvement in different neoplastic diseases will be presented. The objective of this review is to go through the COX-1-relevant literature in oncology and medicinal chemistry, thus attempting to fill the existing gap and highlighting the potential future studies.

## 2. COX-1 Involvement in Neoplastic Diseases

**Renal cell carcinoma.** The pathophysiological relevance of COX-2 in renal cell carcinoma (RCC) is generally acknowledged, and recently, it has been shown that COX-2 inhibition enhances the efficacy of immunotherapy and tyrosine kinase inhibitor-based treatment [30]. COX-1 expression, however, has been practically unexplored, except for a report on COX-1 mRNA overexpression in an RCC rat model [47]. Interestingly, two recent immunohistochemical investigations have shown a correlation between COX-1 overexpression and poor prognosis in RCC [48], and the validity of the combined use of COX-1 and VEGF in RCC histopathologic prognosis [49].

**Skin cancer.** The role of cyclooxygenase-dependent signaling in skin pathophysiology and non-melanoma carcinogenesis has been recently reviewed [33]. There is accumulating evidence that cyclooxygenase-2 may be involved in the pathogenesis of non-melanoma skin cancer, whereas COX-1 expression appears unaltered with respect to healthy tissue. At preclinical level, however, genetic studies show that the activity of both COX isoforms is mechanistically involved in the basal cell carcinoma pathogenesis [50,51].

**Head and neck cancer.** There are few data regarding COX-1 in head and neck cancer. A comparison of cyclooxygenase-1 expression levels between cancerous tissue from head and neck cancer patients and normal mucosa was performed by immunohistochemistry, Western blotting, and real-time RT-PCR (reverse transcription polymerase chain reaction), showing COX-1 overexpression in cancer cells and no expression in normal mucosa [52]. A progression-associated up-regulation of COX-1 expression was detected by immunohistochemistry in patients with hyperplasia, dysplasia, and carcinoma of oral mucosa [53] and a major expression of either COX-1 or COX-2 was reported in patients with oral squamous cell carcinoma with respect to normal mucosa [54]. More recently, tumor samples from patients with sebaceous gland carcinoma, an aggressive tumor commonly localized at Meibomian or Zeis glands, were examined by high-throughput tissue microarray for the expression of proteins involved in angiogenesis, inflammation, apoptosis, cell proliferation, cell-to-cell contact, and carcinogenesis. High expression of COX-1 and COX-2 was reported in 97% and 82% of patients, respectively [55]. Interesting implications on the COX-1 role derive also from *in vitro* investigations in head and neck squamous cell carcinoma (HNSCC) cell lines, in which the effects of pharmacologic inhibitors of cyclooxygenases were compared to those of small-interfering RNA as far as cell-growth, vascular endothelial growth factor (VEGF) production, and intracellular signaling are concerned [56,57]. The results showed that COX-2 inhibition blocked VEGF productions in some HNSCC cells; in contrast, other COX-2 (and PGE<sub>2</sub>) expressing HNSCC cells showed little response to COX-2 inhibition, this effect depending upon a differential expression of COX-1.

**Esophageal cancer.** Barrett's esophagus, a complication of the chronic gastro-esophageal reflux disease, represents the best-known risk factor for esophageal adenocarcinoma development. Esophageal cancer prevention strategies [58], the role of inflammatory mediators in the disease pathophysiology [24], as well as encouraging experimental and epidemiological data on chemoprophylaxis with NSAIDs in patients with Barrett's esophagus have been recently reviewed [59], along with a body of evidence supporting a role for COX-2 in the pathogenetic sequence leading to esophageal adenocarcinoma [60,61]. Interestingly, a possible involvement of COX-1 in the disease pathophysiology had already been suggested by the co-expression of both COX isoforms and angiogenic (VEGF-A) and lymphangiogenic (VEGF-C) growth factors in primary human tumor samples [62]. More recently, a possible cooperation of COX-1 and COX-2 isoforms has been suggested in an investigation of the PGE<sub>2</sub> pathway in a rat model of esophageal adenocarcinoma induced by

gastroduodenal reflux resulting from esophagojejunostomy [63], as well as by the finding that in the same experimental model indomethacin (a dual COX-1/COX-2 inhibitor) reduced the inflammatory lesions and tumor development, whereas a selective COX-2 inhibitor (MF-tricyclic) was ineffective [64]. Interestingly, a recent meta-analysis of nine observational studies has shown that both low-dose aspirin and non-aspirin COX inhibitor use is associated with a reduced risk of developing esophageal adenocarcinoma in patients with Barrett's esophagus [65].

**Colorectal cancer.** In the colorectal carcinogenesis, the long-term transition process from normal mucosa to benign adenoma and final carcinoma provides the possibility to adopt preventive measures. Many evidences exist underlying the pathophysiological link between chronic inflammation and colorectal cancer [66], and the role of COX-2 in the carcinogenesis process has been extensively investigated [67,68]. COX-1 and COX-2 mRNA expression profiles were examined in tumor tissue in comparison to normal mucosa in stage III (Dukes' C) colorectal cancer patients by Church et al. [69]. In contrast to the general opinion that constitutive COX-1 was not subject to variable expression, an altered regulation of COX-1 expression between normal and malignant tissues was reported, consistent with a COX-1 role in tumorigenesis. A significant correlation between levels of mRNA for COX-1, COX-2, TGF-beta1 and PGES, and those for proangiogenic factors VEGF-A and VEGF-C were also found in primary adenocarcinomas of the small intestine, thus suggesting a role for these factors in the propagation this rare neoplasia [70]. Interestingly, experimental mouse studies using genetic disruption of COX-1 or COX-2 plus *Apc* genes already 15 years ago suggested that both isoforms were involved in intestinal polyp formation [45]. Moreover, up-regulation of COX-1 protects intestinal stem cells from the DNA-damaging effect of azoxymethane and may play a key role in the early phase of intestinal tumorigenesis [71]. Furthermore, analysis of the expression levels of COX-1, COX-2, and mPGES (the downstream prostaglandin E synthase) in COX-1<sup>-/-</sup> and COX-2<sup>-/-</sup> *Apc*Δ716 double-knockout mice revealed that COX-1 was required from the early stage of intestinal polyp development, and that additional expression of COX-2 together with mPGES was necessary for subsequent accelerated growth of polyps [72]. On this basis, a mechanistic cooperation between COX-1 in the early stage of tumorigenesis and COX-2 in the subsequent polyp growth was proposed. In early stage of intestinal carcinogenesis, COX-1-PGE<sub>2</sub> signaling associated with a suppression of the PG-catabolizing enzyme, 15-prostaglandin-dehydrogenase (15-PGDH), has been proposed to occur before COX-2 induction [73]. Recent studies have also shown that COX-1 is required for the maintenance of anchorage-independent growth ability of colon cancer cells (a key feature of malignant phenotype), as well as for tumor promoter-induced transformation of preneoplastic cells [74]. Interestingly, Li et al. in the same work identified a novel selective COX-1 inhibitor, 6-C-(E-phenylethenyl)-naringenin, a derivative of the flavonoid naringenin, also showing its chemopreventive efficacy in a colon cancer xenograft model.

**Breast cancer.** In the breast cancer microenvironment, there is a complex interplay between tumor and stromal cells, and a relevant contribute of COX-2-derived prostanoids is generally acknowledged [24,75,76]. However, both isoforms are expressed in breast cancer clinical samples, COX-1 being primarily localized in stromal cells [77,78]. Up-regulation of COX-1 gene expression in tumor tissues compared to normal tissue has been demonstrated also by whole genome expression analysis of breast carcinomas [79]. Moreover, induction of cell growth arrest and apoptosis are induced in MCF-7 human breast cell line in vitro by the COX-1 inhibitor FR122047 [80], and additive effects on tumor cell growth by COX-1 and COX-2 inhibitors are induced in vitro [81] and in vivo [82].

**Cervical cancer.** Human papilloma virus (HPV) infection of the cervical epithelium is well regarded as the main cause of cervical cancer [83], and a link between HPV E6 and E7 oncoproteins and COX-2 transcription has been established [84,85]. However, current data supporting a benefit for NSAIDs in the treatment of cervical intraepithelial neoplasia—the premalignant cervical lesion—are uncertain [86]. COX-1 up-regulation in cervical carcinoma was reported more than 10 years ago, and COX-1-dependent autocrine/paracrine regulation of COX-2, PGE<sub>2</sub> receptors, and angiogenic factors was demonstrated in vitro [87]. Interestingly, it has been recently demonstrated that seminal



plasma can promote cervical tumor cell growth *in vitro* and *in vivo* through the activation of inflammatory pathways involving both COX isoforms [88] and the expression of angiogenic chemokines [89]. Moreover, in an *in vitro* investigation of the co-regulation of COX-1 and downstream prostaglandin E synthases (mPGES-1, mPGES-2 and cPGES) in several tumor cell lines including cervical cancer, it was found that COX-1 and mPGES-1 messenger RNA (mRNA) are co-regulated and functionally coupled in basal PGE<sub>2</sub> synthesis [90]. Although few clinical investigations of COX isoforms expression in cervical cancer are available [91,92], these experimental results, along with previous findings on the relationships between COX isoforms expression and radiosensitivity of cervical cancer cell lines [93] suggest that also the COX-1 role in cervical cancer prevention and treatment could be reevaluated [94].

**Endometrial cancer.** It is generally accepted that production of prostaglandins by endometrial epithelial cells under resting conditions is regulated through the constitutive expression of COX-1, whereas under stimulated conditions prostaglandin production is a result of the up-regulation of COX-2. A critical role for COX-2 isoform in the maintenance of endometrial tissue during the menstrual cycle as well as in the progression of endometrial cancer has been already observed [95,96], along with the potential clinical benefit of a selective COX-2 inhibitor in COX-2 positive endometrial cancers [97]. However, a possible involvement of COX-1 in the early stage of endometrial cancer development has been suggested, based on a higher COX-1 mRNA expression in patients with WHO (World Health Organization)-grade G1 and G2 endometrial cancer [98]. Moreover, a possible role for COX-1 in endometrial cancerogenesis has been suggested [99]. In a mechanistic investigation of the oxytocin ability of modulating the invasive properties of human endometrial carcinoma cells, the results showed that the hormone increased the invasive properties of tumor cells through the activation of PIK3/AKT pathway; this in turn led to up-regulation of both COX isoforms and subsequent PGE<sub>2</sub> production. Interestingly, both COX isoforms were necessary and acted cooperatively for the oxytocin-induced invasion, COX-1 triggering the pro-invasive matrix metalloproteinase (MMP)-14 (a major MMP-2 activator), and COX-2 up-regulating MMP-2 expression. Importantly, an over-expression of oxytocin receptor was detected in endometrial cancer patients, thus indicating the clinical relevance of the oxytocin pathway in endometrial cancer progression.

**Ovarian cancer.** Epithelial ovarian cancer (EOC), the most lethal gynecological malignancy mainly occurring in older (postmenopausal) woman, is a highly heterogeneous disease [100], and includes distinct histological subtypes. A number of inflammatory cells and mediators are involved in EOC development and progression, and accumulating clinical and experimental evidence strongly suggests that inflammation might represent a unifying pathophysiological mechanism of ovarian carcinogenesis [101,102]. Interestingly, COX-1 was first identified as ovarian cancer marker two decades ago [103]. Since then, COX-1 over-expression has been reported by several groups in multiple human, mouse, as well as hen models of ovarian cancer [104–109]. COX-1 has been suggested as the major enzyme regulating PGE<sub>2</sub> production in ovarian cancer cells [110], and along with COX-2 it plays an essential role in gonadotropin-induced tumor cell migration and invasion [111]. Thus, it is not surprising that both COX-1 and COX-2 have been included in a panel of inflammatory markers that characterize the rapidly growing and highly aggressive (type II) ovarian carcinomas [112]. Accordingly, the expression of COX-1 is higher in ovarian cancer patients with low CD8+ (cytotoxic T cells) and high CD1a+ (dendritic cells) cell density than in those with high CD8+ cell density [113], and COX-1 overexpression is the characterizing element in a cluster of ovarian cancer patients in which the poor prognosis is associated with an immunosuppressive status [114]. However, a role for COX-2 isoform in ovarian cancer has been also reported in clinical [115], as well as experimental investigations [116,117]. A major contribution for clarifying this apparent controversy comes from a very recent large-scale quantitative analysis of the expression of COX isoforms through The Cancer Genome Atlas (TCGA) dataset [118]. This study revealed markedly higher COX-1 expression than COX-2 in high-grade serous ovarian cancer (HGSOC)—the most aggressive EOC histotype—along with higher COX-1 expression in HGSOC tumors than 10 other tumor types in TCGA. Interestingly,

a similar or higher expression of COX-2 isoform was instead observed for endometrioid, mucinous and clear cell tumors in an independent tissue microarray, thus suggesting a more relevant role for COX-2 in other EOC histotypes. Importantly, genetic knockdown of COX-1 in ovarian cancer cell lines resulted in down-regulation of both PG signaling and multiple pro-tumorigenic pathways, thus strongly encouraging further development of methods to selectively target COX-1 in the management of HGSOE tumors. The COX-1 relevance in ovarian cancer is also indirectly witnessed by the major efficacy of aspirin, a stronger COX-1 than COX-2 inhibitor, in COX-1-overexpressing *in vitro* and *in vivo* models of ovarian cancer [119–122].

**Hematological tumors.** COX-2 expression in hematological malignancies, including chronic lymphocytic leukemia, chronic myeloid leukemia, lymphoma, and multiple myeloma, favors tumor cell growth and survival, and represents a poor prognostic indicator [36]. As for myeloid and lymphoid acute leukemia, the transcription of both COX-1 and COX-2 isoforms in human leukemic blast cells from acute myeloid (AML) and acute lymphoid (ALL) patients has been well documented, but only COX-1 protein is expressed and active. COX-1-derived PGE<sub>2</sub> stimulates the spontaneous growth of AML leukemic blasts *in vitro* [123]. Among the various COX-derived metabolites, only PGE<sub>2</sub> appears to be endowed with a stimulating effect on the growth of leukemic blast cells *in vitro* [124], this suggesting a potential benefit stemming from COX-1 selective inhibition. COX-1, but not COX-2, is expressed and enzymatically active also in primary blasts from patients affected by acute promyelocytic leukemia (APL), a distinct AML subtype characterized by the t(15;17) translocation involving the PML gene on chromosome 15 and the retinoic acid receptor- $\alpha$  (RAR- $\alpha$ ) gene on chromosome 17. A major component of APL therapy is all-*trans* retinoic acid (ATRA), based on its ability of activating gene transcription and tumor cell differentiation [125]. The ATRA-induced cell differentiation program is very complex, as shown by systems analysis of transcriptome and proteome [126], and includes also COX-1, but not COX-2, upregulation [127]. In this respect, it should be noted that the COX-1 selective inhibitor SC-560, as well as the non-selective indomethacin, impair both PGE<sub>2</sub> production and ATRA-dependent differentiation of NB4 leukemic cells, a model of acute promyelocytic leukemia, thus suggesting that COX-1-inhibiting NSAIDs should be avoided in APL patients under ATRA treatment [128].


**COX-1 and cancer stem cells.** According to the cancer stem cell (CSC) model, tumor tissues are hierarchically organized as normal tissues, the cancer stem cells being able to fuel or reinitiate tumor growth, giving rise to the progeny which constitutes the bulk of tumor mass. The CSC niche homeostasis is regulated by a complex system of molecular signals, which include also the COX-derived AA metabolites [129,130]. In more detail, COX-2 is up-regulated in CSCs isolated from distinct tumor histotypes (e.g., in breast, colon and bone tumors), is co-expressed with stemness molecular markers, and promotes CSC growth *in vitro* systems, as recently reviewed by Pang L.Y. et al. [131]. This is not surprising, considering the relevance of COX-2-PGE<sub>2</sub> system in stem cell biology of normal tissues [132], as well as the well-known relevance of COX-2 in cancer. However, some experimental evidence of COX-1 involvement in CSC biology also exists. In the azoxymethane (AOM) murine colon cancer model, the early molecular response of intestinal stem cells to genotoxic insult is driven by COX-1-PGE<sub>2</sub> signaling, and results in increased stem cell survival [71]. In breast CSCs, isolated from primary cultures of spontaneous tumors from HER2/Neu transgenic mice, and characterized by proteomic analysis, both COX-1 and COX-2 genes are overexpressed with respect to non-CSCs. Moreover, both COX isoforms belong to an eight-gene signature that correlates with breast cancer patient survival, thus suggesting a role of both isoforms in breast cancers with HER2 overexpression [133]. More recently, the development of anticancer drug resistance has been demonstrated to depend also upon the activation of mesenchymal stem cells (MSCs), which are able of modulating in many ways the response to chemotherapy [134]. A peculiar occurrence after cisplatin treatment is the secretion of specific polyunsaturated fatty acids (12-oxo-5,8,10-heptadecatrienoic acid and hexadeca-4,7,10,13-tetraenoic acid) that are able to confer resistance to various anticancer drugs. Interestingly, the central enzymes involved in the synthesis of MSC-derived chemoprotective factors

are COX-1 and thromboxane synthase [135], thus suggesting that enzyme inhibition could restore cancer cell sensitivity.

To summarize, there is a rich body of evidence suggesting that also COX-1 is involved in multiple aspects of cancer pathophysiology. In most cases, it seems that COX-1 and COX-2 isoforms operate in a coordinated manner, while in specific conditions a prominent role for COX-1 has been demonstrated. Functional consequences of COX-1 activation have been poorly investigated as compared to COX-2 activation, even though the effects of COX-1-PGE<sub>2</sub> signaling on tumor cell phenotype have been described. However, a proper consideration should be given to the distinct functional modifications associated to COX-1 (or COX-2) expression [136], as shown also by a very recent genomic, lipidomic and metabolomic analysis of cyclooxygenase-null murine fibroblasts [137]. This study, which describes the common and dissimilar functional interactions of the COX isoforms at the cellular level by using a systems biology approach, demonstrates that COX-1 up-regulation results in a distinct “eicosanoid storm” along with an “anti-inflammatory, proinflammatory, and redox-activated” signatures. Even though fibroblasts are not cancer cells, these results suggest that COX-1 activity, when placed in pivotal position, can affect cell behavior even beyond eicosanoid metabolism.

### 3. Antitumor Activity of COX-1 Selective Inhibitors

Specific inhibition of COX-2 has been extensively investigated, and many highly selective COX-2 inhibitors are available [138]. In contrast, relatively few COX-1-selective inhibitors have been described [139], and only some of them have been investigated for anticancer activity (Figure 1).



**Figure 1.** COX-1-selective inhibitors investigated for anticancer properties. SC-560, 5-(4-chlorophenyl)-1-(4-methoxyphenyl)-3-(trifluoromethyl)pyrazole; mofezolac, (3,4-bis(4-methoxyphenyl)-5-isoxazolyl)acetic acid; FR122047, 1-((4,5-bis(4-methoxyphenyl)-2-thiazolyl)carbonyl)-4-methylpiperazine.

**SC-560.** SC-560 is a member of diaryl heterocycle class of COX inhibitors, discovered during the COX-2 inhibitor programs, and originally used as a pharmacologic tool to analyze the role of COX-1-derived prostaglandins in inflammation and pain [140]. The pharmacological action of SC-560 has been explored in various pathological conditions, and its antitumor activity has been investigated in multiple *in vitro* and *in vivo* experimental models of ovarian cancer and colorectal cancer, as well as other tumor histotypes, as described in the following.

Based upon a peculiar histopathological pattern showing elevated levels of COX-1 (not COX-2) in the epithelial compartment of ovarian tumor tissue undergoing extensive angiogenesis, the pharmacological action of SC-560 in ovarian cancer models was first investigated by Gupta R.A. et al. [104]. On ovarian cancer cell lines characterized for COX isoforms expression and activity, SC-560 showed *in vitro* antiproliferative effects similar to those of indomethacin and celecoxib (non-COX isoform selective and COX-2-selective, respectively), and only at doses 50–100 times greater than those achieved in *in vivo* systems [140]. In contrast, only SC-560 was found to inhibit arachidonic acid-induced VEGF secretion at low doses, an effect that was dose-dependently reverted by PGE<sub>2</sub>. The *in vivo* tumor growth inhibitory efficacy of SC-560 was then demonstrated by Daikoku T. et al. in genetically engineered ovarian cancer murine models, in which COX-1 overexpression was common to various disease-associated genetic alterations [105,106]. As far as COX-1 downstream targets are concerned, experimental evidence for PPAR $\delta$ –ERK signaling

involvement was provided in murine, as well as in human ovarian cancer [119]. The *in vivo* activity of SC-560 alone or in various combination protocols has been extensively investigated by Li W. et al. in the SKOV-3 xenograft model [141–147]. In this model, SKOV-3 cells predominantly express the COX-1 isoform, as determined by immunohistochemical analysis of tumor samples [142], and SC-560 treatment at COX-1-specific inhibitory dosages produced a slight to moderate reduction of tumor growth. Interestingly, combination treatment of SC-560 with cisplatin [147] or taxol [144–147] generally provided greater efficacy than individual agents on various pharmacological end-points, including angiogenesis, proliferation and apoptosis. Not surprisingly, a combination of SC-560 with the non-selective inhibitor ibuprofen [141] or the COX-2-selective celecoxib [143] also showed major efficacy in the SKOV-3 system, suggesting that the inhibition of both COX-1 and COX-2 could overcome a potential compensation between the two isoforms in the SKOV-3 xenograft biology. From a mechanistic point of view, the *in vivo* effects of SC-560 in the SKOV-3 system may be related to the treatment-dependent reduction of PGE<sub>2</sub> production and to the associated inhibition of angiogenesis. However, a clear evidence for an exclusive or prevailing COX-1 dependence of prostaglandin production in SKOV-3 cells is lacking. Moreover, it cannot be excluded that in *in vivo* systems SC-560 could inhibit also COX-2 [148], and that COX-independent effects could contribute to antitumor activity. Some concern is partly due to the SKOV-3 cell line itself, which in some cases is reported as negative for COX-1 expression [104]. This is not actually unusual in the cyclooxygenase literature; the contrasting results variably depend upon a different source of the cell line, antibody cross-reactivity or other technical issues [149], and distinct experimental conditions (i.e., *in vitro* versus *in vivo*). Interestingly, SC-560 increases also paclitaxel sensitivity of taxane-resistant ovarian cancer cell lines characterized by MDR1/P-glycoprotein upregulation. A similar effect was produced by the COX-2 selective inhibitor NS398, not modified by PGE<sub>2</sub> addition, thus suggesting a prostaglandin- and COX-independent mechanism [150]. The concrete possibility that SC-560 could represent a promising lead for ovarian cancer treatment is also suggested by the SC-560 belonging to a group of small molecule compounds that are potentially able to target ovarian cancer stem cell (OVCSC)-specific genes [151]. This interesting feature has been recently discovered through the characterization of OVCSC-specific gene expression profiling and a co-expression extrapolation with CMAP, a massive repository of gene expression data that provides information on gene expression modification of several cell lines treated with >1000 bioactive compounds [152].

*In vitro* and *in vivo* activity of SC-560 in comparison to celecoxib against colorectal cancer cells was investigated by Grosch S. et al. [153], by evaluating their effects on survival, cell cycle progression, and apoptosis of colon cancer cell characterized for COX-1 and COX-2 expression. Independently from COX status, both SC-560 and celecoxib affected *in vitro* cell survival and induced a G<sub>0</sub>/G<sub>1</sub> block, whereas only celecoxib induced apoptosis. *In vivo*, both compounds were active towards a HCT-15 (COX-2 deficient) xenograft, but devoid of significant effect on HT-29 (COX-2 expressing) tumors, overall indicating a COX-independent mechanism of action. The effects of SC-560 on HT-29 colon cancer cells have been investigated also by Wu W. K. et al. [154], showing that the growth inhibitory effect was accompanied by G<sub>1</sub>-S transition arrest and phosphoinositide 3-kinase (PI3K)-induced autophagy. Proliferation inhibition and cell-cycle progression arrest induced *in vitro* by SC-560 has been investigated also on HCT-116 colon cancer cells, which had been previously characterized by a total absence of COX expression and activity [155]. Lee et al. showed that the growth inhibitory effect towards HCT-116 cells was affected by the cell cycle regulator protein p21CIP1 [156]. Moreover, Sakoguchi-Okada N. et al. showed that the growth inhibitory and apoptotic effect of SC-560 (and other non-selective and COX-2-selective inhibitors) on HCT-116 cells was associated with inhibition of survivin expression and Wnt/beta-catenin signaling pathway [157]. With the aim of identifying possible COX-independent mechanisms of action of NSAIDs, treatment-induced gene expression modifications of HCT-116 cells were investigated by using suppression subtractive hybridization [158]. Interestingly, two cancer-related genes, NAG-1 (officially known as GDF15), and thymosin  $\beta$ -4 (TMSB4X), were induced by SC-560, thus suggesting potentially contrasting effects on its antitumor



activity. NAG-1 codes indeed for a pleiotropic cytokine of the TGF- $\beta$  superfamily expressed in a broad range of cell types and involved in the cellular stress response program. NAG-1 induction effect may either be tumor-growth suppressive or enhancing, depending upon its induction pattern (acute or sustained) and pathophysiological context [159]. The induction of thymosin  $\beta$ -4 too could potentially compromise the anticancer activity of SC-560, being thymosin  $\beta$ -4 mechanistically involved in the metastatic process [160,161] and associated with poor prognosis in CRC [162]. Cyclooxygenase- and prostaglandin-independent growth inhibitory effects of SC-560 have been reported in vitro in cells representing various other tumor histotypes. The growth and cell-cycle progression of human A549, H460, and H358 lung cancer cells is affected by SC-560 at doses that are definitely higher than those required for COX-inhibiting activity. Interestingly, the growth suppression induced by SC-560, but not celecoxib treatment, was associated with reactive oxygen species production [163]. Multiple myeloma cell proliferation is also inhibited by SC-560 at doses that are 10 times higher than those necessary for enzymatic inhibition [164]. SC-560 treatment of HuH-6 and HA22T/VGH human hepatocellular carcinoma cell lines, expressing both COX isoforms at mRNA and protein level, led to growth inhibition and apoptosis, and inhibited anchorage-independent growth of HuH-6 cells in soft agar, an in vitro marker for malignancy of cancer cells [165]. From a mechanistic point of view, treatment of tumor cells with SC-560, as well as with the coxib CAY10404, was associated with activation of ERK1/2 signaling pathway.

**Mofezolac.** Among the few selective COX-1 inhibitors investigated as potential analgesics and anti-platelet agents, mofezolac has been developed and marketed in Japan as a powerful pain killer [166], and its anticancer activity has been investigated almost exclusively in colorectal cancer experimental models. In vitro treatment of a COX-1-expressing RGMI cell line (non-transformed cells derived from rat gastric mucosa) by mofezolac induces a weak apoptotic effect, substantially prostaglandin-synthesis independent, with respect to indomethacin or sodium diclofenac [167], and a similar behavior was also observed in AGS gastric adenocarcinoma cells, treated at concentrations that were similar to those found at gastric mucosa after oral administration [168]. With regard to in vivo models of colorectal cancer, mofezolac treatment suppresses the development of aberrant crypt foci (putative preneoplastic lesions) induced by azoxymethane (AOM) and reduces the number of intestinal polyps in Apc knockout mice. In both murine models, the efficacy of mofezolac was similar to that of nimesulide, this indicating a tumorigenic role for both cyclooxygenase isoforms in this experimental model [169]. Accordingly, a combination treatment with COX-1- and COX-2-selective inhibitors more effectively suppressed polyp growth than either of the single treatments alone in Apc knockout mice [170]. Besides inhibiting preneoplastic lesions, mofezolac significantly reduces the incidence, multiplicity and volume of azoxymethane-induced rat colon carcinomas [171], thus confirming the COX-1 pathophysiological role in the AOM-induced intestinal carcinogenesis. Interestingly, Mofezolac is also effective in suppressing beef tallow-promoted colon carcinogenesis in rats, thus suggesting a potential benefit for populations with high fat intake [172].

**FR122047.** The COX-1-selective inhibitor, 4,5-bis(4-methoxyphenyl)-2-[(1-methylpiperazin-4-yl) carbonyl]thiazole (FR122047) was originally developed as antiplatelet agent devoid of ulcerogenic effects [173], and investigated as analgesic agent [174], or used as tool for studying the involvement of COX-1 as well as the role of prostanoids generated along the COX-1 and COX-2 pathways in various models of inflammation [175,176]. The antitumor activity of FR122047 has been investigated in vitro on MCF-7 breast cancer cells [80]. FR122047 treatment inhibits in vitro cell growth of MCF-7 cells, and induces apoptotic cell death that is mechanistically independent from treatment-associated ROS production, as well as from PGE<sub>2</sub> production inhibition. Mechanisms of MCF-7 cell death induced by FR12207 were further investigated by the same group [177], showing that FR122047 treatment induces caspase-mediated apoptosis and at the same time stimulates a defensive autophagic response of MCF-7 cells. Interestingly, the inhibition of caspase-9 blocks the cytoprotective autophagic process, thus increasing the susceptibility of MCF-7 cells to FR-122047-induced cell death.

In general, *in vitro* and *in vivo* antitumor properties of COX-1-selective inhibitors resemble those of other NSAIDs and coxibs; in some cases, COX-dependent, and in others, COX-independent mechanisms having been reported. Preliminary screening of the anticancer properties of COX inhibitors is usually performed by investigating the *in vitro* effect upon cell proliferation and cell cycle progression of tumor cells characterized for COX expression and activity. COX-1-selective inhibitors, not different from other NSAIDs or coxibs, impair cell growth and proliferation, this biological end-point being most often associated with cell cycle G1-phase arrest and cell death. These effects are usually observed at compound concentrations that are much higher than those of common cytotoxic drugs [178], and very often higher than clinically relevant COX-inhibitor concentrations. The true effective concentration of COX inhibitors in *in vitro* systems could actually be lower than that indicated, owing to a subtractive interaction with the serum proteins of culture medium [179]. However, whilst considering the protein interaction caveat, it is not surprising that COX-1 inhibitors (as well as traditional NSAIDs and coxibs) have weak antiproliferative effects related to enzymatic inhibition. This probably reflects the weak impact of COX-derived PGs in the maintenance of *in vitro* growth and proliferation condition. In the various inflammatory pathological conditions in which the PG activity has been investigated, these molecules do not operate as primary signals of homeostasis modification, but rather as amplification signals at the tissue level [19]. There is no reason to believe that *in vitro* tumor cell proliferation could be a different situation, as shown also by the lack of efficacy of COX inhibitors against resting tumor cells [180]. Therefore, to obtain more useful information from a basic antitumor assay such as cell growth and proliferation inhibition, it would be preferable to use tumor cells that increase their proliferation rate as a result of PG addition, and to demonstrate that this effect is impaired by the COX inhibitor under investigation. In this context, it would be preferable to integrate the COX expression and activity data into a more general knowledge of genes and gene products that are associated with AA metabolism in the tumor cells under investigation. This information is available to the scientific community in public databases, e.g., the Cancer Cell Line Encyclopedia [181], and it is essential for a reasoned choice of tumor cells to be used, also in relation to their suitability as models of human clinical disease [182]. The elegant paper of Wilson A.J. et al. [118] on the relevance of COX-1 in HGSOc witnesses the effectiveness of this strategy. Moreover, treatment effects upon cell cycle progression and the cell survival–death homeostasis could be more conveniently investigated in the same experimental setting by using modern multiparametric imaging technologies [183], thereby obtaining a comprehensive picture of treatment-associated phenotypic modifications. In *in vitro* systems again, specific investigations could be planned by using 3D tumor cell models enriched with stromal cells, to more closely mimic the tissue-like condition in which PGs usually operate. As far as other tumor phenotype hallmarks potentially affected by COX-derived PGs, such as angiogenesis, invasion and metastasis, and immune effects, they all would only be observed in *in vivo* systems, although some useful indication can be provided by *in vitro* assays measuring cell adhesion properties, migration, invasion, and colony formation [178]. As far as the COX-independent mechanism, which are often described for COX-1-selective inhibitors, they could also be of interest. Unlike what occurred with the COX-independent pharmacological actions of coxibs [184], which gave rise to substantial research and development activity [43], the results so far reported for COX-1-selective inhibitors have not triggered systematic research on this topic. This is not surprising, considering the small number COX-1-selective compounds available. On the other hand, a COX-independent mechanism occurs in an unpredictable manner, and sometimes in a different context, as in the case of the ability of mofezolac to interact with c-myc oncoprotein [185].

#### 4. Conclusions

The involvement of COX-1 activity in cancer seems to have, in most cases, a pathophysiological role that is consistent and coordinated with COX-2, as in other inflammation-associated pathological conditions. However, in some cases, as exemplified by serous ovarian carcinoma, COX-1 overexpression plays a pivotal role, stimulating more in-depth studies on its function in

this specific condition, and suggesting that other histopathological and molecular subtypes of cancer disease could share this feature. However, it seems clear that the analysis of functional implications of COX-1-signaling, as well as of the pharmacological action of COX-1-selective inhibitors, should not be restricted to the COX pathway, and to the effects of prostaglandins that are already known for their ability to affect the tumor phenotype. A knowledge-based choice of the most appropriate tumor cell models, and a major effort in investigating the COX-1 issue in the more general context of the arachidonic acid metabolic network by using systems biology approaches, should be strongly encouraged.

**Author Contributions:** Conceptualization, M.C.; Writing–Original draft, M.C. and A.P.; Writing–Review and Editing, M.C. and A.P.; Funding Acquisition, M.C.

**Funding:** This research was funded by PRIN, 2009WCNS5C.

**Acknowledgments:** The authors thank Federica Campanella for her help in the literature research.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Balkwill, F.; Mantovani, A. Inflammation and cancer: Back to Virchow? *Lancet* **2001**, *357*, 539–545. [[CrossRef](#)]
2. Hanahan, D.; Weinberg, R.A. Hallmarks of cancer: The next generation. *Cell* **2011**, *144*, 646–674. [[CrossRef](#)] [[PubMed](#)]
3. Ben-Neriah, Y.; Karin, M. Inflammation meets cancer, with NF- $\kappa$ B as the matchmaker. *Nat. Immunol.* **2011**, *12*, 715–723. [[CrossRef](#)] [[PubMed](#)]
4. Crusz, S.M.; Balkwill, F.R. Inflammation and cancer: Advances and new agents. *Nat. Rev. Clin. Oncol.* **2015**, *12*, 584–596. [[CrossRef](#)] [[PubMed](#)]
5. Coussens, L.M.; Zitvogel, L.; Palucka, A.K. Neutralizing tumor-promoting chronic inflammation: A magic bullet? *Science* **2013**, *339*, 286–291. [[CrossRef](#)] [[PubMed](#)]
6. Dennis, E.A.; Norris, P.C. Eicosanoid storm in infection and inflammation. *Nat. Rev. Immunol.* **2015**, *15*, 511–523. [[CrossRef](#)] [[PubMed](#)]
7. Hirata, T.; Narumiya, S. Prostanoid receptors. *Chem. Rev.* **2011**, *111*, 6209–6230. [[CrossRef](#)] [[PubMed](#)]
8. Botha, J.H.; Bobinson, K.M.; Ramchurren, N.; Reddi, K.; Norman, R.J. Human esophageal carcinoma cell lines: Prostaglandin production, biological properties, and behavior in nude mice. *J. Natl. Cancer Inst.* **1986**, *76*, 1053–1056. [[PubMed](#)]
9. Wang, D.; Dubois, R.N. Eicosanoids and cancer. *Nat. Rev. Cancer* **2010**, *10*, 181–193. [[CrossRef](#)] [[PubMed](#)]
10. Nakanishi, M.; Rosenberg, D.W. Multifaceted roles of PGE2 in inflammation and cancer. *Semin. Immunopathol.* **2013**, *35*, 123–137. [[CrossRef](#)] [[PubMed](#)]
11. Sha, W.; Brüne, B.; Weigert, A. The multi-faceted roles of prostaglandin E2 in cancer-infiltrating mononuclear phagocyte biology. *Immunobiology* **2012**, *217*, 1225–1232. [[CrossRef](#)] [[PubMed](#)]
12. Oshima, H.; Oshima, M. The inflammatory network in the gastrointestinal tumor microenvironment: Lessons from mouse models. *J. Gastroenterol.* **2012**, *47*, 97–106. [[CrossRef](#)] [[PubMed](#)]
13. Kalinski, P. Regulation of immune responses by prostaglandin E2. *J. Immunol.* **2012**, *188*, 21–28. [[CrossRef](#)] [[PubMed](#)]
14. Vong, L.; Ferraz, J.G.; Panaccione, R.; Beck, P.L.; Wallace, J.L. A pro-resolution mediator, prostaglandin D(2), is specifically up-regulated in individuals in long-term remission from ulcerative colitis. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 12023–12027. [[CrossRef](#)] [[PubMed](#)]
15. Li, H.; Liu, K.; Boardman, L.A.; Zhao, Y.; Wang, L.; Sheng, Y.; Oi, N.; Limburg, P.J.; Bode, A.M.; Dong, Z. Circulating prostaglandin biosynthesis in colorectal cancer and potential clinical significance. *EBioMedicine* **2015**, *2*, 165–171. [[CrossRef](#)] [[PubMed](#)]
16. Dovizio, M.; Tacconelli, S.; Ricciotti, E.; Bruno, A.; Maier, T.J.; Anzellotti, P.; Di Francesco, L.; Sala, P.; Signoroni, S.; Bertario, L.; et al. Effects of celecoxib on prostanoid biosynthesis and circulating angiogenesis proteins in familial adenomatous polyposis. *J. Pharmacol. Exp. Ther.* **2012**, *341*, 242–250. [[CrossRef](#)] [[PubMed](#)]

17. Liu, Q.; Tao, B.; Liu, G.; Chen, G.; Zhu, Q.; Yu, Y.; Yu, Y.; Xiong, H. Thromboxane A2 receptor inhibition suppresses multiple myeloma cell proliferation by inducing p38/c-Jun N-terminal Kinase (JNK) Mitogen-activated Protein Kinase (MAPK)-mediated G2/M progression delay and cell apoptosis. *J. Biol. Chem.* **2016**, *291*, 4779–4792. [[CrossRef](#)] [[PubMed](#)]
18. Li, X.; Tai, H.H. Activation of thromboxane A(2) receptors induces orphan nuclear receptor Nurr1 expression and stimulates cell proliferation in human lung cancer cells. *Carcinogenesis* **2009**, *30*, 1606–1613. [[CrossRef](#)] [[PubMed](#)]
19. Ricciotti, E.; FitzGerald, G.A. Prostaglandins and inflammation. *Arterioscler. Thromb. Vasc. Biol.* **2011**, *31*, 986–1000. [[CrossRef](#)] [[PubMed](#)]
20. Smith, W.L.; DeWitt, D.L.; Garavito, R.M. Cyclooxygenases: Structural, cellular, and molecular biology. *Annu. Rev. Biochem.* **2000**, *69*, 145–182. [[CrossRef](#)] [[PubMed](#)]
21. Zidar, N.; Odar, K.; Glavac, D.; Jerse, M.; Zupanc, T.; Stajer, D. Cyclooxygenase in normal human tissues—Is COX-1 really a constitutive isoform, and COX-2 an inducible isoform? *J. Cell. Mol. Med.* **2009**, *13*, 3753–3763. [[CrossRef](#)] [[PubMed](#)]
22. Kang, Y.J.; Mbonye, U.R.; DeLong, C.J.; Wada, M.; Smith, W.L. Regulation of intracellular cyclooxygenase levels by gene transcription and protein degradation. *Prog. Lipid Res.* **2007**, *46*, 108–125. [[CrossRef](#)] [[PubMed](#)]
23. Lutz, C.S.; Cornett, A.L. Regulation of genes in the arachidonic acid metabolic pathway by RNA processing and RNA-mediated mechanisms. *Wiley Interdiscip. Rev. RNA* **2013**, *4*, 593–605. [[CrossRef](#)] [[PubMed](#)]
24. Zhang, M.; Zhou, S.; Zhang, L.; Ye, W.; Wen, Q.; Wang, J. Role of cancer-related inflammation in esophageal cancer. *Crit. Rev. Eukaryot. Gene Expr.* **2013**, *23*, 27–35. [[CrossRef](#)] [[PubMed](#)]
25. Cheng, J.; Fan, X.M. Role of cyclooxygenase-2 in gastric cancer development and progression. *World J. Gastroenterol.* **2013**, *19*, 7361–7368. [[CrossRef](#)] [[PubMed](#)]
26. Cathcart, M.C.; O’Byrne, K.J.; Reynolds, J.V.; O’Sullivan, J.; Pidgeon, G.P. COX-derived prostanoid pathways in gastrointestinal cancer development and progression: Novel targets for prevention and intervention. *Biochim. Biophys. Acta* **2012**, *1825*, 49–63. [[CrossRef](#)] [[PubMed](#)]
27. Knab, L.M.; Grippo, P.J.; Bentrem, D.J. Involvement of eicosanoids in the pathogenesis of pancreatic cancer: The roles of cyclooxygenase-2 and 5-lipoxygenase. *World J. Gastroenterol.* **2014**, *20*, 10729–10739. [[CrossRef](#)] [[PubMed](#)]
28. Glover, J.A.; Hughes, C.M.; Cantwell, M.M.; Murray, L.J. A systematic review to establish the frequency of cyclooxygenase-2 expression in normal breast epithelium, ductal carcinoma in situ, microinvasive carcinoma of the breast and invasive breast cancer. *Br. J. Cancer* **2011**, *105*, 13–17. [[CrossRef](#)] [[PubMed](#)]
29. Parida, S.; Mandal, M. Inflammation induced by human papillomavirus in cervical cancer and its implication in prevention. *Eur. J. Cancer Prev.* **2014**, *23*, 432–448. [[CrossRef](#)] [[PubMed](#)]
30. Kaminska, K.; Szczylik, C.; Lian, F.; Czarnecka, A.M. The role of prostaglandin E2 in renal cell cancer development: Future implications for prognosis and therapy. *Future Oncol.* **2014**, *10*, 2177–2187. [[CrossRef](#)] [[PubMed](#)]
31. Shao, N.; Feng, N.; Wang, Y.; Mi, Y.; Li, T.; Hua, L. Systematic review and meta-analysis of COX-2 expression and polymorphisms in prostate cancer. *Mol. Biol. Rep.* **2012**, *39*, 10997–11004. [[CrossRef](#)] [[PubMed](#)]
32. Gakis, G. The role of inflammation in bladder cancer. *Adv. Exp. Med. Biol.* **2014**, *816*, 183–196. [[CrossRef](#)] [[PubMed](#)]
33. Elmets, C.A.; Ledet, J.J.; Athar, M. Cyclooxygenases: Mediators of UV-induced skin cancer and potential targets for prevention. *J. Investig. Dermatol.* **2014**, *134*, 2497–2502. [[CrossRef](#)] [[PubMed](#)]
34. Rundhaug, J.E.; Simper, M.S.; Surh, I.; Fischer, S.M. The role of the EP receptors for prostaglandin E2 in skin and skin cancer. *Cancer Metastasis Rev.* **2011**, *30*, 465–480. [[CrossRef](#)] [[PubMed](#)]
35. Mendes, R.A.; Carvalho, J.F.; van der Waal, I. An overview on the expression of cyclooxygenase-2 in tumors of the head and neck. *Oral Oncol.* **2009**, *45*, e124–e128. [[CrossRef](#)] [[PubMed](#)]
36. Ramon, S.; Woeller, C.F.; Phipps, R.P. The influence of Cox-2 and bioactive lipids on hematological cancers. *Curr. Angiogenes.* **2013**, *2*, 135–142. [[CrossRef](#)] [[PubMed](#)]
37. Nuvoli, B.; Galati, R. Cyclooxygenase-2, epidermal growth factor receptor, and aromatase signaling in inflammation and mesothelioma. *Mol. Cancer Ther.* **2013**, *12*, 844–852. [[CrossRef](#)] [[PubMed](#)]
38. Greenhough, A.; Smartt, H.J.; Moore, A.E.; Roberts, H.R.; Williams, A.C.; Paraskeva, C.; Kaidi, A. The COX-2/PGE2 pathway: Key roles in the hallmarks of cancer and adaptation to the tumour microenvironment. *Carcinogenesis* **2009**, *30*, 377–386. [[CrossRef](#)] [[PubMed](#)]



39. Harris, R.E. Cyclooxygenase-2 (cox-2) and the inflammogenesis of cancer. *Subcell. Biochem.* **2007**, *42*, 93–126. [[CrossRef](#)] [[PubMed](#)]
40. Dixon, D.A.; Blanco, F.F.; Bruno, A.; Patrignani, P. Mechanistic aspects of COX-2 expression in colorectal neoplasia. *Recent Results Cancer Res.* **2013**, *191*, 7–37. [[CrossRef](#)] [[PubMed](#)]
41. Cebola, I.; Peinado, M.A. Epigenetic deregulation of the COX pathway in cancer. *Prog. Lipid Res.* **2012**, *51*, 301–313. [[CrossRef](#)] [[PubMed](#)]
42. Fischer, S.M.; Hawk, E.T.; Lubet, R.A. Coxibs and other nonsteroidal anti-inflammatory drugs in animal models of cancer chemoprevention. *Cancer Prev. Res.* **2011**, *4*, 1728–1735. [[CrossRef](#)] [[PubMed](#)]
43. Gurpinar, E.; Grizzle, W.E.; Piazza, G.A. NSAIDs inhibit tumorigenesis, but how? *Clin. Cancer Res.* **2014**, *20*, 1104–1113. [[CrossRef](#)] [[PubMed](#)]
44. Rouzer, C.A.; Marnett, L.J. Cyclooxygenases: Structural and functional insights. *J. Lipid Res.* **2009**, *50*, S29–S34. [[CrossRef](#)] [[PubMed](#)]
45. Chulada, P.C.; Thompson, M.B.; Mahler, J.F.; Doyle, C.M.; Gaul, B.W.; Lee, C.; Tiano, H.F.; Morham, S.G.; Smithies, O.; Langenbach, R. Genetic disruption of Ptgs-1, as well as Ptgs-2, reduces intestinal tumorigenesis in Min mice. *Cancer Res.* **2000**, *60*, 4705–4708. [[PubMed](#)]
46. Tiano, H.F.; Loftin, C.D.; Akunda, J.; Lee, C.A.; Spalding, J.; Sessoms, A.; Dunson, D.B.; Rogan, E.G.; Morham, S.G.; Smart, R.C.; et al. Deficiency of either cyclooxygenase (COX)-1 or COX-2 alters epidermal differentiation and reduces mouse skin tumorigenesis. *Cancer Res.* **2002**, *62*, 3395–3401. [[PubMed](#)]
47. Okamoto, T.; Hara, A.; Hino, O. Down-regulation of cyclooxygenase-2 expression but up-regulation of cyclooxygenase-1 in renal carcinomas of the Eker (TSC2 gene mutant) rat model. *Cancer Sci.* **2003**, *94*, 22–25. [[CrossRef](#)] [[PubMed](#)]
48. Yu, Z.H.; Zhang, Q.; Wang, Y.D.; Chen, J.; Jiang, Z.M.; Shi, M.; Guo, X.; Qin, J.; Cui, G.H.; Cai, Z.M.; et al. Overexpression of cyclooxygenase-1 correlates with poor prognosis in renal cell carcinoma. *Asian Pac. J. Cancer Prev.* **2013**, *14*, 3729–3734. [[CrossRef](#)] [[PubMed](#)]
49. Osman, W.M.; Youssef, N.S. Combined use of COX-1 and VEGF immunohistochemistry refines the histopathologic prognosis of renal cell carcinoma. *Int. J. Clin. Exp. Pathol.* **2015**, *8*, 8165–8177. [[PubMed](#)]
50. Tang, J.Y.; Aszterbaum, M.; Athar, M.; Barsanti, F.; Cappola, C.; Estevez, N.; Hebert, J.; Hwang, J.; Khaimskiy, Y.; Kim, A.; et al. Basal cell carcinoma chemoprevention with nonsteroidal anti-inflammatory drugs in genetically predisposed PTCH1+/- humans and mice. *Cancer Prev. Res.* **2010**, *3*, 25–34. [[CrossRef](#)] [[PubMed](#)]
51. Müller-Decker, K. Cyclooxygenase-dependent signaling is causally linked to non-melanoma skin carcinogenesis: Pharmacological, genetic, and clinical evidence. *Cancer Metastasis Rev.* **2011**, *30*, 343–361. [[CrossRef](#)] [[PubMed](#)]
52. Erovic, B.M.; Woegerbauer, M.; Pammer, J.; Selzer, E.; Grasl, M.C.; Thurnher, D. Strong evidence for up-regulation of cyclooxygenase-1 in head and neck cancer. *Eur. J. Clin. Investig.* **2008**, *38*, 61–66. [[CrossRef](#)] [[PubMed](#)]
53. Mauro, A.; Lipari, L.; Leone, A.; Tortorici, S.; Burruano, F.; Provenzano, S.; Gerbino, A.; Buscemi, M. Expression of cyclooxygenase-1 and cyclooxygenase-2 in normal and pathological human oral mucosa. *Folia Histochem. Cytobiol.* **2010**, *48*, 555–563. [[CrossRef](#)] [[PubMed](#)]
54. Pannone, G.; Sanguedolce, F.; De Maria, S.; Farina, E.; Lo Muzio, L.; Serpico, R.; Emanuelli, M.; Rubini, C.; De Rosa, G.; Staibano, S.; et al. Cyclooxygenase isozymes in oral squamous cell carcinoma: A real-time RT-PCR study with clinic pathological correlations. *Int. J. Immunopathol. Pharmacol.* **2007**, *20*, 317–324. [[CrossRef](#)] [[PubMed](#)]
55. Erovic, B.M.; Al Habeeb, A.; Harris, L.; Goldstein, D.P.; Kim, D.; Ghazarian, D.; Irish, J.C. Identification of novel target proteins in sebaceous gland carcinoma. *Head Neck* **2013**, *35*, 642–648. [[CrossRef](#)] [[PubMed](#)]
56. Park, S.W.; Kim, H.S.; Hah, J.H.; Kim, K.H.; Heo, D.S.; Sung, M.W. Differential effects between cyclooxygenase-2 inhibitors and siRNA on vascular endothelial growth factor production in head and neck squamous cell carcinoma cell lines. *Head Neck* **2010**, *32*, 1534–1543. [[CrossRef](#)] [[PubMed](#)]
57. Park, S.W.; Kim, H.S.; Choi, M.S.; Kim, J.E.; Jeong, W.J.; Heo, D.S.; Sung, M.W. The influence of cyclooxygenase-1 expression on the efficacy of cyclooxygenase-2 inhibition in head and neck squamous cell carcinoma cell lines. *Anticancer Drugs* **2011**, *22*, 416–423. [[CrossRef](#)] [[PubMed](#)]

58. Akiyama, J.; Alexandre, L.; Baruah, A.; Buttar, N.; Chandra, R.; Clark, A.B.; Hart, A.R.; Hawk, E.; Kandioler, D.; Kappel, S.; et al. Strategy for prevention of cancers of the esophagus. *Ann. N. Y. Acad. Sci.* **2014**, *1325*, 108–126. [[CrossRef](#)] [[PubMed](#)]
59. Tsibouris, P.; Vlachou, E.; Isaacs, P.E. Role of chemoprophylaxis with either NSAIDs or statins in patients with Barrett's esophagus. *World J. Gastrointest. Pharmacol. Ther.* **2014**, *5*, 27–39. [[CrossRef](#)] [[PubMed](#)]
60. Morris, C.D.; Armstrong, G.R.; Bigley, G.; Green, H.; Attwood, S.E. Cyclooxygenase-2 expression in the Barrett's metaplasia–dysplasia–adenocarcinoma sequence. *Am. J. Gastroenterol.* **2001**, *96*, 990–996. [[CrossRef](#)] [[PubMed](#)]
61. Taddei, A.; Fabbroni, V.; Pini, A.; Lucarini, L.; Ringressi, M.N.; Fantappiè, O.; Bani, D.; Messerini, L.; Masini, E.; Bechi, P. Cyclooxygenase-2 and inflammation mediators have a crucial role in reflux-related esophageal histological changes and Barrett's esophagus. *Dig. Dis. Sci.* **2014**, *59*, 949–957. [[CrossRef](#)] [[PubMed](#)]
62. Von Rahden, B.H.; Stein, H.J.; Pühringer, F.; Koch, I.; Langer, R.; Piontek, G.; Siewert, J.R.; Höfler, H.; Sarbia, M. Coexpression of cyclooxygenases (COX-1, COX-2) and vascular endothelial growth factors (VEGF-A, VEGF-C) in esophageal adenocarcinoma. *Cancer Res.* **2005**, *65*, 5038–5044. [[CrossRef](#)] [[PubMed](#)]
63. Piazzuelo, E.; Santander, S.; Cebrián, C.; Jiménez, P.; Pastor, C.; García-González, M.A.; Esteva, F.; Esquivias, P.; Ortego, J.; Lanás, A. Characterization of the prostaglandin E2 pathway in a rat model of esophageal adenocarcinoma. *Curr. Cancer Drug Targets* **2012**, *12*, 132–143. [[CrossRef](#)] [[PubMed](#)]
64. Esquivias, P.; Morandeira, A.; Escartín, A.; Cebrián, C.; Santander, S.; Esteva, F.; García-González, M.A.; Ortego, J.; Lanás, A.; Piazzuelo, E. Indomethacin but not a selective cyclooxygenase-2 inhibitor inhibits esophageal adenocarcinogenesis in rats. *World J. Gastroenterol.* **2012**, *18*, 4866–4874. [[CrossRef](#)] [[PubMed](#)]
65. Zhang, S.; Zhang, X.Q.; Ding, X.W.; Yang, R.K.; Huang, S.L.; Kastelein, F.; Bruno, M.; Yu, X.J.; Zhou, D.; Zou, X.P. Cyclooxygenase inhibitors use is associated with reduced risk of esophageal adenocarcinoma in patients with Barrett's esophagus: A meta-analysis. *Br. J. Cancer* **2014**, *110*, 2378–2388. [[CrossRef](#)] [[PubMed](#)]
66. Janakiram, N.B.; Rao, C.V. The role of inflammation in colon cancer. *Adv. Exp. Med. Biol.* **2014**, *816*, 25–52. [[CrossRef](#)] [[PubMed](#)]
67. Arber, N.; Eagle, C.J.; Spicak, J.; Rácz, I.; Dite, P.; Hajer, J.; Zavoral, M.; Lechuga, M.J.; Gerletti, P.; Tang, J.; et al. PreSAP Trial Investigators. Celecoxib for the prevention of colorectal adenomatous polyps. *N. Engl. J. Med.* **2006**, *355*, 885–895. [[CrossRef](#)] [[PubMed](#)]
68. Kraus, S.; Naumov, I.; Arber, N. COX-2 active agents in the chemoprevention of colorectal cancer. *Recent Results Cancer Res.* **2013**, *191*, 95–103. [[CrossRef](#)] [[PubMed](#)]
69. Church, R.D.; Yu, J.; Fleshman, J.W.; Shannon, W.D.; Govindan, R.; McLeod, H.L. RNA profiling of cyclooxygenases 1 and 2 in colorectal cancer. *Br. J. Cancer* **2004**, *91*, 1015–1018. [[CrossRef](#)] [[PubMed](#)]
70. Von Rahden, B.H.; Brücher, B.L.; Langner, C.; Siewert, J.R.; Stein, H.J.; Sarbia, M. Expression of cyclo-oxygenase 1 and 2, prostaglandin E synthase and transforming growth factor beta1, and their relationship with vascular endothelial growth factors A and C, in primary adenocarcinoma of the small intestine. *Br. J. Surg.* **2006**, *93*, 1424–1432. [[CrossRef](#)] [[PubMed](#)]
71. Riehl, T.E.; George, R.J.; Sturmoski, M.A.; May, R.; Dieckgraefe, B.; Anant, S.; Houchen, C.W. Azoxymethane protects intestinal stem cells and reduces crypt epithelial mitosis through a COX-1 dependent mechanism. *Am. J. Physiol. Gastrointest. Liver Physiol.* **2006**, *291*, G1062–G1070. [[CrossRef](#)] [[PubMed](#)]
72. Takeda, H.; Sonoshita, M.; Oshima, H.; Sugihara, K.; Chulada, P.C.; Langenbach, R.; Oshima, M.; Taketo, M.M. Cooperation of cyclooxygenase 1 and cyclooxygenase 2 in intestinal polyposis. *Cancer Res.* **2003**, *63*, 4872–4877. [[PubMed](#)]
73. Smartt, H.J.; Greenhough, A.; Ordóñez-Morán, P.; Talero, E.; Cherry, C.A.; Wallam, C.A.; Parry, L.; Al Kharusi, M.; Roberts, H.R.; Mariadason, J.M.; et al.  $\beta$ -catenin represses expression of the tumour suppressor 15-prostaglandin dehydrogenase in the normal intestinal epithelium and colorectal tumour cells. *Gut* **2012**, *61*, 1306–13014. [[CrossRef](#)] [[PubMed](#)]
74. Li, H.; Zhu, F.; Chen, H.; Cheng, K.W.; Zykova, T.; Oi, N.; Lubet, R.A.; Bode, A.M.; Wang, M.; Dong, Z. 6-C-(E-phenylethenyl)-naringenin suppresses colorectal cancer growth by inhibiting cyclooxygenase-1. *Cancer Res.* **2014**, *74*, 243–252. [[CrossRef](#)] [[PubMed](#)]
75. Vona-Davis, L.; Rose, D.P. The obesity-inflammation-eicosanoid axis in breast cancer. *J. Mammary Gland Biol. Neoplasia* **2013**, *18*, 291–307. [[CrossRef](#)] [[PubMed](#)]

76. Hoellen, F.; Kelling, K.; Dittmer, C.; Diedrich, K.; Friedrich, M.; Thill, M. Impact of cyclooxygenase-2 in breast cancer. *Anticancer Res.* **2011**, *31*, 4359–4367. [[PubMed](#)]
77. Hwang, D.; Scollard, D.; Byrne, J.; Levine, E. Expression of cyclooxygenase-1 and cyclooxygenase-2 in human breast cancer. *J. Natl. Cancer Inst.* **1998**, *90*, 455–460. [[CrossRef](#)] [[PubMed](#)]
78. Fahlén, M.; Zhang, H.; Löfgren, L.; Masironi, B.; von Schoultz, E.; von Schoultz, B.; Sahlin, L. Expression of cyclooxygenase-1 and cyclooxygenase-2, syndecan-1 and connective tissue growth factor in benign and malignant breast tissue from premenopausal women. *Gynecol. Endocrinol.* **2017**, *33*, 353–358. [[CrossRef](#)] [[PubMed](#)]
79. Haakensen, V.D.; Bjørø, T.; Lüders, T.; Riis, M.; Bukholm, I.K.; Kristensen, V.N.; Troester, M.A.; Homen, M.M.; Ursin, G.; Børresen-Dale, A.L.; et al. Serum estradiol levels associated with specific gene expression patterns in normal breast tissue and in breast carcinomas. *BMC Cancer* **2011**, *11*, 332. [[CrossRef](#)] [[PubMed](#)]
80. Jeong, H.S.; Kim, J.H.; Choi, H.Y.; Lee, E.R.; Cho, S.G. Induction of cell growth arrest and apoptotic cell death in human breast cancer MCF-7 cells by the COX-1 inhibitor FR122047. *Oncol. Rep.* **2010**, *24*, 351–356. [[CrossRef](#)] [[PubMed](#)]
81. McFadden, D.W.; Riggs, D.R.; Jackson, B.J.; Cunningham, C. Additive effects of Cox-1 and Cox-2 inhibition on breast cancer in vitro. *Int. J. Oncol.* **2006**, *29*, 1019–1023. [[CrossRef](#)] [[PubMed](#)]
82. Kundu, N.; Fulton, A.M. Selective cyclooxygenase (COX)-1 or COX-2 inhibitors control metastatic disease in a murine model of breast cancer. *Cancer Res.* **2002**, *62*, 2343–2346. [[PubMed](#)]
83. Boccardo, E.; Lepique, A.P.; Villa, L.L. The role of inflammation in HPV carcinogenesis. *Carcinogenesis* **2010**, *31*, 1905–1912. [[CrossRef](#)] [[PubMed](#)]
84. Subbaramaiah, K.; Dannenberg, A.J. Cyclooxygenase-2 transcription is regulated by human papillomavirus 16 E6 and E7 oncoproteins: Evidence of a corepressor/coactivator exchange. *Cancer Res.* **2007**, *67*, 3976–3985. [[CrossRef](#)] [[PubMed](#)]
85. Young, J.L.; Jazaeri, A.A.; Darus, C.J.; Modesitt, S.C. Cyclooxygenase-2 in cervical neoplasia: A review. *Gynecol. Oncol.* **2008**, *109*, 140–145. [[CrossRef](#)] [[PubMed](#)]
86. Grabosch, S.M.; Shariff, O.M.; Wulff, J.L.; Helm, C.W. Non-steroidal anti-inflammatory agents to induce regression and prevent the progression of cervical intraepithelial neoplasia. *Cochrane Database Syst. Rev.* **2014**, CD004121. [[CrossRef](#)] [[PubMed](#)]
87. Sales, K.J.; Katz, A.A.; Howard, B.; Soeters, R.P.; Millar, R.P.; Jabbour, H.N. Cyclooxygenase 1 is up-regulated in cervical carcinomas: Autocrine/paracrine regulation of cyclooxygenase-2, prostaglandin E receptors and angiogenic factors by Cyclooxygenase-1. *Cancer Res.* **2002**, *62*, 424–432. [[PubMed](#)]
88. Sutherland, J.R.; Sales, K.J.; Jabbour, H.N.; Katz, A.A. Seminal plasma enhances cervical adenocarcinoma cell proliferation and tumour growth in vivo. *PLoS ONE* **2012**, *7*, e33848. [[CrossRef](#)] [[PubMed](#)]
89. Sales, K.J.; Sutherland, J.R.; Jabbour, H.N.; Katz, A.A. Seminal plasma induces angiogenic chemokine expression in cervical cancer cells and regulates vascular function. *Biochim. Biophys. Acta* **2012**, *1823*, 1789–1795. [[CrossRef](#)] [[PubMed](#)]
90. Radilova, H.; Libra, A.; Holasova, S.; Safarova, M.; Viskova, A.; Kunc, F.; Buncek, M. COX-1 is coupled with mPGES-1 and ABCC4 in human cervix cancer cells. *Mol. Cell. Biochem.* **2009**, *330*, 131–140. [[CrossRef](#)] [[PubMed](#)]
91. Jung, Y.W.; Kim, S.W.; Kim, S.; Kim, J.H.; Cho, N.H.; Kim, J.W.; Kim, Y.T. Prevalence and clinical relevance of cyclooxygenase-1 and -2 expression in stage IIB cervical adenocarcinoma. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **2010**, *148*, 62–66. [[CrossRef](#)] [[PubMed](#)]
92. Athavale, R.; Clooney, K.; O'Hagan, J.; Shawki, H.; Clark, A.H.; Green, J.A. COX-1 and COX-2 expression in stage I and II invasive cervical carcinoma: Relationship to disease relapse and long-term survival. *Int. J. Gynecol. Cancer* **2006**, *16*, 1303–1308. [[CrossRef](#)] [[PubMed](#)]
93. Jeon, Y.T.; Song, Y.C.; Kim, S.H.; Wu, H.G.; Kim, I.H.; Park, I.A.; Kim, J.W.; Park, N.H.; Kang, S.B.; Lee, H.P.; et al. Influences of cyclooxygenase-1 and -2 expression on the radiosensitivities of human cervical cancer cell lines. *Cancer Lett.* **2007**, *256*, 33–38. [[CrossRef](#)] [[PubMed](#)]
94. Kim, H.S.; Kim, T.; Kim, M.K.; Suh, D.H.; Chung, H.H.; Song, Y.S. Cyclooxygenase-1 and -2: Molecular targets for cervical neoplasia. *J. Cancer Prev.* **2013**, *18*, 123–134. [[CrossRef](#)] [[PubMed](#)]
95. Ohno, S.; Ohno, Y.; Suzuki, N.; Inagawa, H.; Kohchi, C.; Soma, G.; Inoue, M. Multiple roles of cyclooxygenase-2 in endometrial cancer. *Anticancer Res.* **2005**, *25*, 3679–3687. [[PubMed](#)]

96. Nasir, A.; Boulware, D.; Kaiser, H.E.; Lancaster, J.M.; Coppola, D.; Smith, P.V.; Hakam, A.; Siegel, S.E.; Bodey, B. Cyclooxygenase-2 (COX-2) expression in human endometrial carcinoma and precursor lesions and its possible use in cancer chemoprevention and therapy. *In Vivo* **2007**, *21*, 35–43. [[PubMed](#)]
97. Hasegawa, K.; Torii, Y.; Ishii, R.; Oe, S.; Kato, R.; Udagawa, Y. Effects of a selective COX-2 inhibitor in patients with uterine endometrial cancers. *Arch. Gynecol. Obstet.* **2011**, *284*, 1515–1521. [[CrossRef](#)] [[PubMed](#)]
98. Sugimoto, T.; Koizumi, T.; Sudo, T.; Yamaguchi, S.; Kojima, A.; Kumagai, S.; Nishimura, R. Correlative expression of cyclooxygenase-1 (Cox-1) and human epidermal growth factor receptor type-2 (Her-2) in endometrial cancer. *Kobe J. Med. Sci.* **2007**, *53*, 177–187. [[PubMed](#)]
99. Déry, M.C.; Chaudhry, P.; Leblanc, V.; Parent, S.; Fortier, A.M.; Asselin, E. Oxytocin increases invasive properties of endometrial cancer cells through phosphatidylinositol 3-kinase/AKT-dependent up-regulation of cyclooxygenase-1, -2, and X-linked inhibitor of apoptosis protein. *Biol. Reprod.* **2011**, *85*, 1133–1142. [[CrossRef](#)] [[PubMed](#)]
100. Vaughan, S.; Coward, J.I.; Bast, R.C., Jr.; Berchuck, A.; Berek, J.S.; Brenton, J.D.; Coukos, G.; Crum, C.C.; Drapkin, R.; Etemadmoghadam, D.; et al. Rethinking ovarian cancer: Recommendations for improving outcomes. *Nat. Rev. Cancer* **2011**, *11*, 719–725. [[CrossRef](#)] [[PubMed](#)]
101. Shan, W.; Liu, J. Inflammation, a hidden path to breaking the spell of ovarian cancer. *Cell Cycle* **2009**, *8*, 3107–3111. [[CrossRef](#)] [[PubMed](#)]
102. Macciò, A.; Madeddu, C. Inflammation and ovarian cancer. *Cytokine* **2012**, *58*, 133–147. [[CrossRef](#)] [[PubMed](#)]
103. Lee, G.; Ng, H.T. Clinical evaluations of a new ovarian cancer marker, COX-1. *Int. J. Gynaecol. Obstet.* **1995**, *49*, S27–S32. [[CrossRef](#)]
104. Gupta, R.A.; Tejada, L.V.; Tong, B.J.; Das, S.K.; Morrow, J.D.; Dey, S.K.; DuBois, R.N. Cyclooxygenase-1 is overexpressed and promotes angiogenic growth factor production in ovarian cancer. *Cancer Res.* **2003**, *63*, 906–911. [[PubMed](#)]
105. Daikoku, T.; Wang, D.; Tranguch, S.; Morrow, J.D.; Orsulic, S.; DuBois, R.N.; Dey, S.K. Cyclooxygenase-1 is a potential target for prevention and treatment of ovarian epithelial cancer. *Cancer Res.* **2005**, *65*, 3735–3744. [[CrossRef](#)] [[PubMed](#)]
106. Daikoku, T.; Tranguch, S.; Trofimova, I.N.; Dinulescu, D.M.; Jacks, T.; Nikitin, A.Y.; Connolly, D.C.; Dey, S.K. Cyclooxygenase-1 is overexpressed in multiple genetically engineered mouse models of epithelial ovarian cancer. *Cancer Res.* **2006**, *66*, 2527–2531. [[CrossRef](#)] [[PubMed](#)]
107. Hales, D.B.; Zhuge, Y.; Lagman, J.A.; Ansenberger, K.; Mahon, C.; Barua, A.; Luborsky, J.L.; Bahr, J.M. Cyclooxygenases expression and distribution in the normal ovary and their role in ovarian cancer in the domestic hen (*Gallus domesticus*). *Endocrine* **2008**, *33*, 235–244. [[CrossRef](#)] [[PubMed](#)]
108. Urick, M.E.; Johnson, P.A. Cyclooxygenase 1 and 2 mRNA and protein expression in the *Gallus domesticus* model of ovarian cancer. *Gynecol. Oncol.* **2006**, *103*, 673–678. [[CrossRef](#)] [[PubMed](#)]
109. Eilati, E.; Pan, L.; Bahr, J.M.; Hales, D.B. Age dependent increase in prostaglandin pathway coincides with onset of ovarian cancer in laying hens. *Prostaglandins Leukot. Essent. Fat. Acids* **2012**, *87*, 177–184. [[CrossRef](#)] [[PubMed](#)]
110. Kino, Y.; Kojima, F.; Kiguchi, K.; Igarashi, R.; Ishizuka, B.; Kawai, S. Prostaglandin E2 production in ovarian cancer cell lines is regulated by cyclooxygenase-1, not cyclooxygenase-2. *Prostaglandins Leukot. Essent. Fat. Acids* **2005**, *73*, 103–111. [[CrossRef](#)] [[PubMed](#)]
111. Lau, M.T.; Wong, A.S.; Leung, P.C. Gonadotropins induce tumor cell migration and invasion by increasing cyclooxygenases expression and prostaglandin E(2) production in human ovarian cancer cells. *Endocrinology* **2010**, *151*, 2985–2993. [[CrossRef](#)] [[PubMed](#)]
112. Ali-Fehmi, R.; Semaan, A.; Sethi, S.; Arabi, H.; Bandyopadhyay, S.; Hussein, Y.R.; Diamond, M.P.; Saed, G.; Morris, R.T.; Munkarah, A.R. Molecular typing of epithelial ovarian carcinomas using inflammatory markers. *Cancer* **2011**, *117*, 301–309. [[CrossRef](#)] [[PubMed](#)]
113. Liu, M.; Matsumura, N.; Mandai, M.; Li, K.; Yagi, H.; Baba, T.; Suzuki, A.; Hamanishi, J.; Fukuhara, K.; Konishi, I. Classification using hierarchical clustering of tumor-infiltrating immune cells identifies poor prognostic ovarian cancers with high levels of COX expression. *Mod. Pathol.* **2009**, *22*, 373–384. [[CrossRef](#)] [[PubMed](#)]
114. Hamanishi, J.; Mandai, M.; Abiko, K.; Matsumura, N.; Baba, T.; Yoshioka, Y.; Kosaka, K.; Konishi, I. The comprehensive assessment of local immune status of ovarian cancer by the clustering of multiple immune factors. *Clin. Immunol.* **2011**, *141*, 338–347. [[CrossRef](#)] [[PubMed](#)]



115. Lee, J.Y.; Myung, S.K.; Song, Y.S. Prognostic role of cyclooxygenase-2 in epithelial ovarian cancer: A meta-analysis of observational studies. *Gynecol. Oncol.* **2013**, *129*, 613–619. [[CrossRef](#)] [[PubMed](#)]
116. Kim, H.J.; Yim, G.W.; Nam, E.J.; Kim, Y.T. Synergistic effect of COX-2 inhibitor on paclitaxel-induced apoptosis in the human ovarian cancer cell line OVCAR-3. *Cancer Res. Treat.* **2014**, *46*, 81–92. [[CrossRef](#)] [[PubMed](#)]
117. Guo, F.J.; Tian, J.Y.; Jin, Y.M.; Wang, L.; Yang, R.Q.; Cui, M.H. Effects of cyclooxygenase-2 gene silencing on the biological behavior of SKOV3 ovarian cancer cells. *Mol. Med. Rep.* **2015**, *11*, 59–66. [[CrossRef](#)] [[PubMed](#)]
118. Wilson, A.J.; Fadare, O.; Beeghly-Fadiel, A.; Son, D.S.; Liu, Q.; Zhao, S.; Saskowski, J.; Uddin, M.J.; Daniel, C.; Crews, B.; et al. Aberrant over-expression of COX-1 intersects multiple pro-tumorigenic pathways in high-grade serous ovarian cancer. *Oncotarget* **2015**, *6*, 21353–21368. [[CrossRef](#)] [[PubMed](#)]
119. Daikoku, T.; Tranguch, S.; Chakrabarty, A.; Wang, D.; Khabele, D.; Orsulic, S.; Morrow, J.D.; Dubois, R.N.; Dey, S.K. Extracellular signal-regulated kinase is a target of cyclooxygenase-1-peroxisome proliferator-activated receptor-delta signaling in epithelial ovarian cancer. *Cancer Res.* **2007**, *67*, 5285–5292. [[CrossRef](#)] [[PubMed](#)]
120. Sonnemann, J.; Hüls, I.; Sigler, M.; Palani, C.D.; Hong, L.T.T.; Völker, U.; Kroemer, H.K.; Beck, J.F. Histone deacetylase inhibitors and aspirin interact synergistically to induce cell death in ovarian cancer cells. *Oncol. Rep.* **2008**, *20*, 219–224. [[CrossRef](#)] [[PubMed](#)]
121. Cho, M.; Kabir, S.M.; Dong, Y.; Lee, E.; Rice, V.M.; Khabele, D.; Son, D.S. Aspirin blocks EGF-stimulated cell viability in a COX-1 dependent manner in ovarian cancer cells. *J. Cancer* **2013**, *4*, 671–678. [[CrossRef](#)] [[PubMed](#)]
122. Son, D.S.; Wilson, A.J.; Parl, A.K.; Khabele, D. The effects of the histone deacetylase inhibitor romidepsin (FK228) are enhanced by aspirin (ASA) in COX-1 positive ovarian cancer cells through augmentation of p21. *Cancer Biol. Ther.* **2010**, *9*, 928–935. [[CrossRef](#)] [[PubMed](#)]
123. Truffinet, V.; Donnard, M.; Vincent, C.; Faucher, J.L.; Bordessoule, D.; Turlure, P.; Trimoreau, F.; Denizot, Y. Cyclooxygenase-1, but not -2, in blast cells of patients with acute leukemia. *Int. J. Cancer* **2007**, *121*, 924–927. [[CrossRef](#)] [[PubMed](#)]
124. Fiancette, R.; Vincent-Fabert, C.; Guerin, E.; Trimoreau, F.; Denizot, Y. Lipid mediators and human leukemic blasts. *J. Oncol.* **2011**, *2011*, 389021. [[CrossRef](#)] [[PubMed](#)]
125. McCulloch, D.; Brown, C.; Iland, H. Retinoic acid and arsenic trioxide in the treatment of acute promyelocytic leukemia: Current perspectives. *Onco Targets Ther.* **2017**, *10*, 1585–1601. [[CrossRef](#)] [[PubMed](#)]
126. Zhou, G.B.; Li, G.; Chen, S.J.; Chen, Z. From dissection of disease pathogenesis to elucidation of mechanisms of targeted therapies: Leukemia research in the genomic era. *Acta Pharmacol. Sin.* **2007**, *28*, 1434–1449. [[CrossRef](#)] [[PubMed](#)]
127. Rocca, B.; Morosetti, R.; Habib, A.; Maggiano, N.; Zassadowski, F.; Ciabattini, G.; Chomienne, C.; Papp, B.; Ranelletti, F.O. Cyclooxygenase-1, but not -2, is upregulated in NB4 leukemic cells and human primary promyelocytic blasts during differentiation. *Leukemia* **2004**, *18*, 1373–1379. [[CrossRef](#)] [[PubMed](#)]
128. Habib, A.; Hamade, E.; Mahfouz, R.; Nasrallah, M.S.; de Thé, H.; Bazarbachi, A. Arsenic trioxide inhibits ATRA-induced prostaglandin E2 and cyclooxygenase-1 in NB4 cells, a model of acute promyelocytic leukemia. *Leukemia* **2008**, *22*, 1125–1130. [[CrossRef](#)] [[PubMed](#)]
129. Lau, E.Y.; Ho, N.P.; Lee, T.K. Cancer stem cells and their microenvironment: Biology and therapeutic implications. *Stem Cells Int.* **2017**, *2017*, 3714190. [[CrossRef](#)] [[PubMed](#)]
130. Peitzsch, C.; Tyutyunnykova, A.; Pantel, K.; Dubrovska, A. Cancer stem cells: The root of tumor recurrence and metastases. *Semin. Cancer Biol.* **2017**, *44*, 10–24. [[CrossRef](#)] [[PubMed](#)]
131. Pang, L.Y.; Hurst, E.A.; Argyle, D.J. Cyclooxygenase-2: A role in cancer stem cell survival and repopulation of cancer cells during therapy. *Stem Cells Int.* **2016**, *2016*, 2048731. [[CrossRef](#)] [[PubMed](#)]
132. Goessling, W.; North, T.E.; Loewer, S.; Lord, A.M.; Lee, S.; Stoick-Cooper, C.L.; Weidinger, G.; Puder, M.; Daley, G.Q.; Moon, R.T.; et al. Genetic interaction of PGE2 and Wnt signaling regulates developmental specification of stem cells and regeneration. *Cell* **2009**, *136*, 1136–1147. [[CrossRef](#)] [[PubMed](#)]
133. Kanojia, D.; Zhou, W.; Zhang, J.; Jie, C.; Lo, P.K.; Wang, Q.; Chen, H. Proteomic profiling of cancer stem cells derived from primary tumors of HER2/Neu transgenic mice. *Proteomics* **2012**, *12*, 3407–3415. [[CrossRef](#)] [[PubMed](#)]
134. Lazennec, G.; Lam, P.Y. Recent discoveries concerning the tumor—Mesenchymal stem cell interactions. *Biochim. Biophys. Acta* **2016**, *1866*, 290–299. [[CrossRef](#)] [[PubMed](#)]

135. Roodhart, J.M.; Daenen, L.G.; Stigter, E.C.; Prins, H.J.; Gerrits, J.; Houthuijzen, J.M.; Gerritsen, M.G.; Schipper, H.S.; Backer, M.J.; van Amersfoort, M.; et al. Mesenchymal stem cells induce resistance to chemotherapy through the release of platinum-induced fatty acids. *Cancer Cell* **2011**, *20*, 370–383. [[CrossRef](#)] [[PubMed](#)]
136. Loftin, C.D.; Tiano, H.F.; Langenbanch, R. Phenotypes of the COX-deficient mice indicate physiological and pathophysiological roles for COX-1 and COX-2. *Prostaglandins Other Lipid Mediat.* **2002**, *68–69*, 177–185. [[CrossRef](#)]
137. Islam, A.B.; Dave, M.; Amin, S.; Jensen, R.V.; Amin, A.R. Genomic, lipidomic and metabolomic analysis of cyclooxygenase-null cells: Eicosanoid storm, cross talk, and compensation by COX-1. *Genom. Proteom. Bioinform.* **2016**, *14*, 81–93. [[CrossRef](#)] [[PubMed](#)]
138. Tortorella, M.D.; Zhang, Y.; Talley, J. Desirable Properties for 3rd Generation Cyclooxygenase-2 Inhibitors. *Mini Rev. Med. Chem.* **2016**, *16*, 1284–1289. [[CrossRef](#)] [[PubMed](#)]
139. Vitale, P.; Panella, A.; Scilimati, A.; Perrone, M.G. COX-1 inhibitors: Beyond structure toward therapy. *Med. Res. Rev.* **2016**, *36*, 641–671. [[CrossRef](#)] [[PubMed](#)]
140. Smith, C.J.; Zhang, Y.; Koboldt, C.M.; Muhammad, J.; Zweifel, B.S.; Shaffer, A.; Talley, J.J.; Masferrer, J.L.; Seibert, K.; Isakson, P.C. Pharmacological analysis of cyclooxygenase-1 in inflammation. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 13313–13318. [[CrossRef](#)] [[PubMed](#)]
141. Li, W.; Xu, R.J.; Lin, Z.Y.; Zhuo, G.C.; Zhang, H.H. Effects of a cyclooxygenase-1-selective inhibitor in a mouse model of ovarian cancer, administered alone or in combination with ibuprofen, a nonselective cyclooxygenase inhibitor. *Med. Oncol.* **2009**, *26*, 170–177. [[CrossRef](#)] [[PubMed](#)]
142. Li, W.; Ji, Z.L.; Zhuo, G.C.; Xu, R.J.; Wang, J.; Jiang, H.R. Effects of a selective cyclooxygenase-1 inhibitor in SKOV-3 ovarian carcinoma xenograft-bearing mice. *Med. Oncol.* **2010**, *27*, 98–104. [[CrossRef](#)] [[PubMed](#)]
143. Li, W.; Wang, J.; Jiang, H.R.; Xu, X.L.; Zhang, J.; Liu, M.L.; Zhai, L.Y. Combined effects of cyclooxygenase-1 and cyclooxygenase-2 selective inhibitors on ovarian carcinoma in vivo. *Int. J. Mol. Sci.* **2011**, *12*, 668–681. [[CrossRef](#)] [[PubMed](#)]
144. Li, W.; Liu, M.L.; Cai, J.H.; Tang, Y.X.; Zhai, L.Y.; Zhang, J. Effect of the combination of a cyclooxygenase-1 selective inhibitor and taxol on proliferation, apoptosis and angiogenesis of ovarian cancer in vivo. *Oncol. Lett.* **2012**, *4*, 168–174. [[CrossRef](#)] [[PubMed](#)]
145. Li, W.; Cai, J.H.; Zhang, J.; Tang, Y.X.; Wan, L. Effects of cyclooxygenase inhibitors in combination with taxol on expression of cyclin D1 and Ki-67 in a xenograft model of ovarian carcinoma. *Int. J. Mol. Sci.* **2012**, *13*, 9741–9753. [[CrossRef](#)] [[PubMed](#)]
146. Li, W.; Tang, Y.X.; Wan, L.; Cai, J.H.; Zhang, J. Effects of combining Taxol and cyclooxygenase inhibitors on the angiogenesis and apoptosis in human ovarian cancer xenografts. *Oncol. Lett.* **2013**, *5*, 923–928. [[CrossRef](#)] [[PubMed](#)]
147. Li, W.; Wan, L.; Zhai, L.Y.; Wang, J. Effects of SC-560 in combination with cisplatin or taxol on angiogenesis in human ovarian cancer xenografts. *Int. J. Mol. Sci.* **2014**, *5*, 19265–19280. [[CrossRef](#)] [[PubMed](#)]
148. Brenneis, C.; Maier, T.J.; Schmidt, R.; Hofacker, A.; Zulauf, L.; Jakobsson, P.J.; Scholich, K.; Geisslinger, G. Inhibition of prostaglandin E2 synthesis by SC-560 is independent of cyclooxygenase 1 inhibition. *FASEB J.* **2006**, *20*, 1352–1360. [[CrossRef](#)] [[PubMed](#)]
149. Saed, G.M. Immunohistochemical staining of cyclooxygenases with monoclonal antibodies. *Methods Mol. Biol.* **2008**, *477*, 219–228. [[CrossRef](#)] [[PubMed](#)]
150. Lee, J.P.; Hahn, H.S.; Hwang, S.J.; Choi, J.Y.; Park, J.S.; Lee, I.H.; Kim, T.J. Selective cyclooxygenase inhibitors increase paclitaxel sensitivity in taxane-resistant ovarian cancer by suppressing P-glycoprotein expression. *J. Gynecol. Oncol.* **2013**, *24*, 273–279. [[CrossRef](#)] [[PubMed](#)]
151. Huang, Y.; Ju, B.; Tian, J.; Liu, F.; Yu, H.; Xiao, H.; Liu, X.; Liu, W.; Yao, Z.; Hao, Q. Ovarian cancer stem cell-specific gene expression profiling and targeted drug prescreening. *Oncol. Rep.* **2014**, *31*, 1235–1248. [[CrossRef](#)] [[PubMed](#)]
152. Lee, J.K.; Havaleshko, D.M.; Cho, H.; Weinstein, J.N.; Kaldjian, E.P.; Karpovich, J.; Grimshaw, A.; Theodorescu, D. A strategy for predicting the chemosensitivity of human cancer and its application to drug discovery. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 13086–13091. [[CrossRef](#)] [[PubMed](#)]
153. Grösch, S.; Tegeder, I.; Niederberger, E.; Bräutigam, L.; Geisslinger, G. COX-2 independent induction of cell cycle arrest and apoptosis in colon cancer cells by the selective COX-2 inhibitor celecoxib. *FASEB J.* **2001**, *15*, 2742–2744. [[CrossRef](#)] [[PubMed](#)]

154. Wu, W.K.; Sung, J.J.; Wu, Y.C.; Li, H.T.; Yu, L.; Li, Z.J.; Cho, C.H. Inhibition of cyclooxygenase-1 lowers proliferation and induces macroautophagy in colon cancer cells. *Biochem. Biophys. Res. Commun.* **2009**, *382*, 79–84. [[CrossRef](#)] [[PubMed](#)]
155. Sheng, H.; Shao, J.; Kirkland, S.C.; Isakson, P.; Coffey, R.J.; Morrow, J.; Beauchamp, R.D.; DuBois, R.N. Inhibition of human colon cancer cell growth by selective inhibition of cyclooxygenase-2. *J. Clin. Investig.* **1997**, *99*, 2254–2259. [[CrossRef](#)] [[PubMed](#)]
156. Lee, E.; Choi, M.K.; Han, I.O.; Lim, S.J. Role of p21CIP1 as a determinant of SC560 response in human HCT116 colon carcinoma cells. *Exp. Mol. Med.* **2006**, *38*, 325–331. [[CrossRef](#)] [[PubMed](#)]
157. Sakoguchi-Okada, N.; Takahashi-Yanaga, F.; Fukada, K.; Shiraishi, F.; Taba, Y.; Miwa, Y.; Morimoto, S.; Iida, M.; Sasaguri, T. Celecoxib inhibits the expression of survivin via the suppression of promoter activity in human colon cancer cells. *Biochem. Pharmacol.* **2007**, *73*, 1318–1329. [[CrossRef](#)] [[PubMed](#)]
158. Jain, A.K.; Moore, S.M.; Yamaguchi, K.; Eling, T.E.; Baek, S.J. Selective nonsteroidal anti-inflammatory drugs induce thymosin beta-4 and alter actin cytoskeletal organization in human colorectal cancer cells. *J. Pharmacol. Exp. Ther.* **2004**, *311*, 885–891. [[CrossRef](#)] [[PubMed](#)]
159. Moon, Y. NSAID-activated gene 1 and its implications for mucosal integrity and intervention beyond NSAIDs. *Pharmacol. Res.* **2017**, *121*, 122–128. [[CrossRef](#)] [[PubMed](#)]
160. Wang, W.S.; Chen, P.M.; Hsiao, H.L.; Wang, H.S.; Liang, W.Y.; Su, Y. Overexpression of the thymosin beta-4 gene is associated with increased invasion of SW480 colon carcinoma cells and the distant metastasis of human colorectal carcinoma. *Oncogene* **2004**, *23*, 6666–6671. [[CrossRef](#)] [[PubMed](#)]
161. Piao, Z.; Hong, C.S.; Jung, M.R.; Choi, C.; Park, Y.K. Thymosin  $\beta$ 4 induces invasion and migration of human colorectal cancer cells through the ILK/AKT/ $\beta$ -catenin signaling pathway. *Biochem. Biophys. Res. Commun.* **2014**, *452*, 858–864. [[CrossRef](#)] [[PubMed](#)]
162. Gemoll, T.; Strohkamp, S.; Schillo, K.; Thorns, C.; Habermann, J.K. MALDI-imaging reveals thymosin beta-4 as an independent prognostic marker for colorectal cancer. *Oncotarget* **2015**, *6*, 43869–43880. [[CrossRef](#)] [[PubMed](#)]
163. Lee, E.; Choi, M.K.; Youk, H.J.; Kim, C.H.; Han, I.O.; Yoo, B.C.; Lee, M.K.; Lim, S.J. 5-(4-chlorophenyl)-1-(4-methoxyphenyl)-3-trifluoromethylpyrazole acts in a reactive oxygen species-dependent manner to suppress human lung cancer growth. *J. Cancer Res. Clin. Oncol.* **2006**, *132*, 223–233. [[CrossRef](#)] [[PubMed](#)]
164. Ding, J.; Tsuboi, K.; Hoshikawa, H.; Goto, R.; Mori, N.; Katsukawa, M.; Hiraki, E.; Yamamoto, S.; Abe, M.; Ueda, N. Cyclooxygenase isozymes are expressed in human myeloma cells but not involved in anti-proliferative effect of cyclooxygenase inhibitors. *Mol. Carcinog.* **2006**, *45*, 250–259. [[CrossRef](#)] [[PubMed](#)]
165. Lampiasi, N.; Foderà, D.; D’Alessandro, N.; Cusimano, A.; Azzolina, A.; Tripodo, C.; Florena, A.M.; Minervini, M.I.; Notarbartolo, M.; Montalto, G.; et al. The selective cyclooxygenase-1 inhibitor SC-560 suppresses cell proliferation and induces apoptosis in human hepatocellular carcinoma cells. *Int. J. Mol. Med.* **2006**, *17*, 245–252. [[CrossRef](#)] [[PubMed](#)]
166. Goto, K.; Ochi, H.; Yasunaga, Y.; Matsuyuki, H.; Imayoshi, T.; Kusuhara, H.; Okumoto, T. Analgesic effect of mofezolac, a non-steroidal anti-inflammatory drug, against phenylquinone-induced acute pain in mice. *Prostaglandins Other Lipid Mediat.* **1998**, *56*, 245–254. [[CrossRef](#)]
167. Kusuhara, H.; Matsuyuki, H.; Matsuura, M.; Imayoshi, T.; Okumoto, T.; Matsui, H. Induction of apoptotic DNA fragmentation by nonsteroidal anti-inflammatory drugs in cultured rat gastric mucosal cells. *Eur. J. Pharmacol.* **1998**, *360*, 273–280. [[CrossRef](#)]
168. Kusuhara, H.; Komatsu, H.; Sumichika, H.; Sugahara, K. Reactive oxygen species are involved in the apoptosis induced by nonsteroidal anti-inflammatory drugs in cultured gastric cells. *Eur. J. Pharmacol.* **1999**, *383*, 331–337. [[CrossRef](#)]
169. Kitamura, T.; Kawamori, T.; Uchiya, N.; Itoh, M.; Noda, T.; Matsuura, M.; Sugimura, T.; Wakabayashi, K. Inhibitory effects of mofezolac, a cyclooxygenase-1 selective inhibitor, on intestinal carcinogenesis. *Carcinogenesis* **2002**, *23*, 1463–1466. [[CrossRef](#)] [[PubMed](#)]
170. Kitamura, T.; Itoh, M.; Noda, T.; Matsuura, M.; Wakabayashi, K. Combined effects of cyclooxygenase-1 and cyclooxygenase-2 selective inhibitors on intestinal tumorigenesis in adenomatous polyposis coli gene knockout mice. *Int. J. Cancer* **2004**, *109*, 576–580. [[CrossRef](#)] [[PubMed](#)]

171. Niho, N.; Kitamura, T.; Takahashi, M.; Mutoh, M.; Sato, H.; Matsuura, M.; Sugimura, T.; Wakabayashi, K. Suppression of azoxymethane-induced colon cancer development in rats by a cyclooxygenase-1 selective inhibitor, mofezolac. *Cancer Sci.* **2006**, *97*, 1011–1014. [[CrossRef](#)] [[PubMed](#)]
172. Miao, L.; Shiraishi, R.; Fujise, T.; Kuroki, T.; Kakimoto, T.; Sakata, Y.; Takashima, T.; Iwakiri, R.; Fujimoto, K.; Shi, R.; et al. Chemopreventive effect of mofezolac on beef tallow diet/azoxymethane-induced colon carcinogenesis in rats. *Hepatogastroenterology* **2011**, *58*, 81–88. [[PubMed](#)]
173. Tanaka, A.; Sakai, H.; Motoyama, Y.; Ishikawa, T.; Takasugi, H. Antiplatelet agents based on cyclooxygenase inhibition without ulcerogenesis. Evaluation and synthesis of 4,5-bis(4-methoxyphenyl)-2-substituted-thiazoles. *J. Med. Chem.* **1994**, *37*, 1189–1199. [[CrossRef](#)] [[PubMed](#)]
174. Ochi, T.; Motoyama, Y.; Goto, T. The analgesic effect profile of FR122047, a selective cyclooxygenase-1 inhibitor, in chemical nociceptive models. *Eur. J. Pharmacol.* **2000**, *391*, 49–54. [[CrossRef](#)]
175. Ochi, T.; Goto, T. Differential effect of FR122047, a selective cyclo-oxygenase-1 inhibitor, in rat chronic models of arthritis. *Br. J. Pharmacol.* **2002**, *135*, 782–788. [[CrossRef](#)] [[PubMed](#)]
176. Ochi, T.; Ohkubo, Y.; Mutoh, S. Role of cyclooxygenase-2, but not cyclooxygenase-1, on type II collagen-induced arthritis in DBA/1J mice. *Biochem. Pharmacol.* **2003**, *66*, 1055–1060. [[CrossRef](#)]
177. Jeong, H.S.; Choi, H.Y.; Lee, E.R.; Kim, J.H.; Jeon, K.; Lee, H.J.; Cho, S.G. Involvement of caspase-9 in autophagy-mediated cell survival pathway. *Biochim. Biophys. Acta* **2011**, *1813*, 80–90. [[CrossRef](#)] [[PubMed](#)]
178. Boccarelli, A.; Pannunzio, A.; Coluccia, M. The Challenge of Establishing Reliable Screening Tests for Selecting Anticancer Metal Compounds. In *Bioinorganic Medicinal Chemistry*; Alessio, E., Ed.; Wiley-VCH Verlag GmbH & Co., KGaA: Weinheim, Germany, 2011; pp. 175–196. ISBN 978-3-527-32631-0.
179. Lin, J.H.; Cocchetto, D.M.; Duggan, D.E. Protein binding as a primary determinant of the clinical pharmacokinetic properties of non-steroidal anti-inflammatory drugs. *Clin. Pharmacokinet.* **1987**, *12*, 402–432. [[CrossRef](#)] [[PubMed](#)]
180. Duan, W.; Zhang, L. Cyclooxygenase inhibitors not inhibit resting lung cancer A549 cell proliferation. *Prostaglandins Leukot. Essent. Fat. Acids.* **2006**, *74*, 317–321. [[CrossRef](#)] [[PubMed](#)]
181. Cancer Cell Line Encyclopedia (CCLE). Available online: <http://www.broadinstitute.org/ccle> (accessed on 4 September 2018).
182. Domcke, S.; Sinha, R.; Levine, D.A.; Sander, C.; Schultz, N. Evaluating cell lines as tumour models by comparison of genomic profiles. *Nat. Commun.* **2013**, *4*, 2126. [[CrossRef](#)] [[PubMed](#)]
183. Bray, M.A.; Singh, S.; Han, H.; Davis, C.T.; Borgeson, B.; Hartland, C.; Kost-Alimova, M.; Gustafsdottir, S.M.; Gibson, C.C.; Carpenter, A.E. Cell Painting a high-content image-based assay for morphological profiling using multiplexed fluorescent dyes. *Nat. Protoc.* **2016**, *11*, 1757–1774. [[CrossRef](#)] [[PubMed](#)]
184. Grösch, S.; Maier, T.J.; Schiffmann, S.; Geisslinger, G. Cyclooxygenase-2 (COX-2)-independent anticarcinogenic effects of selective COX-2 inhibitors. *J. Natl. Cancer Inst.* **2006**, *98*, 736–747. [[CrossRef](#)] [[PubMed](#)]
185. National Center for Biotechnology Information. PubChem BioAssay Database, AID=624141. Available online: <https://pubchem.ncbi.nlm.nih.gov/bioassay/624141> (accessed on 4 September 2018).

