

Article

# The Fifth Coefficient of Bazilevič Functions

Oh Sang Kwon <sup>1</sup>, Derek Keith Thomas <sup>2</sup> and Young Jae Sim <sup>1,\*</sup> 

<sup>1</sup> Department of Mathematics, Kyungsung University, Busan 48434, Korea; oskwon@ks.ac.kr

<sup>2</sup> Department of Mathematics, Swansea University, Bay Campus, Swansea, SA1 8EN, UK; d.k.thomas@swansea.ac.uk

\* Correspondence: yjsim@ks.ac.kr

Received: 12 July 2020; Accepted: 25 July 2020; Published: 26 July 2020



**Abstract:** Let  $f \in \mathcal{A}$ , the class of normalized analytic functions defined in the unit disk  $\mathbb{D}$ , and be given by  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$  for  $z \in \mathbb{D}$ . This paper presents a new approach to finding bounds for  $|a_n|$ . As an application, we find the sharp bound for  $|a_5|$  for the class  $B_1(\alpha)$  of Bazilevič functions when  $\alpha \geq 1$ .

**Keywords:** univalent functions, Bazilevič functions, coefficient problems

**MSC:** 30C45; 30C50

## 1. Introduction

Let  $\mathcal{A}$  denote the class of analytic functions  $f$  in the unit disk  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$  normalized by  $f(0) = 0 = f'(0) - 1$ . Then for  $z \in \mathbb{D}$ ,  $f \in \mathcal{A}$  has the following representation

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n. \quad (1)$$

Denote by  $\mathcal{S}$ , the subset of  $\mathcal{A}$  consisting of univalent functions in  $\mathbb{D}$ .

We remark at the outset that in a great number of the more familiar subclasses of  $\mathcal{S}$ , sharp bounds have been found for the coefficients  $|a_n|$ , when  $2 \leq n \leq 4$ , but bounds when  $n = 5$  and beyond are much more difficult to obtain. (See, e.g., [1]).

Denote by  $\mathcal{S}^*$ , the class of starlike functions defined as follows.

**Definition 1.** Let  $f \in \mathcal{A}$ . Then  $f \in \mathcal{S}^*$  if, and only if, for  $z \in \mathbb{D}$ ,

$$\operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > 0.$$

An application of the method introduced in this paper to estimate the fifth coefficient of functions in  $\mathcal{A}$ , concerns the  $B_1(\alpha)$  Bazilevič functions defined as follows.

**Definition 2.** Let  $f \in \mathcal{A}$ . Then  $f \in B_1(\alpha)$  if, and only if, for  $\alpha \geq 0$ , and  $z \in \mathbb{D}$ ,

$$\operatorname{Re} \left\{ f'(z) \left( \frac{f(z)}{z} \right)^{\alpha-1} \right\} > 0. \quad (2)$$

We note that  $B_1(0) = \mathcal{S}^*$ , and each of the above classes are necessarily subclasses of  $\mathcal{S}$ . Apart from  $\alpha = 0$ , where  $|a_n| \leq n$  for  $n \geq 2$ , we also note that sharp bounds for  $|a_n|$  are known for  $f \in B_1(\alpha)$

when  $\alpha \geq 0$  only when  $2 \leq n \leq 4$ , [2], and only partial solutions are known for  $|a_n|$  when  $n \geq 5$  [3,4] for  $\alpha \geq 1$ .

It was conjectured in [4], that when  $\alpha \geq 1$ , the sharp bound for  $|a_n|$  when  $n \geq 2$  is given by

$$|a_n| \leq \frac{2}{n-1+\alpha'}$$

and a partial solution to this problem in the case  $n = 5$  was given in [3].

In this paper, we illustrate our method by giving a complete solution to finding the sharp bound for  $|a_5|$  when  $f \in \mathcal{B}_1(\alpha)$  for  $\alpha \geq 1$ .

### 2. Auxiliary Results

Denote by  $\mathcal{P}$ , the class of analytic functions  $p$  with positive real part on  $\mathbb{D}$  given by

$$p(z) = 1 + \sum_{n=1}^{\infty} p_n z^n. \tag{3}$$

**Lemma 1** ([5]). *If the functions*

$$1 + \sum_{n=1}^{\infty} b_n z^n \quad \text{and} \quad 1 + \sum_{n=1}^{\infty} c_n z^n$$

*belong to  $\mathcal{P}$ , then the same is true of the function*

$$1 + \frac{1}{2} \sum_{n=1}^{\infty} b_n c_n z^n.$$

**Lemma 2** ([6]). *Let  $h(z) = 1 + \beta_1 z + \beta_2 z^2 + \dots$  and  $1 + G(z) = 1 + d_1 z + d_2 z^2 + \dots$  be functions in  $\mathcal{P}$ ,  $\epsilon_0 = 1$ , and*

$$\epsilon_n = \frac{1}{2^n} \left[ 1 + \frac{1}{2} \sum_{k=1}^n \binom{n}{k} \beta_k \right], \quad n \in \mathbb{N}. \tag{4}$$

*If  $A_n$  ( $n \in \mathbb{N}$ ) is defined by*

$$\sum_{n=1}^{\infty} (-1)^{n+1} \epsilon_{n-1} G^n(z) = \sum_{n=1}^{\infty} A_n z^n,$$

*then  $|A_n| \leq 2$ .*

We first outline the method of proof.

Let  $p \in \mathcal{P}$  be in the form (3), and

$$\Psi = p_4 + B_1 p_1^4 + B_2 p_1^2 p_2 + B_3 p_1 p_3 + B_4 p_2^2 \tag{5}$$

with  $B_i \in \mathbb{C}$ ,  $i \in \{1, 2, 3, 4\}$ . Assume that there exists  $q \in \mathcal{P}$  of the form  $q(z) = 1 + \sum_{n=1}^{\infty} b_n z^n$ . Then by

Lemma 1 the function

$$1 + \frac{1}{2} (p(z) - 1) * (q(z) - 1) = 1 + \sum_{n=1}^{\infty} \frac{1}{2} b_n p_n z^n$$

also belongs to  $\mathcal{P}$ . Let

$$1 + G(z) = 1 + \frac{1}{2} (p(z) - 1) * (q(z) - 1) = 1 + \sum_{n=1}^{\infty} v_n z^n.$$

Then  $1 + G(z) \in \mathcal{P}$ , and  $v_n = b_n p_n / 2$ ,  $n \in \mathbb{N}$ .

Now assume that  $h(z) = 1 + \sum_{n=1}^{\infty} u_n z^n \in \mathcal{P}$ . Then by Lemma 2 we obtain  $|A_4| \leq 2$ , where

$$A_4 = \frac{1}{2}\epsilon_0 b_4 p_4 - \frac{1}{4}\epsilon_1 b_2^2 p_2^2 - \frac{1}{2}\epsilon_1 b_1 b_3 p_1 p_3 + \frac{3}{8}\epsilon_2 b_1^2 b_2 p_1^2 p_2 - \frac{1}{16}\epsilon_3 b_1^4 p_1^4. \tag{6}$$

Here,  $\epsilon_i, i \in \{1, 2, 3, 4\}$ , are given by

$$\begin{aligned} \epsilon_0 &= 1, \\ \epsilon_1 &= \frac{1}{2} \left( 1 + \frac{1}{2} u_1 \right), \\ \epsilon_2 &= \frac{1}{4} \left( 1 + u_1 + \frac{1}{2} u_2 \right), \\ \epsilon_3 &= \frac{1}{8} \left( 1 + \frac{3}{2} u_1 + \frac{3}{2} u_2 + \frac{1}{2} u_3 \right). \end{aligned}$$

Hence we have the following.

(A) Let  $p \in \mathcal{P}$  be in the form (3). If there exist  $q, h \in \mathcal{P}$  such that  $q$  and  $h$  are represented by

$$q(z) = 1 + \sum_{n=1}^{\infty} b_n z^n, \quad h(z) = 1 + \sum_{n=1}^{\infty} u_n z^n,$$

respectively, with

$$\begin{cases} b_4 = 2 \\ B_1 = -\frac{1}{128} \left( 1 + \frac{3}{2} u_1 + \frac{3}{2} u_2 + \frac{1}{2} u_3 \right) b_1^4 \\ B_2 = \frac{3}{32} \left( 1 + u_1 + \frac{1}{2} u_2 \right) b_1^2 b_2 \\ B_3 = -\frac{1}{4} \left( 1 + \frac{1}{2} u_1 \right) b_1 b_3 \\ B_4 = -\frac{1}{8} \left( 1 + \frac{1}{2} u_1 \right) b_2^2, \end{cases}$$

then  $|\Psi| = |A_4| \leq 2$ , where  $\Psi$  and  $A_4$  are given by (5) and (6), respectively.

We now recall a recent result of Cho et al. [7], where they obtained the following parametric formulas for the initial coefficients of Carathéodory functions (see also [8]). We recall the Möbius transformation  $\psi_\zeta : \mathbb{D} \rightarrow \mathbb{D}, \zeta \in \mathbb{D}$ , defined by

$$\psi_\zeta(z) = \frac{z - \zeta}{1 - \bar{\zeta}z} \tag{7}$$

and let

$$L(z) = \frac{1+z}{1-z}, \quad z \in \mathbb{D}. \tag{8}$$

**Lemma 3 ([7]).** If  $p \in \mathcal{P}$  is of the form (3), then

$$p_1 = 2\zeta_1, \tag{9}$$

$$p_2 = 2\zeta_1^2 + 2(1 - |\zeta_1|^2)\zeta_2 \tag{10}$$

and

$$p_3 = 2\zeta_1^3 + 4(1 - |\zeta_1|^2)\zeta_1\zeta_2 - 2(1 - |\zeta_1|^2)\bar{\zeta}_1\zeta_2^2 + 2(1 - |\zeta_1|^2)(1 - |\zeta_2|^2)\zeta_3 \tag{11}$$

for some  $\zeta_i \in \bar{\mathbb{D}}, i \in \{1, 2, 3\}$ . For  $\zeta_1 \in \mathbb{T}$ , there is a unique function  $p \in \mathcal{P}$  with  $p_1$  as in (9), namely,

$$p(z) = \frac{1 + \zeta_1 z}{1 - \zeta_1 z}, \quad z \in \mathbb{D}.$$

For  $\zeta_1 \in \mathbb{D}$  and  $\zeta_2 \in \mathbb{T}$ , there is a unique function  $p = L \circ \omega \in \mathcal{P}$  with  $p_1$  and  $p_2$  as in (9)-(10), where

$$\omega(z) = z\psi_{-\zeta_1}(\zeta_2 z), \quad z \in \mathbb{D},$$

i.e.,

$$p(z) = \frac{1 + (\bar{\zeta}_1 \zeta_2 + \zeta_1)z + \zeta_2 z^2}{1 + (\bar{\zeta}_1 \zeta_2 - \zeta_1)z - \zeta_2 z^2}, \quad z \in \mathbb{D}.$$

For  $\zeta_1, \zeta_2 \in \mathbb{D}$  and  $\zeta_3 \in \mathbb{T}$ , there is a unique function  $p = L \circ \omega \in \mathcal{P}$  with  $p_1, p_2$  and  $p_3$  as in (9)-(11), where

$$\omega(z) = z\psi_{-\zeta_1}(z\psi_{-\zeta_2}(\zeta_3 z)), \quad z \in \mathbb{D},$$

i.e.,

$$p(z) = \frac{1 + (\bar{\zeta}_2 \zeta_3 + \bar{\zeta}_1 \zeta_2 + \zeta_1)z + (\bar{\zeta}_1 \zeta_3 + \zeta_1 \bar{\zeta}_2 \zeta_3 + \zeta_2)z^2 + \zeta_3 z^3}{1 + (\bar{\zeta}_2 \zeta_3 + \bar{\zeta}_1 \zeta_2 - \zeta_1)z + (\bar{\zeta}_1 \zeta_3 - \zeta_1 \bar{\zeta}_2 \zeta_3 - \zeta_2)z^2 - \zeta_3 z^3}, \quad z \in \mathbb{D}.$$

Conversely, if  $\zeta_1, \zeta_2 \in \mathbb{D}$  and  $\zeta_3 \in \bar{\mathbb{D}}$  are given, then we can construct a (unique) function  $p \in \mathcal{P}$  of the form (3) so that  $p_i, i \in \{1, 2, 3\}$ , satisfying the identities in (9)-(11). For this, we define

$$\omega(z) = \omega_{\zeta_1, \zeta_2, \zeta_3}(z) = z\psi_{-\zeta_1}(z\psi_{-\zeta_2}(\zeta_3 z)), \quad z \in \mathbb{D}, \tag{12}$$

where  $\psi_\zeta$  is the function defined as in (7). Then  $\omega \in \mathcal{B}$ . Moreover, if we define  $p(z) = (1 + \omega(z))/(1 - \omega(z)), z \in \mathbb{D}$ , then  $p$  is represented by (3), where  $p_1, p_2$  and  $p_3$  satisfy the identities in (9)-(11) (see the proof of ([7], Lemma 2.4)).

Assume that the function  $q(z) = 1 + \sum_{n=1}^{\infty} b_n z^n \in \mathcal{P}$  is constructed by  $\zeta_1, \zeta_2 \in \mathbb{D}, \zeta_3 \in \bar{\mathbb{D}}$ , and the

function  $h(z) = 1 + \sum_{n=1}^{\infty} u_n z^n \in \mathcal{P}$  is constructed by  $\zeta_1, \zeta_2 \in \mathbb{D}, \zeta_3 \in \bar{\mathbb{D}}$ . Namely,  $q = L \circ \omega_1$  and  $h = L \circ \omega_2$ , where  $L$  is the function defined by (8),  $\omega_1(z) = z\psi_{-\zeta_1}(z\psi_{-\zeta_2}(\zeta_3 z))$  and  $\omega_2(z) = z\psi_{-\zeta_1}(z\psi_{-\zeta_2}(\zeta_3 z))$ . Then by combining the above argument, we conclude (B) below.

(B) Let  $p \in \mathcal{P}$  be in the form (3). If there exist  $\zeta_1, \zeta_2, \zeta_1, \zeta_2 \in \mathbb{D}, \zeta_3, \zeta_3 \in \bar{\mathbb{D}}$  satisfying the following conditions

$$\left\{ \begin{array}{l} b_1 = 2\zeta_1 \\ b_2 = 2\zeta_1^2 + 2(1 - |\zeta_1|^2)\zeta_2 \\ b_3 = 2\zeta_1^3 + 4(1 - |\zeta_1|^2)\zeta_1\zeta_2 - 2(1 - |\zeta_1|^2)\bar{\zeta}_1\zeta_2^2 + 2(1 - |\zeta_1|^2)(1 - |\zeta_2|^2)\zeta_3 \\ u_1 = 2\zeta_1 \\ u_2 = 2\zeta_1^2 + 2(1 - |\zeta_1|^2)\zeta_2 \\ u_3 = 2\zeta_1^3 + 4(1 - |\zeta_1|^2)\zeta_1\zeta_2 - 2(1 - |\zeta_1|^2)\bar{\zeta}_1\zeta_2^2 + 2(1 - |\zeta_1|^2)(1 - |\zeta_2|^2)\zeta_3 \\ b_4 = 2 \\ B_1 = -\frac{1}{128} \left( 1 + \frac{3}{2}u_1 + \frac{3}{2}u_2 + \frac{1}{2}u_3 \right) b_1^4 \\ B_2 = \frac{3}{32} \left( 1 + u_1 + \frac{1}{2}u_2 \right) b_1^2 b_2 \\ B_3 = -\frac{1}{4} \left( 1 + \frac{1}{2}u_1 \right) b_1 b_3 \\ B_4 = -\frac{1}{8} \left( 1 + \frac{1}{2}u_1 \right) b_2^2, \end{array} \right.$$

then  $|\Psi| = |A_4| \leq 2$ , where  $\Psi$  and  $A_4$  are given by (5) and (6), respectively.

Since the system of equations in (B) has many solutions, we now place some restrictions on the parameters sufficient for our purpose.

We fix

$$q(z) = q_\tau(z) = \frac{1 + 2\tau z + z^2}{1 - z^2}.$$

Then if  $\tau \in [-1, 1]$ ,  $q \in \mathcal{P}$  and is given by  $q(z) = 1 + \sum_{n=1}^{\infty} b_n z^n$ , with

$$b_1 = b_3 = 2\tau, \quad \text{and} \quad b_2 = b_4 = 2.$$

We also assume that  $\zeta_i, i \in \{1, 2, 3\}$  take real values. Then the identities for  $u_i, i \in \{1, 2, 3\}$  become

$$\begin{cases} u_1 = 2\zeta_1 \\ u_2 = 2\zeta_1^2 + 2(1 - \zeta_1^2)\zeta_2 \\ u_3 = 2\zeta_1^3 + 4(1 - \zeta_1^2)\zeta_1\zeta_2 - 2(1 - |\zeta_1|^2)\zeta_1\zeta_2^2 + 2(1 - \zeta_1^2)(1 - \zeta_2^2)\zeta_3. \end{cases}$$

Thus we are able to conclude the following.

(C) Let  $p \in \mathcal{P}$  be in the form (3). If there exist  $\zeta_1, \zeta_2 \in (-1, 1)$  and  $\zeta_3, \tau \in [-1, 1]$  such that

$$\begin{cases} u_1 = 2\zeta_1 \\ u_2 = 2\zeta_1^2 + 2(1 - \zeta_1^2)\zeta_2 \\ u_3 = 2\zeta_1^3 + 4(1 - \zeta_1^2)\zeta_1\zeta_2 - 2(1 - |\zeta_1|^2)\zeta_1\zeta_2^2 + 2(1 - \zeta_1^2)(1 - \zeta_2^2)\zeta_3 \\ b_4 = 2 \\ B_1 = -\frac{1}{8} \left( 1 + \frac{3}{2}u_1 + \frac{3}{2}u_2 + \frac{1}{2}u_3 \right) \tau^4 \\ B_2 = \frac{3}{4} \left( 1 + u_1 + \frac{1}{2}u_2 \right) \tau^2 \\ B_3 = - \left( 1 + \frac{1}{2}u_1 \right) \tau^2 \\ B_4 = -\frac{1}{2} \left( 1 + \frac{1}{2}u_1 \right), \end{cases}$$

then  $|\Psi| = |A_4| \leq 2$ , where  $\Psi$  and  $A_4$  are given by (5) and (6), respectively.

### 3. The Fifth Coefficient of $\mathcal{B}_1(\alpha)$ Bazilevič Functions

**Lemma 4.** ([9] Cohn’s rule) Let  $t(z) = a_0 + a_1z + \dots + a_nz^n$  be a polynomial of degree  $n$  and

$$t^*(z) = z^n \overline{t(1/\bar{z})} = \bar{a}_n + \bar{a}_{n-1}z + \dots + \bar{a}_0z^n.$$

Let  $r$  and  $s$  be the number of zeros of  $t$  inside and on the unit circle  $|z| = 1$ , respectively. If  $|a_0| < |a_n|$ , then

$$t_1(z) = \frac{\bar{a}_n t(n) - a_0 t^*(z)}{z}$$

is a polynomial of degree  $n - 1$  and has  $r_1 = r - 1$  and  $s_1 = s$  number of zeros inside and on the unit circle  $|z| = 1$ , respectively.

We now use the above method to find the sharp bound for  $|a_5|$  when  $\alpha \geq 1$ .

**Theorem 1.** Let  $f \in \mathcal{B}_1(\alpha)$ , and be given by (1), then  $|a_5| \leq 2/(4 + \alpha)$ , provided  $\alpha \geq 1$ . The inequality is sharp, with extreme function  $f \in \mathcal{A}$  defined by

$$f'(z) \left( \frac{f(z)}{z} \right)^{\alpha-1} = \frac{1 + z^5}{1 - z^5}. \tag{13}$$

**Proof.** From (2) we can write

$$f'(z) \left( \frac{f(z)}{z} \right)^{\alpha-1} = p(z), \quad (14)$$

for some function  $p \in \mathcal{P}$  of the form (3). Putting the series (1) and (3) into (14) by equating the coefficients we get

$$p_1 = (1 + \alpha)a_2, \quad p_2 = \frac{1}{2}(2 + \alpha)[(-1 + \alpha)a_2^2 + 2a_3], \quad (15)$$

$$p_3 = \frac{1}{6}(3 + \alpha)[(2 - 3\alpha + \alpha^2)a_2^3 + 6(-1 + \alpha)a_2a_3 + 6a_4] \quad (16)$$

and

$$p_4 = \frac{1}{24}(4 + \alpha)[(-6 + 11\alpha - 6\alpha^2 + \alpha^3)a_2^4 + 12(3 - 3\alpha + \alpha^2)a_2^2a_3 + 24(-1 + \alpha)a_2a_4 + 12((-1 + \alpha)a_3^2 + 2a_5)]. \quad (17)$$

From the equalities (15), (16) and (17),  $a_5$  can be written as

$$a_5 = \frac{1}{4 + \alpha} \cdot \Psi, \quad (18)$$

where

$$\Psi = p_4 + (4 + \alpha)(1 - \alpha)(\mu_1 p_1 p_3 + \mu_2 p_2^2 + \mu_3 p_1^2 p_2 + \mu_4 p_1^4),$$

with

$$\mu_1 = \frac{1}{(1 + \alpha)(3 + \alpha)}, \quad \mu_2 = \frac{1}{2(2 + \alpha)^2},$$

$$\mu_3 = \frac{1 - 2\alpha}{2(1 + \alpha)^2(2 + \alpha)}, \quad \text{and} \quad \mu_4 = \frac{(3\alpha - 1)(2\alpha - 1)}{24(1 + \alpha)^4}.$$

Thus it is enough to show that  $|\Psi| \leq 2$ .

When  $\alpha = 1$ , it is clear that  $|\Psi| \leq 2$  holds trivially, and so we can assume that  $\alpha > 1$ .

Put

$$\zeta_1 = -\frac{25 + 28\alpha + 10\alpha^2 + \alpha^3}{(3 + 4\alpha + \alpha^2)^2},$$

$$\zeta_2 = \frac{12 + 42\alpha + 32\alpha^2 + 9\alpha^3 + \alpha^4}{3(34 + 52\alpha + 32\alpha^2 + 9\alpha^3 + \alpha^4)},$$

and

$$\zeta_3 = -\frac{256 + 684\alpha + 770\alpha^2 + 438\alpha^3 + 134\alpha^4 + 21\alpha^5 + \alpha^6}{4(5 + 3\alpha + \alpha^2)(57 + 156\alpha + 163\alpha^2 + 82\alpha^3 + 20\alpha^4 + 2\alpha^5)},$$

then  $|\zeta_i| < 1$  ( $i = 1, 2, 3$ ) when  $\alpha > 1$ .

Now let  $\omega \in \mathbb{B}$  be defined by (12). Thus the function  $k$  defined by  $k = L \circ \omega$ , where  $L$  is given by (8), belongs to  $\mathcal{P}$ , with  $k(0) = 1$ .

Setting

$$k(z) = 1 + \sum_{n=1}^{\infty} \beta_n z^n,$$

we obtain

$$\beta_1 = -\frac{2(25 + 28\alpha + 10\alpha^2 + \alpha^3)}{(3 + 4\alpha + \alpha^2)^2},$$

$$\beta_2 = \frac{2(187 + 262\alpha + 173\alpha^2 + 94\alpha^3 + 41\alpha^4 + 10\alpha^5 + \alpha^6)}{3(1 + \alpha)^4(3 + \alpha)^2},$$

and

$$\beta_3 = -\frac{2(427 + 696\alpha + 688\alpha^2 + 641\alpha^3 + 427\alpha^4 + 161\alpha^5 + 30\alpha^6 + 2\alpha^7)}{3(1 + \alpha)^6(3 + \alpha)^2}.$$

With  $\epsilon_n$  ( $n \in \mathbb{N}$ ) as in (4), we obtain

$$\epsilon_1 = \frac{(2 + \alpha)^2(-4 + 3\alpha + \alpha^2)}{2(3 + 4\alpha + \alpha^2)^2}, \quad (19)$$

$$\epsilon_2 = \frac{(2 + \alpha)^3(4 - 11\alpha + 5\alpha^2 + 2\alpha^3)}{6(1 + \alpha)^4(3 + \alpha)^2}, \quad (20)$$

and

$$\epsilon_3 = \frac{(2 + \alpha)^4(-4 + 23\alpha - 38\alpha^2 + 13\alpha^3 + 6\alpha^4)}{24(1 + \alpha)^6(3 + \alpha)^2}. \quad (21)$$

Now define  $q$  by

$$q(z) = \frac{1 + 2\tau z + 2\tau^2 z^2 + 2\tau z^3 + z^4}{1 - z^4} = 1 + \sum_{n=1}^{\infty} c_n z^n$$

with  $\tau := \sqrt{3 + 4\alpha + \alpha^2}/(2 + \alpha)$ . Then  $0 < \tau < 1$  for  $\alpha > 1$ , and

$$c_1 = c_3 = 2\tau, \quad c_2 = 2\tau^2 \quad \text{and} \quad c_4 = 2. \quad (22)$$

We shall show that  $q$  belongs  $\mathcal{P}$ . Let

$$\omega(z) = \frac{q(z) - 1}{q(z) + 1} = \frac{zA(z)}{B(z)}, \quad (23)$$

where  $A$  and  $B$  are polynomials of degree 3, and defined by

$$A(z) = \tau + \tau^2 z + \tau z^2 + z^3 \quad \text{and} \quad B(z) = 1 + \tau z + \tau^2 z^2 + \tau z^3,$$

respectively. Then it holds that

$$B(z) = z^3 \overline{A(1/\bar{z})}. \quad (24)$$

Define a function  $\tilde{A}$  by  $\tilde{A}(z) = (A(z) - \tau B(z))/z$ . Then a computation gives

$$\tilde{A}(z) = (1 - \tau^2)(\tau + z)z = 0.$$

Therefore  $\tilde{A}$  has two zeros in  $\mathbb{D}$ . By Lemma 4, the function  $A$  has exactly three zeros, say  $w_1, w_2$  and  $w_3$ , in  $\mathbb{D}$ . Hence, from (23) and (24), we have

$$\omega(z) = z \prod_{i=1}^3 \frac{z - w_i}{1 - \bar{w}_i z},$$

which implies that  $\omega$  belongs to  $\mathcal{B}$ . It follows from (23) that  $q \in \mathcal{P}$  as we asserted.

Since  $p \in \mathcal{P}$ , by Lemma 1 the function

$$h(z) := 1 + \frac{1}{2} \sum_{n=1}^{\infty} c_n p_n z^n$$

also belongs to  $\mathcal{P}$ . Therefore from Lemma 2 with  $d_n = c_n p_n / 2$  we have  $|A_4| \leq 2$ , where

$$A_4 = \frac{1}{2}c_4 p_4 - \frac{1}{4}\epsilon_1(c_2^2 p_2^2 + 2c_1 c_3 p_1 p_3) + \frac{3}{8}\epsilon_2 c_1^2 c_2 p_1^2 p_2 - \frac{1}{16}\epsilon_3 c_1^4 p_1^4.$$

Finally, (19)–(22) shows that  $A_4 = \Psi$ , and so  $|\Psi| \leq 2$ . Thus it follows from (18) that the inequality  $|a_5| \leq 2/(4 + \alpha)$  holds.

Let  $f \in \mathcal{B}_1(\alpha)$  be the function defined by (13). Then, by equating coefficients in (13), we get  $a_2 = a_3 = a_4 = 0$  and  $a_5 = 2/(4 + \alpha)$ , which shows that this result is sharp. This completes the proof of Theorem 1.  $\square$

#### 4. Conclusions

There are several instances in the literature where only partial solutions are known for the bounds for  $|a_5|$  for functions in subclasses of  $\mathcal{S}$  (again, see [1]). Applying the method introduced in this paper may well provide improved, or complete solutions to some of these.

**Author Contributions:** All authors have equal contributions. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was partially supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIP; Ministry of Science, ICT & Future Planning) (No. NRF-2017R1C1B5076778).

**Conflicts of Interest:** The authors declare no conflict of interest.

#### References

1. Thomas, D.K.; Tuneski, N.; Vasudevarao, A. *Univalent Functions: A Primer*; De Gruyter Studies in Mathematics 69; Walter de Gruyter GmbH: Berlin, Germany, 2018.
2. Singh, R. On Bazilevič Functions. *Proc. Am. Math. Soc.* **1973**, *38*, 261–271.
3. Cho, N.E.; Kumar, V. On a conjecture for Bazilevič functions. *Bull. Malaysian Math. Soc.* **2019**. [[CrossRef](#)]
4. Marjono, J.S.; Thomas, D.K. The fifth and sixth coefficients of Bazilevič Functions  $\mathcal{B}_1(\alpha)$ . *Mediterr. J. Math.* **2017**, *14*, 158. [[CrossRef](#)]
5. Schur, I. Über Potenzreihen, die im Innern des Einheitskreises beschränkt sind. *J. Reine Angew. Math.* **1917**, *147*, 205–232.
6. Nehari, Z.; Netanyahu, E. On the coefficients of meromorphic schlicht functions. *Proc. Am. Math. Soc.* **1957**, *8*, 15–23. [[CrossRef](#)]
7. Cho, N.E.; Kowalczyk, V.; Lecko, A. The sharp bounds of some coefficient functionals over the class of functions convex in the direction of the imaginary axis. *Bull. Aust. Math. Soc.* **2019**, *100*, 86–96. [[CrossRef](#)]
8. Li, M.; Sugawa, T. Schur parameters and the Carathéodory class. *Results Math.* **2019**, *74*, 185. [[CrossRef](#)]
9. Rahman Q. I.; Schmeisser, G. *Analytic theory of polynomials*, London Mathematical Society Monographs. New Series, 26. The Clarendon Press, Oxford University Press, Oxford, 2002.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).