

Article

\mathcal{L} -Single Valued Extremally Disconnected Ideal Neutrosophic Topological Spaces

Fahad Alsharari 

Department of Mathematics, College of Science and Human Studies, Hotat Sudair, Majmaah University, Majmaah 11952, Saudi Arabia; f.alsharari@mu.edu.sa

Abstract: This paper aims to mark out new concepts of r -single valued neutrosophic sets, called r -single valued neutrosophic \mathcal{L} -closed and \mathcal{L} -open sets. The definition of \mathcal{L} -single valued neutrosophic irresolute mapping is provided and its characteristic properties are discussed. Moreover, the concepts of \mathcal{L} -single valued neutrosophic extremally disconnected and \mathcal{L} -single valued neutrosophic normal spaces are established. As a result, a useful implication diagram between the r -single valued neutrosophic ideal open sets is obtained. Finally, some kinds of separation axioms, namely r -single valued neutrosophic ideal- R_i (r -SVNIR $_i$, for short), where $i = \{0, 1, 2, 3\}$, and r -single valued neutrosophic ideal- T_j (r -SVNIT $_j$, for short), where $j = \{1, 2, 2\frac{1}{2}, 3, 4\}$, are introduced. Some of their characterizations, fundamental properties, and the relations between these notions have been studied.

Keywords: r -single valued neutrosophic \mathcal{L} -closed; \mathcal{L} -single valued neutrosophic irresolute mapping; \mathcal{L} -single valued neutrosophic extremally disconnected; \mathcal{L} -single valued neutrosophic normal; r -SVNIR $_i$; r -SVNIT $_j$



Citation: Alsharari, F. \mathcal{L} -Single Valued Extremally Disconnected Ideal Neutrosophic Topological Spaces. *Symmetry* **2021**, *13*, 53. <https://doi.org/10.3390/sym13010053>

Received: 12 November 2020
Accepted: 29 December 2020
Published: 31 December 2020

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2020 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

In 1999, Smarandache introduced the concept of a neutrosophy [1]. It has been used at various axes of mathematical theories and applications. In recent decades, the theory made an outstanding advancement in the field of topological spaces. Salama et al. and Hur et al. [2–6], for example, among many others, wrote their works in fuzzy neutrosophic topological spaces (FNTS), following Chang [7]'s discoveries in the way of fuzzy topological spaces (FTS).

Šostak, in 1985 [8], marked out a new definition of fuzzy topology as a crisp subfamily of family of fuzzy sets, which seems to be a drawback in the process of fuzzification of the concept of topological spaces. Yan, Wang, Nanjing, Liang, and Yan [9,10] developed a parallel theory in the context of intuitionistic I -fuzzy topological spaces.

The idea of “single-valued neutrosophic set” [11] was set out by Wang in 2010. Gayyar [12], in his 2016 paper, foregrounded the concept of a “smooth neutrosophic topological spaces”. The ordinary single-valued neutrosophic topology was presented by Kim [13]. Recently, Saber et al. [14,15] familiarized the concepts of single-valued neutrosophic ideal open local function, single-valued neutrosophic topological space, and the connectedness and stratification of single-valued neutrosophic topological spaces.

Neutrosophy, and especially neutrosophic sets, are powerful, general, and formal frameworks that generalize the concept of the ordinary sets, fuzzy sets, and intuitionistic fuzzy sets from philosophical point of view. This paper sets out to introduce and examine a new class of sets called r -single valued \mathcal{L} -closed in the single valued neutrosophic topological spaces in Šostak's sense. More precisely, different attributes, like \mathcal{L} -single valued neutrosophic irresolute mapping, \mathcal{L} -single valued neutrosophic extremally disconnected, \mathcal{L} -single valued neutrosophic normal spaces, and some kinds of separation axioms, were developed. It can be fairly claimed that we have achieved expressive definitions, distinguished theorems, important lemmas, and counterexamples to investigate, in-depth, our

consequences and to find out the best results. It is notable to say that different crucial notions in single valued neutrosophic topology were generalized in this article. Different attributes, like extremally disconnected and some kinds of separation axioms, which have a significant impact on the overall topology’s notions, were also studied.

It is notable to say that the application aspects to this area of research can be further pointed to. There are many applications of neutrosophic theories in many branches of sciences. Possible applications are to control engineering and to Geographical Information Systems, and so forth, and could be secured, as mentioned by many authors, such as Reference [16–20].

In this study, \tilde{X} is assumed to be a nonempty set, $\zeta = [0, 1]$ and $\zeta_0 = (0, 1]$. For $\alpha \in \zeta$, $\tilde{\alpha}(v) = \alpha$ for all $v \in \tilde{X}$. The family of all single-valued neutrosophic sets on \tilde{X} is denoted by $\zeta^{\tilde{X}}$.

2. Preliminaries

This section is devoted to provide a complete survey and trace previous studies related to the idea of this research article.

Definition 1 ([21]). Let \tilde{X} be a non-empty set. A neutrosophic set (briefly, NS) in \tilde{X} is an object having the form

$$\sigma_n = \{ \langle v, \tilde{\rho}_{\sigma_n}(v), \tilde{q}_{\sigma_n}(v), \tilde{\eta}_{\sigma_n}(v) \rangle : v \in \tilde{X} \},$$

where

$$\tilde{\rho} : \tilde{X} \rightarrow]^{-0, 1^+}[, \tilde{q} : \tilde{X} \rightarrow]^{-0, 1^+}[, \tilde{\eta} : \tilde{X} \rightarrow]^{-0, 1^+}[$$

and

$$^{-0} \leq \tilde{\rho}_{\sigma_n}(v) + \tilde{q}_{\sigma_n}(v) + \tilde{\eta}_{\sigma_n}(v) \leq 3^+$$

represent the degree of membership (namely $\tilde{\rho}_{\sigma_n}(v)$), the degree of indeterminacy (namely $\tilde{q}_{\sigma_n}(v)$), and the degree of non-membership (namely $\tilde{\eta}_{\sigma_n}(v)$), respectively, of any $v \in \tilde{X}$ to the set σ_n .

Definition 2 ([11]). Let \tilde{X} be a space of points (objects), with a generic element in \tilde{X} denoted by v . Then, σ_n is called a single valued neutrosophic set (briefly, SVNS) in \tilde{X} , if σ_n has the form $\sigma_n = \langle \tilde{\rho}_{\sigma_n}, \tilde{q}_{\sigma_n}, \tilde{\eta}_{\sigma_n} \rangle$, where $\tilde{\rho}_{\sigma_n}, \tilde{q}_{\sigma_n}, \tilde{\eta}_{\sigma_n} : \tilde{X} \rightarrow [0, 1]$. In this case, $\tilde{\rho}_{\sigma_n}, \tilde{q}_{\sigma_n}, \tilde{\eta}_{\sigma_n}$ are called truth membership function, indeterminacy membership function, and falsity membership function, respectively.

Let \tilde{X} be a nonempty set and $\zeta = [0, 1]$ and $\zeta_0 = (0, 1]$. A single-valued neutrosophic set σ_n on \tilde{X} is a mapping defined as $\sigma_n = \langle \tilde{\rho}_{\sigma_n}, \tilde{q}_{\sigma_n}, \tilde{\eta}_{\sigma_n} \rangle : \tilde{X} \rightarrow \zeta$ such that $0 \leq \tilde{\rho}_{\sigma_n}(v) + \tilde{q}_{\sigma_n}(v) + \tilde{\eta}_{\sigma_n}(v) \leq 3$.

We denote the single-valued neutrosophic sets $\langle 0, 1, 1 \rangle$ and $\langle 1, 0, 0 \rangle$ by $\tilde{0}$ and $\tilde{1}$, respectively.

Definition 3 ([11]). Let $\sigma_n = \langle \tilde{\rho}_{\sigma_n}, \tilde{q}_{\sigma_n}, \tilde{\eta}_{\sigma_n} \rangle$ be an SVNS on \tilde{X} . The complement of the set σ_n (briefly σ_n^c) is defined as follows:

$$\tilde{\rho}_{\sigma_n^c}(v) = \tilde{\eta}_{\sigma_n}(v), \quad \tilde{q}_{\sigma_n^c}(v) = [\tilde{q}_{\sigma_n}]^c(v), \quad \tilde{\eta}_{\sigma_n^c}(v) = \tilde{\rho}_{\sigma_n}(v).$$

Definition 4 ([22,23]). Let \tilde{X} be a non-empty set and let $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ be given by $\sigma_n = \langle \tilde{\rho}_{\sigma_n}, \tilde{q}_{\sigma_n}, \tilde{\eta}_{\sigma_n} \rangle$ and $\gamma_n = \langle \tilde{\rho}_{\gamma_n}, \tilde{q}_{\gamma_n}, \tilde{\eta}_{\gamma_n} \rangle$. Then:

- (1) We say that $\sigma_n \subseteq \gamma_n$ if $\tilde{\rho}_{\sigma_n} \leq \tilde{\rho}_{\gamma_n}, \tilde{q}_{\sigma_n} \geq \tilde{q}_{\gamma_n}, \tilde{\eta}_{\sigma_n} \geq \tilde{\eta}_{\gamma_n}$.
- (2) The intersection of σ_n and γ_n denoted by $\sigma_n \cap \gamma_n$ is an SVNS and is given by

$$\sigma_n \cap \gamma_n = \langle \tilde{\rho}_{\sigma_n} \cap \tilde{\rho}_{\gamma_n}, \tilde{q}_{\sigma_n} \cup \tilde{q}_{\gamma_n}, \tilde{\eta}_{\sigma_n} \cup \tilde{\eta}_{\gamma_n} \rangle.$$

- (3) The union of σ_n and γ_n denoted by $\sigma_n \cup \gamma_n$ is an SVNS and is given by

$$\sigma_n \cup \gamma_n = \langle \tilde{\rho}_{\sigma_n} \cup \tilde{\rho}_{\gamma_n}, \tilde{q}_{\sigma_n} \cap \tilde{q}_{\gamma_n}, \tilde{\eta}_{\sigma_n} \cap \tilde{\eta}_{\gamma_n} \rangle.$$

For any arbitrary family $\{\sigma_n\}_{i \in j} \subseteq \zeta^{\tilde{X}}$ of SVNS, the union and intersection are given by

$$(4) \quad \bigcap_{i \in j} [\sigma_n]_i = \langle \bigcap_{i \in j} \tilde{\rho}_{[\sigma_n]_i}, \bigcup_{i \in j} \tilde{q}_{[\sigma_n]_i}, \bigcup_{i \in j} \tilde{\eta}_{[\sigma_n]_i} \rangle,$$

$$(5) \quad \bigcup_{i \in j} [\sigma_n]_i = \langle \bigcup_{i \in j} \tilde{\rho}_{[\sigma_n]_i}, \bigcap_{i \in j} \tilde{q}_{[\sigma_n]_i}, \bigcap_{i \in j} \tilde{\eta}_{[\sigma_n]_i} \rangle.$$

Definition 5 ([12]). A single-valued neutrosophic topological space is an ordered quadruple $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}}, \tilde{\tau}^{\tilde{q}}, \tilde{\tau}^{\tilde{\eta}})$ where $\tilde{\tau}^{\tilde{\rho}}, \tilde{\tau}^{\tilde{q}}, \tilde{\tau}^{\tilde{\eta}} : \zeta^{\tilde{X}} \rightarrow \zeta$ are mappings satisfying the following axioms:

$$(SVNT1) \quad \tilde{\tau}^{\tilde{\rho}}(\tilde{0}) = \tilde{\tau}^{\tilde{\rho}}(\tilde{1}) = 1 \text{ and } \tilde{\tau}^{\tilde{\rho}}(\tilde{0}) = \tilde{\tau}^{\tilde{\rho}}(\tilde{1}) = \tilde{\tau}^{\tilde{\eta}}(\tilde{0}) = \tilde{\tau}^{\tilde{\eta}}(\tilde{1}) = 0,$$

$$(SVNT2) \quad \tilde{\tau}^{\tilde{\rho}}(\sigma_n \cap \gamma_n) \geq \tilde{\tau}^{\tilde{\rho}}(\sigma_n) \cap \tilde{\tau}^{\tilde{\rho}}(\gamma_n), \quad \tilde{\tau}^{\tilde{q}}(\sigma_n \cap \gamma_n) \leq \tilde{\tau}^{\tilde{q}}(\sigma_n) \cup \tilde{\tau}^{\tilde{q}}(\gamma_n),$$

$$\tilde{\tau}^{\tilde{\eta}}(\sigma_n \cap \gamma_n) \leq \tilde{\tau}^{\tilde{\eta}}(\sigma_n) \cup \tilde{\tau}^{\tilde{\eta}}(\gamma_n), \text{ for all } \sigma_n, \gamma_n \in \zeta^{\tilde{X}},$$

$$(SVNT3) \quad \tilde{\tau}^{\tilde{\rho}}(\bigcup_{j \in \Gamma} [\sigma_n]_j) \geq \bigcap_{j \in \Gamma} \tilde{\tau}^{\tilde{\rho}}([\sigma_n]_j), \quad \tilde{\tau}^{\tilde{q}}(\bigcup_{j \in \Gamma} [\sigma_n]_j) \leq \bigcup_{j \in \Gamma} \tilde{\tau}^{\tilde{q}}([\sigma_n]_j),$$

$$\tilde{\tau}^{\tilde{\eta}}(\bigcup_{j \in \Gamma} [\sigma_n]_j) \leq \bigcup_{j \in \Gamma} \tilde{\tau}^{\tilde{\eta}}([\sigma_n]_j) \text{ for all } \{[\sigma_n]_j, j \in \Gamma\} \in \zeta^{\tilde{X}}.$$

The quadruple $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}}, \tilde{\tau}^{\tilde{q}}, \tilde{\tau}^{\tilde{\eta}})$ is called a single-valued neutrosophic topological space (SVNTS, for short). We will occasionally write $\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}$ for $(\tau^{\tilde{\rho}}, \tau^{\tilde{q}}, \tau^{\tilde{\eta}})$ and it will cause no ambiguity

Definition 6 ([14]). Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}}, \tilde{\tau}^{\tilde{q}}, \tilde{\tau}^{\tilde{\eta}})$ be an SVNTS. Then, for every $\sigma_n \in \zeta^{\tilde{X}}$ and $r \in \zeta_0$, the single valued neutrosophic closure and the single valued neutrosophic interior of σ_n are defined by:

$$C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\sigma_n, s) = \bigcap \{ \gamma_n \in \zeta^{\tilde{X}} : \sigma_n \leq \gamma_n, \quad \tau^{\tilde{\rho}}([\gamma_n]^c) \geq r, \quad \tau^{\tilde{q}}([\gamma_n]^c) \leq 1 - r, \quad \tau^{\tilde{\eta}}([\gamma_n]^c) \leq 1 - r \},$$

$$\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\sigma_n, s) = \bigcup \{ \gamma_n \in \zeta^{\tilde{X}} : \sigma_n \geq \gamma_n, \quad \tau^{\tilde{\rho}}(\gamma_n) \geq r, \quad \tau^{\tilde{q}}(\gamma_n) \leq 1 - r, \quad \tau^{\tilde{\eta}}(\gamma_n) \leq 1 - r \}.$$

Definition 7 ([24]). Let $(\tilde{X}, \tau^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNTS and $r \in \zeta_0, \sigma_n \in \zeta^{\tilde{X}}$. Then,

$$(1) \quad \sigma_n \text{ is } r\text{-single valued neutrosophic semiopen (} r\text{-SVNSO, for short) iff } \sigma_n \leq C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\sigma_n, r), r),$$

$$(2) \quad \sigma_n \text{ is } r\text{-single valued neutrosophic } \beta\text{-open (} r\text{-SVN}\beta\text{O, for short) iff } \sigma_n \leq C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\sigma_n, r), r), r).$$

The complement of r -SVNSO (resp. r -SVN β O) is said to be an r -SVNSC (resp. r -SVN β C), respectively.

Definition 8 ([14]). Let \tilde{X} be a nonempty set and $v \in \tilde{X}$. If $s \in (0, 1], t \in [0, 1)$ and $p \in [0, 1)$. Then, the single-valued neutrosophic point $x_{s,t,p}$ in \tilde{X} is given by

$$x_{s,t,p}(\kappa) = \begin{cases} (s, t, p), & \text{if } x = v, \\ (0, 1, 1), & \text{otherwise.} \end{cases}$$

We say $x_{s,t,p} \in \sigma_n$ iff $s < \tilde{\rho}_{\sigma_n}(v)$, $t \geq \tilde{q}_{\sigma_n}(v)$ and $p \geq \tilde{\eta}_{\sigma_n}(v)$. To avoid the ambiguity, we denote the set of all neutrosophic points by $pt(\zeta^{\tilde{X}})$.

A single-valued neutrosophic set σ_n is said to be quasi-coincident with another single-valued neutrosophic set γ_n , denoted by $\sigma_n q \gamma_n$, if there exists an element $v \in \tilde{X}$ such that

$$\tilde{\rho}_{\sigma_n}(v) + \tilde{\rho}_{\gamma_n}(v) > 1, \quad \tilde{q}_{\sigma_n}(v) + \tilde{q}_{\gamma_n}(v) \leq 1, \quad \tilde{\eta}_{\sigma_n}(v) + \tilde{\eta}_{\gamma_n}(v) \leq 1.$$

Definition 9 ([14]). A mapping $\mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}} = \mathcal{I}^{\tilde{\rho}}, \mathcal{I}^{\tilde{q}}, \mathcal{I}^{\tilde{\eta}} : \zeta^{\tilde{X}} \rightarrow \zeta$ is called single-valued neutrosophic ideal (SVNI) on \tilde{X} if it satisfies the following conditions:

$$(I_1) \quad \mathcal{I}^{\tilde{\rho}}(\tilde{0}) = 1 \text{ and } \mathcal{I}^{\tilde{q}}(\tilde{0}) = \mathcal{I}^{\tilde{\eta}}(\tilde{0}) = 0.$$

$$(I_2) \quad \text{If } \sigma_n \leq \gamma_n, \text{ then } \mathcal{I}^{\tilde{\rho}}(\gamma_n) \leq \mathcal{I}^{\tilde{\rho}}(\sigma_n), \mathcal{I}^{\tilde{q}}(\gamma_n) \geq \mathcal{I}^{\tilde{q}}(\sigma_n), \text{ and } \mathcal{I}^{\tilde{\eta}}(\gamma_n) \geq \mathcal{I}^{\tilde{\eta}}(\sigma_n), \text{ for } \gamma_n, \sigma_n \in \zeta^{\tilde{X}}.$$

$$(I_3) \quad \mathcal{I}^{\tilde{\rho}}(\sigma_n \cup \gamma_n) \geq \mathcal{I}^{\tilde{\rho}}(\sigma_n) \cap \mathcal{I}^{\tilde{\rho}}(\gamma_n), \mathcal{I}^{\tilde{q}}(\sigma_n \cup \gamma_n) \leq \mathcal{I}^{\tilde{q}}(\sigma_n) \cup \mathcal{I}^{\tilde{q}}(\gamma_n) \text{ and } \mathcal{I}^{\tilde{\eta}}(\sigma_n \cup \gamma_n) \leq \mathcal{I}^{\tilde{\eta}}(\sigma_n) \cup \mathcal{I}^{\tilde{\eta}}(\gamma_n), \text{ for each } \sigma_n, \gamma_n \in \zeta^{\tilde{X}}.$$

The triple $(\tilde{X}, \tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is called a single valued neutrosophic ideal topological space in Šostak's sense (SVNITS, for short).

Definition 10 ([14]). Let $(\tilde{X}, \tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS for each $\sigma_n \in \zeta^{\tilde{X}}$. Then, the single valued neutrosophic ideal open local function $[\sigma_n]_r^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ of σ_n is the union of all single-valued neutrosophic points $x_{s,t,k}$ such that, if $\gamma_n \in Q_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(x_{s,t,k}, r)$ and $\mathcal{I}^{\tilde{\rho}}(\zeta_n) \geq r$, $\mathcal{I}^{\tilde{q}}(\zeta_n) \leq 1 - r$, $\mathcal{I}^{\tilde{\eta}}(\zeta_n) \leq 1 - r$, then there is at least one $v \in \tilde{X}$ for which $\tilde{\rho}_{\sigma_n}(v) + \tilde{\rho}_{\gamma_n}(v) - 1 > \tilde{\rho}_{\zeta_n}(v)$, $\tilde{q}_{\sigma_n}(v) + \tilde{q}_{\gamma_n}(v) - 1 \leq \tilde{q}_{\zeta_n}(v)$, and $\tilde{\eta}_{\sigma_n}(v) + \tilde{\eta}_{\gamma_n}(v) - 1 \leq \tilde{\eta}_{\zeta_n}(v)$.

Occasionally, we will write $[\sigma_n]_r^{\tilde{\rho}\tilde{q}\tilde{\eta}}$ for $[\sigma_n]_r^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$, and it will cause no ambiguity.

Remark 1 ([14]). Let $(\tilde{X}, \tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS and $\sigma_n \in \zeta^{\tilde{X}}$. Then,

$$\text{Cl}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r) = \sigma_n \cup [\sigma_n]_r^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \quad \text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r) = \sigma_n \cap [(\sigma_n^c)_r^{\tilde{\rho}\tilde{q}\tilde{\eta}}]^c.$$

It is clear that $\text{Cl}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}$ is a single-valued neutrosophic closure operator and $(\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is the single-valued neutrosophic topology generated by $\text{Cl}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}$, i.e.,

$$\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\mathcal{I})(\sigma_n) = \bigcup \{r \mid \text{Cl}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n^c, r) = \sigma_n^c\}.$$

Theorem 1 ([14]). Let $\{[\sigma_n]_i\}_{i \in J} \subset \zeta^{\tilde{X}}$ be a family of single-valued neutrosophic sets on \tilde{X} and $(\tilde{X}, \tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an r -SVNITS. Then,

- (1) $(\bigcup ([\sigma_n]_i)_r^{\tilde{\rho}\tilde{q}\tilde{\eta}} : i \in J) \leq (\bigcup [\sigma_n]_i : i \in j)_r^{\tilde{\rho}\tilde{q}\tilde{\eta}}$,
- (2) $(\bigcap ([\sigma_n]_i) : i \in j)_r^{\tilde{\rho}\tilde{q}\tilde{\eta}} \geq (\bigcap ([\sigma_n]_i)_r^{\tilde{\rho}\tilde{q}\tilde{\eta}} : i \in J)$.

Theorem 2 ([14]). Let $(\tilde{X}, \tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS and $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$, $r \in \zeta_0$. Then,

- (1) $\text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n \vee \gamma_n, r) \leq \text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r) \vee \text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\gamma_n, r)$,
- (2) $\text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r) \leq \text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r) \leq \sigma_n \leq \text{Cl}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r) \leq \text{C}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r)$,
- (3) $\text{Cl}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}([\sigma_n]^c, r) = [\text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r)]^c$, and $[\text{Cl}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r)]^c = \text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}([\sigma_n]^c, r)$,
- (4) $\text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n \wedge \gamma_n, r) = \text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r) \wedge \text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\gamma_n, r)$.

3. \mathcal{L} -Single Valued Neutrosophic Ideal Irresolute Mapping

This section provides the definitions of the r -single-valued neutrosophic \mathcal{L} -open set (SVN \mathcal{L} O, for short), the r -single-valued neutrosophic \mathcal{L} -closed set (SVN \mathcal{L} C, for short) and the \mathcal{L} -single valued neutrosophic ideal irresolute mapping (\mathcal{L} -SVNI-irresolute, for short), in the sense of Šostak. To understand the aim of this section, it is essential to clarify its content and elucidate the context in which the definitions, theorems, and examples are performed. Some results follow.

Definition 11. Let $(\tilde{X}, \tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an r -SVNITS for every $\sigma_n \in \zeta^{\tilde{X}}$ and $r \in \zeta_0$. Then, σ_n is called r -SVN \mathcal{L} C iff $\text{Cl}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r) = \sigma_n$. The complement of the r -SVN \mathcal{L} C is called r -SVN \mathcal{L} O.

Proposition 1. Let $(\tilde{X}, \tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an r -SVNITS and $\sigma_n \in \zeta^{\tilde{X}}$. Then,

- (1) σ_n is r -SVN \mathcal{L} C iff $[\sigma_n]_r^{\tilde{\rho}\tilde{q}\tilde{\eta}} \leq \sigma_n$,
- (2) σ_n is r -SVN \mathcal{L} O iff $([\sigma_n]_r^{\tilde{\rho}\tilde{q}\tilde{\eta}})^c \geq [\sigma_n]^c$,
- (3) If $\tau^{\tilde{\rho}}([\sigma_n]^c) \geq r$, $\tau^{\tilde{q}}([\sigma_n]^c) \leq 1 - r$, $\tau^{\tilde{\eta}}([\sigma_n]^c) \leq 1 - r$, then σ_n is r -SVN \mathcal{L} C,
- (4) If $\tau^{\tilde{\rho}}(\sigma_n) \geq r$, $\tau^{\tilde{q}}(\sigma_n) \leq 1 - r$, $\tau^{\tilde{\eta}}(\sigma_n) \leq 1 - r$, then σ_n is r -SVN \mathcal{L} O,
- (5) If σ_n is r -SVN \mathcal{L} C (resp. r -SVN β C), then $\text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}([\sigma_n]_r^{\tilde{\rho}\tilde{q}\tilde{\eta}}, r) \leq \sigma_n$ (resp. $\text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}([\text{int}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r)]_r^{\tilde{\rho}\tilde{q}\tilde{\eta}}) \leq \sigma_n$).

Proof. The proof of (1) and (2) are straightforward from Definition 11.

(3) Let $\tau^{\tilde{\rho}}([\sigma_n]^c) \geq r$, $\tau^{\tilde{q}}([\sigma_n]^c) \leq 1 - r$, $\tau^{\tilde{\eta}}([\sigma_n]^c) \leq 1 - r$. Then,

$$\sigma_n = \text{C}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r) \geq \text{Cl}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}\tilde{q}\tilde{\eta}}(\sigma_n, r) = \sigma_n \cup [\sigma_n]_r^{\tilde{\rho}\tilde{q}\tilde{\eta}} \geq [\sigma_n]_r^{\tilde{\rho}\tilde{q}\tilde{\eta}}.$$

Hence, σ_n is an r -SVN $\mathcal{L}C$.

(4) The proof is direct consequence of (1).

(5) Let σ_n be an r -SVN $\mathcal{S}C$. Then,

$$\begin{aligned} \sigma_n \geq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\sigma_n, r), r) &\geq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r) = \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}([\sigma_n \cup [\sigma_n]_r^{\tilde{\rho}}], r) \\ &\geq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}([\sigma_n]_r^{\tilde{\rho}}, r). \end{aligned}$$

The another case is similarly proved. \square

Example 1. Suppose that $\tilde{X} = \{a, b\}$. Define $\varepsilon_n, \gamma_n, \zeta_n \in \zeta^{\tilde{X}}$ as follows:

$$\gamma_n = \langle (0.3, 0.3), (0.3, 0.3), (0.3, 0.3) \rangle; \quad \varepsilon_n = \langle (0.7, 0.7), (0.7, 0.7), (0.7, 0.7) \rangle;$$

$$\zeta_n = \langle (0.2, 0.2), (0.2, 0.2), (0.2, 0.2) \rangle.$$

Define $\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}} : \zeta^{\tilde{X}} \rightarrow \zeta$ as follows:

$$\tilde{\tau}^{\tilde{\rho}}(\sigma_n) = \begin{cases} 1, & \text{if } \sigma_n = \tilde{0}, \\ 1, & \text{if } \sigma_n = \tilde{1}, \\ \frac{1}{3}, & \text{if } \sigma_n = \gamma_n; \\ \frac{1}{3}, & \text{if } \sigma_n = \varepsilon_n; \\ 0, & \text{if otherwise;} \end{cases} \quad \mathcal{I}^{\tilde{\rho}}(\sigma_n) = \begin{cases} 1, & \text{if } \sigma_n = (0, 1, 1), \\ \frac{1}{3}, & \text{if } \sigma_n = \zeta_n, \\ \frac{2}{3}, & \text{if } \tilde{0} < \sigma_n < \zeta_n; \\ 0, & \text{if otherwise;} \end{cases}$$

$$\tilde{\tau}^{\tilde{q}}(\sigma_n) = \begin{cases} 0, & \text{if } \sigma_n = \tilde{0}, \\ 0, & \text{if } \sigma_n = \tilde{1}, \\ \frac{2}{3}, & \text{if } \sigma_n = \gamma_n; \\ \frac{2}{3}, & \text{if } \sigma_n = \varepsilon_n; \\ 1, & \text{if otherwise;} \end{cases} \quad \mathcal{I}^{\tilde{q}}(\sigma_n) = \begin{cases} 0, & \text{if } \sigma_n = (0, 1, 1), \\ \frac{2}{3}, & \text{if } \sigma_n = \zeta_n, \\ \frac{1}{3}, & \text{if } \tilde{0} < \sigma_n < \zeta_n; \\ 1, & \text{if otherwise;} \end{cases}$$

$$\tilde{\tau}^{\tilde{\eta}}(\sigma_n) = \begin{cases} 0, & \text{if } \sigma_n = \tilde{0}, \\ 0, & \text{if } \sigma_n = \tilde{1}, \\ \frac{2}{3}, & \text{if } \sigma_n = \gamma_n; \\ \frac{2}{3}, & \text{if } \sigma_n = \varepsilon_n; \\ 1, & \text{if otherwise;} \end{cases} \quad \mathcal{I}^{\tilde{\eta}}(\sigma_n) = \begin{cases} 0, & \text{if } \sigma_n = (0, 1, 1), \\ \frac{2}{3}, & \text{if } \sigma_n = \zeta_n, \\ \frac{1}{3}, & \text{if } \tilde{0} < \sigma_n < \zeta_n; \\ 1, & \text{if otherwise.} \end{cases}$$

- (1) $G_n = \langle (0.6, 0.6), (0.6, 0.6), (0.6, 0.6) \rangle$ is $\frac{1}{3}$ -SVN $\mathcal{L}C$ but $\tilde{\tau}^{\tilde{\rho}}([G_n]^c) \not\geq \frac{1}{3}$, $\tilde{\tau}^{\tilde{q}}([G_n]^c) \not\leq \frac{2}{3}$, and $\tilde{\tau}^{\tilde{\eta}}([G_n]^c) \not\leq \frac{2}{3}$,
- (2) $G_n = \langle (0.6, 0.6), (0.6, 0.6), (0.6, 0.6) \rangle \geq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}([G_n]_{\frac{1}{3}}^{\tilde{\rho}}, \frac{1}{3}) = \tilde{0}$ but G_n is not is $\frac{1}{3}$ -SVN $\mathcal{S}C$.

Lemma 1. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS. Then, we have the following.

- (1) Every intersection of r -SVN $\mathcal{L}C$'s is r -SVN $\mathcal{L}C$.
- (2) Every union of r -SVN $\mathcal{L}O$'s is r -SVN $\mathcal{L}O$.

Proof. (1) Let $\{[\sigma_n]_i\}_{i \in j}$ be a family of r -SVN $\mathcal{L}C$'s. Then, for every $i \in j$, we obtain $[\sigma_n]_i = \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}}([\sigma_n]_i, r)$, and, by Theorem 1(2), we have

$$\begin{aligned} \bigcap_{i \in j} [\sigma_n]_i &= \bigcap_{i \in j} \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}}([\sigma_n]_i, r) = \bigcap_{i \in j} ([\sigma_n]_i \cup ([\sigma_n]_i)_r^{\tilde{\rho}}) \geq \bigcap_{i \in j} [\sigma_n]_i \cup \bigcap_{i \in j} ([\sigma_n]_i)_r^{\tilde{\rho}} \\ &\geq \bigcap_{i \in j} [\sigma_n]_i \cup \left(\bigcap_{i \in j} [\sigma_n]_i \right)_r^{\tilde{\rho}} = \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{\rho}}\left(\bigcap_{i \in j} [\sigma_n]_i, r\right). \end{aligned}$$

Therefore, $\bigcap_{i \in j} [\sigma_n]_i$ is r -SVN $\mathcal{L}C$.

(2) From Theorem 1(1). \square

Lemma 2. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ be an SVNITS for each $r \in \zeta_0$. Then,

- (1) For each r -SVNLO $\sigma_n \in \zeta^{\tilde{X}}$, $\sigma_n q \gamma_n$ iff $\sigma_n q \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r)$,
- (2) $x_{s,t,k} q \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r)$ iff $\sigma_n q \gamma_n$ for every r -SVNLO $\sigma_n \in \zeta^{\tilde{X}}$ with $x_{s,t,k} \in \sigma_n$.

Proof. (1) Let σ_n be an r -SVNLO and $\sigma_n \bar{q} \gamma_n$. Then, for any $v \in \tilde{X}$, we obtain

$$\tilde{\rho}_{\sigma_n}(v) + \tilde{\rho}_{\gamma_n}(v) > 1, \quad \tilde{\sigma}_{\sigma_n}(v) + \tilde{\sigma}_{\gamma_n}(v) \leq 1, \quad \tilde{\eta}_{\sigma_n}(v) + \tilde{\eta}_{\gamma_n}(v) \leq 1.$$

This implies that $\tilde{\rho}_{\gamma_n} \leq \tilde{\rho}_{[\sigma_n]^c}$, $\tilde{\sigma}_{\gamma_n} \geq \tilde{\sigma}_{[\sigma_n]^c}$ and $\tilde{\eta}_{\gamma_n} \geq \tilde{\eta}_{[\sigma_n]^c}$; hence, $\gamma_n \leq [\sigma_n]^c$. Since σ_n is r -SVNLO, $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r) \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}([\sigma_n]^c, r) = [\sigma_n]^c$, it follows that $\sigma_n \bar{q} \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r)$.

(2) Let $x_{s,t,k} q \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r)$. Then, $\sigma_n q \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r)$ with $x_{s,t,k} \in \sigma_n$. By (1), we have $\gamma_n q \sigma_n$ for each r -SVNLO $\sigma_n \in \zeta^{\tilde{X}}$. On the other hand, let $\sigma_n \bar{q} \gamma_n$. Then, $\gamma_n \leq [\sigma_n]^c$. Since σ_n is r -SVNLO,

$$\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r) \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}([\sigma_n]^c, r) = [\sigma_n]^c \quad \text{and} \quad \sigma_n \bar{q} \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r).$$

Since $x_{s,t,k} \in \sigma_n$, we obtain $x_{s,t,k} \bar{q} \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r)$ \square

Definition 12. Suppose that $f : (\tilde{X}, \tilde{\tau}_1^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}_1^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}) \rightarrow (\tilde{Y}, \tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is a mapping. Then,

- (1) f is called \mathcal{L} -SVNI-irresolute iff $f^{-1}(\sigma_n)$ is r -SVNLO in \tilde{X} for any r -SVNLO σ_n in \tilde{Y} ,
- (2) f is called \mathcal{L} -SVNI-irresolute open iff $f(\sigma_n)$ is r -SVNLO in \tilde{Y} for any r -SVNLO σ_n in \tilde{X} ,
- (3) f is called \mathcal{L} -SVNI-irresolute closed iff $f(\sigma_n)$ is r -SVNLC in \tilde{Y} for any r -SVNLC σ_n in \tilde{X} .

Theorem 3. Let $f : (\tilde{X}, \tilde{\tau}_1^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}_1^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}) \rightarrow (\tilde{Y}, \tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ be a mapping. Then, the following conditions are equivalent:

- (1) f is \mathcal{L} -SVNI-irresolute,
- (2) $f^{-1}(\sigma_n)$ is r -SVNLC, for each r -SVNLC $\sigma_n \in \tilde{Y}$,
- (3) $f(\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\sigma_n, r)) \leq \text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(\sigma_n), r)$ for each $\sigma_n \in \zeta^{\tilde{X}}$, $r \in \zeta_0$,
- (4) $\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f^{-1}(\gamma_n), r) \leq f^{-1}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r))$ for each $\gamma_n \in \zeta^{\tilde{Y}}$, $r \in \zeta_0$.

Proof. (1) \Rightarrow (2): Let σ_n be an r -SVNLC in \tilde{Y} . Then, $[\sigma_n]^c$ is r -SVNLO in \tilde{Y} by (1), we obtain $f^{-1}([\sigma_n]^c)$ is r -SVNLO. But, $f^{-1}([\sigma_n]^c) = [f^{-1}(\sigma_n)]^c$. Then, $f^{-1}(\sigma_n)$ is r -SVNLC in \tilde{X} .

(2) \Rightarrow (3): For each $\sigma_n \in \zeta^{\tilde{X}}$ and $r \in \zeta_0$, since $\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(\sigma_n), r)) = \text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(\sigma_n), r)$. From Definition 11, $\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(\sigma_n), r)$ is r -SVNLC in \tilde{Y} . By (2), $f^{-1}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(\sigma_n), r))$ is r -SVNLC in \tilde{X} . Since

$$\sigma_n \leq f^{-1}(f(\sigma_n)) \leq f^{-1}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(\sigma_n), r)),$$

by Definition 11, we get,

$$\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\sigma_n, r) \leq \text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f^{-1}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(\sigma_n), r)), r) = f^{-1}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(\sigma_n), r)).$$

Hence,

$$f(\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\sigma_n, r)) \leq f(f^{-1}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(\sigma_n), r))) \leq \text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(\sigma_n), r).$$

(3) \Rightarrow (4): For each $\gamma_n \in \zeta^{\tilde{Y}}$ and $r \in \zeta_0$, put $\sigma_n = f^{-1}(\gamma_n)$. By (3),

$$f(\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f^{-1}(\gamma_n), r)) \leq \text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(f(f^{-1}(\gamma_n)), r) \leq \text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}(\gamma_n, r).$$

It implies that $CI_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f^{-1}(\gamma_n), r) \leq f^{-1}(CI_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\gamma_n, r))$.

(4)⇒(1): Let γ_n be an r -SVN $\mathcal{L}\mathcal{O}$ in \tilde{Y} . Then, $[\gamma_n]^c$ is an r -SVN $\mathcal{L}\mathcal{C}$ in \tilde{Y} . Hence, $CI_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}([\gamma_n]^c, r) = [\gamma_n]^c$, and, by (4), we have,

$$f^{-1}([\gamma_n]^c) = f^{-1}(CI_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}([\gamma_n]^c, r)) \geq CI_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f^{-1}([\gamma_n]^c), r).$$

On the other hand, $f^{-1}([\gamma_n]^c) \leq CI_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f^{-1}([\gamma_n]^c), r)$. Thus, $f^{-1}([\gamma_n]^c) = CI_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f^{-1}([\gamma_n]^c), r)$, that is $f^{-1}([\gamma_n]^c)$ is an r -SVN $\mathcal{L}\mathcal{C}$ set in \tilde{X} . Hence, $f^{-1}(\gamma_n)$ is an r -SVN $\mathcal{L}\mathcal{O}$ set in \tilde{X} . □

Theorem 4. Let $f : (\tilde{X}, \tilde{\tau}_1^{\tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}, \mathcal{I}_1^{\tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}) \rightarrow (\tilde{Y}, \tilde{\tau}_2^{\tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}, \mathcal{I}_2^{\tilde{\rho}, \tilde{\sigma}, \tilde{\eta}})$ be a mapping. Then, the following conditions are equivalent:

- (1) f is \mathcal{L} -SVNI-irresolute open,
- (2) $f(\text{int}_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\sigma_n, r)) \leq \text{int}_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f(\sigma_n), r)$ for each $\sigma_n \in \tilde{\zeta}^{\tilde{X}}$, $r \in \tilde{\zeta}_0$,
- (3) $\text{int}_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f^{-1}(\gamma_n), r) \leq f^{-1}(\text{int}_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\gamma_n, r))$ for each $\gamma_n \in \tilde{\zeta}^{\tilde{Y}}$, $r \in \tilde{\zeta}_0$,
- (4) For any $\gamma_n \in \tilde{\zeta}^{\tilde{Y}}$ and any r -SVN $\mathcal{L}\mathcal{C}$ $\sigma_n \in \tilde{\zeta}^{\tilde{X}}$ with $f^{-1}(\gamma_n) \leq \sigma_n$, there exists an r -SVN $\mathcal{L}\mathcal{C}$ $\zeta_n \in \tilde{\zeta}^{\tilde{Y}}$ with $\gamma_n \leq \zeta_n$ such that $f^{-1}(\zeta_n) \leq \sigma_n$.

Proof. (1)⇒(2): For every $\sigma_n \in \tilde{\zeta}^{\tilde{X}}$, $r \in \tilde{\zeta}_0$ and $\text{int}_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\sigma_n, r) \leq \sigma_n$ from Theorem 2(2), we have $f(\text{int}_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\sigma_n, r)) \leq f(\sigma_n)$. By (1), $f(\text{int}_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\sigma_n, r))$ is r -SVN $\mathcal{L}\mathcal{O}$ in \tilde{Y} . Hence,

$$f(\text{int}_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\sigma_n, r)) = \text{int}_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f(\text{int}_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\sigma_n, r))) \leq \text{int}_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f(\sigma_n), r).$$

(2)⇒(3): For each $\gamma_n \in \tilde{\zeta}^{\tilde{Y}}$ and $r \in \tilde{\zeta}_0$, put $\sigma_n = f^{-1}(\gamma_n)$ from (2),

$$f(\text{int}_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f^{-1}(\gamma_n), r)) \leq \text{int}_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f(f^{-1}(\gamma_n)), r) \leq \text{int}_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\gamma_n, r).$$

It implies that

$$\text{int}_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f^{-1}(\gamma_n), r) \leq f^{-1}(f(\text{int}_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f^{-1}(\gamma_n), r))) \leq f^{-1}(\text{int}_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\gamma_n, r)).$$

(3)⇒(4): Obvious.

(4)⇒(1): Let ε_n be an r -SVN $\mathcal{L}\mathcal{O}$ in \tilde{X} . Put $\gamma_n = [f(\varepsilon_n)]^c$ and $\sigma_n = [\varepsilon_n]^c$ such that σ_n is r -SVN $\mathcal{L}\mathcal{C}$ in \tilde{X} . We obtain

$$f^{-1}(\gamma_n) = f^{-1}([f(\varepsilon_n)]^c) = [f^{-1}(f(\varepsilon_n))]^c \leq [\varepsilon_n]^c = \sigma_n.$$

From (4), there exists r -SVN $\mathcal{L}\mathcal{O}$ $\zeta_n \in \tilde{\zeta}^{\tilde{Y}}$ with $\gamma_n \leq \zeta_n$ such that $f^{-1}(\zeta_n) \leq \sigma_n = [\varepsilon_n]^c$. It implies $\varepsilon_n \leq [f^{-1}(\zeta)]^c = f^{-1}([\zeta_n]^c)$. Thus, $f(\varepsilon_n) \leq f(f^{-1}([\zeta_n]^c)) \leq [\zeta_n]^c$. On the other hand, since $\gamma_n \leq \zeta_n$, we have

$$f(\varepsilon_n) = [\gamma_n]^c \geq [\zeta_n]^c.$$

Hence, $f(\varepsilon_n) = [\zeta_n]^c$, that is, $f(\varepsilon_n)$ is r -SVN $\mathcal{L}\mathcal{O}$ in \tilde{Y} . □

Theorem 5. Let $f : (\tilde{X}, \tilde{\tau}_1^{\tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}, \mathcal{I}_1^{\tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}) \rightarrow (\tilde{Y}, \tilde{\tau}_2^{\tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}, \mathcal{I}_2^{\tilde{\rho}, \tilde{\sigma}, \tilde{\eta}})$ be a mapping. Then, the following conditions are equivalent:

- (1) f is \mathcal{L} -SVNI-irresolute closed.
- (2) $f(CI_{\tau_1}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(\gamma_n, r)) \leq CI_{\tau_2}^{\mathcal{L}, \tilde{\rho}, \tilde{\sigma}, \tilde{\eta}}(f(\gamma_n), r)$ for each $\gamma_n \in \tilde{\zeta}^{\tilde{X}}$, $r \in \tilde{\zeta}_0$.

Proof. Obvious. \square

Theorem 6. Let $f : (\tilde{X}, \tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}) \rightarrow (\tilde{Y}, \tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be a bijective mapping. Then, the following conditions are equivalent:

- (1) f is \mathcal{L} -SVNI-irresolute closed,
- (2) $\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(f^{-1}(\sigma_n), r) \leq f^{-1}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r))$ for each $\sigma_n \in \zeta^{\tilde{Y}}$, $r \in \zeta_0$.

Proof. (1) \Rightarrow (2) : Suppose that f is an \mathcal{L} -SVNI-irresolute closed. From Theorem 5(2), we claim that, for each $\gamma_n \in \zeta^{\tilde{X}}$ and $r \in \zeta_0$,

$$f(\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\gamma_n, r)) \leq \text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(f(\gamma_n), r).$$

Now, for all $\sigma_n \in \zeta^{\tilde{Y}}$, $r \in \zeta_0$, put $\gamma_n = f^{-1}(\sigma_n)$, since f is onto, it implies that $f(f^{-1}(\sigma_n)) = \sigma_n$. Thus,

$$f(\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(f^{-1}(\sigma_n), r)) \leq \text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(f(f^{-1}(\sigma_n)), r) = \text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r).$$

Again, since f is onto, it follows:

$$\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(f^{-1}(\sigma_n), r) = f^{-1}(f(\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(f^{-1}(\sigma_n), r))) \leq f^{-1}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)).$$

(2) \Rightarrow (1) : Put $\sigma_n = f(\gamma_n)$. By the injection of f , we get

$$\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\gamma_n, r) = \text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(f^{-1}(f(\gamma_n)), r) \leq f^{-1}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(f(\gamma_n), r)),$$

for the reason that f is onto, which implies that

$$f(\text{CI}_{\tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\gamma_n, r)) \leq f(f^{-1}(\text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(f(\gamma_n), r))) = \text{CI}_{\tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(f(\gamma_n), r).$$

\square

4. \mathcal{L} -Single Valued Neutrosophic Extremely Disconnected and \mathcal{L} -Single Valued Neutrosophic Normal

This section is devoted to introducing \mathcal{L} -single valued neutrosophic extremally disconnected (\mathcal{L} -SVNE-disconnected, for short) and \mathcal{L} -single valued neutrosophic normal (\mathcal{L} -SVN-normal, for short), in the sense of Šostak. These definitions and their components, together with a set of criteria for identifying the spaces, are provided to illustrate how the ideas are applied.

Definition 13. An SVNITS $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is called \mathcal{L} -SVNE-disconnected if $\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \geq r$, $\tilde{\tau}^{\tilde{q}}(\text{CI}_{\tilde{\tau}^{\tilde{q}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r$ for each $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \geq r$, $\tilde{\tau}^{\tilde{q}}(\sigma_n) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\sigma_n) \leq 1 - r$.


Definition 14. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS and $r \in \zeta_0$. Then, $\sigma_n \in \zeta^{\tilde{X}}$ is said to be:

- (1) r -single valued neutrosophic semi-ideal open set (r -SVNSIO) iff $\sigma_n \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\sigma_n, r), r)$,
- (2) r -single valued neutrosophic pre-ideal open set (r -SVNPIO) iff $\sigma_n \leq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r)$,
- (3) r -single valued neutrosophic α -ideal open set (r -SVN α IO) iff $\sigma_n \leq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\sigma_n, r), r), r)$,
- (4) r -single valued neutrosophic β -ideal open set (r -SVN β IO) iff $\sigma_n \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r), r)$,
- (5) r -single valued neutrosophic β -ideal open (r -SVN β IO) iff $\sigma_n \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r), r)$,

- (6) r -single valued neutrosophic regular ideal open set (r -SVNRIO) iff $\sigma_n = \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r)$.

The complement of r -SVNSIO (resp. r -SVNPIO, r -SVN α IO, r -SVN β IO, r -SVNS β IO, r -SVNRIO) are called r -SVNSIC (resp. r -SVNPIC, r -SVN α IC, r -SVN β IC, r -SVNS β IC, r -SVNRIC).

Remark 2. The following diagram can be easily obtained from the above definition:



Theorem 7. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ be an SVNITS and $r \in \zeta_0$. Then, the following properties are equivalent:

- (1) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is \mathcal{L} -SVNE-disconnected,
- (2) $\tilde{\tau}^{\tilde{\rho}}([\text{int}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{\rho}}(\sigma_n, r)]^c) \geq r$, $\tilde{\tau}^{\tilde{\sigma}}([\text{int}_{\tilde{\tau}^{\tilde{\sigma}}}^{\tilde{\sigma}}(\sigma_n, r)]^c) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}([\text{int}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{\eta}}(\sigma_n, r)]^c) \leq 1 - r$ for each $\tilde{\tau}^{\tilde{\rho}}([\sigma_n]^c) \geq r$, $\tilde{\tau}^{\tilde{\sigma}}([\sigma_n]^c) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}([\sigma_n]^c) \leq 1 - r$,
- (3) $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n, r), r) \leq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r)$, for each $\sigma_n \in \zeta^{\tilde{X}}$,
- (4) Every r -SVNSIO set is r -SVNPIO,
- (5) $\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{\rho}}(\sigma_n, r)) \geq r$, $\tilde{\tau}^{\tilde{\sigma}}(\text{CI}_{\tilde{\tau}^{\tilde{\sigma}}}^{\tilde{\sigma}}(\sigma_n, r)) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{\eta}}(\sigma_n, r)) \leq 1 - r$ for each r -SVNS β IO $\sigma_n \in \zeta^{\tilde{X}}$,
- (6) Every r -SVNS β IO set is r -SVNPIO,
- (7) For each $\sigma_n \in \zeta^{\tilde{X}}$, σ_n is r -SVN α IO set iff it is r -SVNSIO.

Proof. (1) \Rightarrow (2): The proof is direct consequence of Definition 14.

(2) \Rightarrow (3): For each $\sigma_n \in \zeta^{\tilde{X}}$, $\tilde{\tau}^{\tilde{\rho}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}}}(\sigma_n, r)) \geq r$, $\tilde{\tau}^{\tilde{\sigma}}(\text{int}_{\tilde{\tau}^{\tilde{\sigma}}}(\sigma_n, r)) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\text{int}_{\tilde{\tau}^{\tilde{\eta}}}(\sigma_n, r)) \leq 1 - r$, and, by (2), we have

$$\begin{aligned}
 \tilde{\tau}^{\tilde{\rho}}([\text{int}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{\rho}}([\text{int}_{\tilde{\tau}^{\tilde{\rho}}}(\sigma_n, r)]^c, r)]^c) &\geq r, & \tilde{\tau}^{\tilde{\sigma}}([\text{int}_{\tilde{\tau}^{\tilde{\sigma}}}^{\tilde{\sigma}}([\text{int}_{\tilde{\tau}^{\tilde{\sigma}}}(\sigma_n, r)]^c, r)]^c) &\leq 1 - r, \\
 \tilde{\tau}^{\tilde{\eta}}([\text{int}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{\eta}}([\text{int}_{\tilde{\tau}^{\tilde{\eta}}}(\sigma_n, r)]^c, r)]^c) &\leq 1 - r.
 \end{aligned}$$

Thus,

$$\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{\rho}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}}}(\sigma_n, r), r)) \geq r, \quad \tilde{\tau}^{\tilde{\sigma}}(\text{CI}_{\tilde{\tau}^{\tilde{\sigma}}}^{\tilde{\sigma}}(\text{int}_{\tilde{\tau}^{\tilde{\sigma}}}(\sigma_n, r), r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{\eta}}(\text{int}_{\tilde{\tau}^{\tilde{\eta}}}(\sigma_n, r), r)) \leq 1 - r;$$

hence,

$$\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n, r), r) = \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n, r), r), r) \leq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r).$$

(3) \Rightarrow (4): Let σ_n be an r -SVNSIO set. Then, by (4), we have

$$\sigma_n \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n, r), r) \leq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r).$$

Thus, σ_n is an r -SVNPIO set.

(4) \Rightarrow (5): Since σ_n is an r -SVNS β IO set, $\sigma_n \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r), r)$. Then, $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)$ is r -SVNSIO, and, by (4), $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \leq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r)$; hence, $\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{\rho}}(\sigma_n, r)) \geq r$, $\tilde{\tau}^{\tilde{\sigma}}(\text{CI}_{\tilde{\tau}^{\tilde{\sigma}}}^{\tilde{\sigma}}(\sigma_n, r)) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{\eta}}(\sigma_n, r)) \leq 1 - r$.

(5) \Rightarrow (6): Let σ_n be an r -SVN β IO set, then, by (5), $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \leq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r)$. Thus,

$$\sigma_n \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \leq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r).$$

Therefore, σ_n is an r -SVNPIO set.

(6) \Rightarrow (7): Let σ_n be an r -SVNSIO. Then, σ_n is r -SVNS β IO, by (6), σ_n is an r -SVNPIO set. Since σ_n is r -SVNSIO and r -SVNPIO, σ_n is r -SVN α IO.

(7) \Rightarrow (1): Suppose that $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \geq r$, $\tilde{\tau}^{\tilde{q}}(\sigma_n) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\sigma_n) \leq 1 - r$, then $CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)$ is r -SVNSIO, and, by (7), $CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)$ is r -SVN α IO. Hence,

$$CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r) \leq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(CI^*(\sigma_n, r), r), r), r) = \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r), r) \leq CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r).$$

Hence,

$$\tilde{\tau}^{\tilde{\rho}}(CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)) \geq r, \quad \tilde{\tau}^{\tilde{q}}(CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r.$$

Thus, $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is \mathcal{L} -SVNE-disconnected. \square

Theorem 8. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS $r \in \zeta_0$ and $\sigma_n \in \zeta^{\tilde{X}}$. Then, the following are equivalent:

- (1) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is \mathcal{L} -SVNE-disconnected,
- (2) $CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)\bar{q}C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\gamma_n, r)$, for every $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \geq r$, $\tilde{\tau}^{\tilde{q}}(\sigma_n) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\sigma_n) \leq 1 - r$ and every r -SVN \mathcal{L} O $\gamma_n \in \zeta^{\tilde{X}}$ with $\sigma_n\bar{q}\gamma_n$,
- (3) $CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r), r), r)\bar{q}C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\gamma_n, r)$, for every $\sigma_n \in \zeta^{\tilde{X}}$ and r -SVN \mathcal{L} O $\gamma_n \in \zeta^{\tilde{X}}$ with $\sigma_n\bar{q}\gamma_n$.

Proof. (1) \Rightarrow (2): Let $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \geq r$, $\tilde{\tau}^{\tilde{q}}(\sigma_n) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\sigma_n) \leq 1 - r$. Then, by (1),

$$\tilde{\tau}^{\tilde{\rho}}(CI_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\sigma_n, r)) \geq r, \quad \tilde{\tau}^{\tilde{q}}(CI_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(CI_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r.$$

Since $[CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)]^c$ is an r -SVN \mathcal{L} O and $CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)\bar{q}[CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)]^c$, it implies that

$$CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)\bar{q}C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}([CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)]^c, r).$$

(2) \Rightarrow (1): Let $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \geq r$, $\tilde{\tau}^{\tilde{q}}(\sigma_n) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\sigma_n) \leq 1 - r$. Since $[CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)]^c$ is an r -SVN \mathcal{L} O, then, by (2),

$$CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)\bar{q}C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}([CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)]^c, r).$$

This implies that $CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r) \leq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r), r) \leq CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)$, so

$$\tilde{\tau}^{\tilde{\rho}}(CI_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\sigma_n, r)) \geq r, \quad \tilde{\tau}^{\tilde{q}}(CI_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(CI_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r.$$

(2) \Rightarrow (3): Suppose that $\sigma_n \in \zeta^{\tilde{X}}$ and γ_n is an r -SVN \mathcal{L} O with $\sigma_n\bar{q}\gamma_n$. Since

$$\tilde{\tau}^{\tilde{\rho}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}}}(CI_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\sigma_n, r), r)) \geq r, \quad \tilde{\tau}^{\tilde{q}}(\text{int}_{\tilde{\tau}^{\tilde{q}}}(CI_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\sigma_n, r), r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(\text{int}_{\tilde{\tau}^{\tilde{\eta}}}(CI_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r), r)) \leq 1 - r.$$

By (2), we have $CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r), r), r)\bar{q}C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\gamma_n, r)$.

(3) \Rightarrow (2): Let $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \geq r$, $\tilde{\tau}^{\tilde{q}}(\sigma_n) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\sigma_n) \leq 1 - r$ and γ_n be an r -SVN \mathcal{L} O with $\sigma_n\bar{q}\gamma_n$. Then, by (3), we obtain $CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r), r), r)\bar{q}C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\gamma_n, r)$. Since

$$CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r) \leq CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\text{iny}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r), r), r),$$

then, we have $CI_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)\bar{q}C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\gamma_n, r)$. \square

Definition 15. An SVNITS $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is called \mathcal{L} -SVN-normal if, for every $[\sigma_n]_1\bar{q}[\sigma_n]_2$ with $\tilde{\tau}^{\tilde{\rho}}([\sigma_n]_1) \geq r$, $\tilde{\tau}^{\tilde{q}}([\sigma_n]_1) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}([\sigma_n]_1) \leq 1 - r$ and $[\sigma_n]_2$ is r -SVN \mathcal{L} O, there exists

$[\gamma_n]_j \in \xi^{\tilde{X}}$, for $j = \{1, 2\}$ with $\tilde{\tau}^{\tilde{\rho}}([\gamma_n]_1^c) \geq r$, $\tilde{\tau}^{\tilde{\rho}}([\gamma_n]_1^c) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}([\gamma_n]_1^c) \leq 1 - r$, $[\gamma_n]_2$ is r -SVN $\mathcal{L}C$ such that $[\sigma_n]_2 \leq [\gamma_n]_1$, $[\sigma_n]_1 \leq [\gamma_n]_2$ and $[\gamma_n]_1 \bar{q}[\gamma_n]_2$.

Theorem 9. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS; then, the following are equivalent:

- (1) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is an \mathcal{L} -SVN-normal.
- (2) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is an \mathcal{L} -SVNE-disconnected.

Proof. (1) \Rightarrow (2): Let $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \geq r$, $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\sigma_n) \leq 1 - r$ and $[\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)]^c$ be an r -SVN $\mathcal{L}O$. Then, $\sigma_n \bar{q}[\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)]^c$. By the \mathcal{L} -SVN-normality of $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$, there exist $[\gamma_n]_i \in \xi^{\tilde{X}}$, for $i = \{1, 2\}$ with

$$\tilde{\tau}^{\tilde{\rho}}([\gamma_n]_1^c) \geq r, \quad \tilde{\tau}^{\tilde{\rho}}([\gamma_n]_1^c) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}([\gamma_n]_1^c) \leq 1 - r,$$

and $[\gamma_n]_2$ r -SVN $\mathcal{L}C$ such that $[\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)]^c \leq [\gamma_n]_1$, $\sigma_n \leq [\gamma_n]_2$ and $[\gamma_n]_1 \bar{q}[\gamma_n]_2$. Since

$$\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r) \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}([\gamma_n]_2, r) = [\gamma_n]_2 \leq [\gamma_n]_1^c \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r),$$

we have $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r) = [\gamma_n]_2$. Since $[\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)]^c \leq [\gamma_n]_1 \leq [\gamma_n]_2^c = [\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)]^c$, so $[\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)]^c = [\gamma_n]_1$. Hence, $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r) = [\gamma_n]_1^c$ and

$$\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \geq r, \quad \tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r.$$

Thus, $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is an \mathcal{L} -SVNE-disconnected.

(2) \Rightarrow (1): Suppose that $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \geq r$, $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\sigma_n) \leq 1 - r$ and γ_n is an r -SVN $\mathcal{L}O$ with $\sigma_n \bar{q}\gamma_n$. By the \mathcal{L} -SVNE-disconnected of $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$, we have

$$\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \geq r, \quad \tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r,$$

and $[\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)]^c$ is r -SVN $\mathcal{L}O$. Since $\sigma_n \bar{q}\gamma_n$, $\sigma_n \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)$ and $\gamma_n \leq [\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)]^c$. Thus, $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is an \mathcal{L} -SVN-normal. \square

Theorem 10. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS, $\sigma_n, \sigma_n \xi^{\tilde{X}}$ and $r \in \xi_0$. Then, the following properties are equivalent:

- (1) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is \mathcal{L} -SVNE-disconnected.
- (2) If σ_n is r -SVNRIO, then σ_n is r -SVN $\mathcal{L}C$.
- (3) If σ_n is r -SVN $\mathcal{L}C$, then σ_n is r -SVN $\mathcal{L}O$.

Proof. (1) \Rightarrow (2): Let σ_n be an r -SVNRIO. Then, $\sigma_n = \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r)$ and $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \geq r$, $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\sigma_n) \leq 1 - r$. By (1),

$$\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \geq r, \quad \tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r.$$

Hence $\sigma_n = \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r) = \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)$.

(2) \Rightarrow (1): Suppose that $\sigma_n = \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r)$, then $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \geq r$, $\tilde{\tau}^{\tilde{\rho}}(\sigma_n) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\sigma_n) \leq 1 - r$, by (2), σ_n is r -SVN $\mathcal{L}C$. This implies that

$$\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r) \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r), r) = \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r) \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r).$$

Thus,

$$\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \geq r, \quad \tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r,$$

then $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is an \mathcal{L} -SVNE-disconnected.

(2) \Leftrightarrow (3): Obvious. \square

Remark 3. The union of two r -SVNRIO sets need not to be an r -SVNRIO.

Theorem 11. If $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is \mathcal{L} -SVNE-disconnected and $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}, r \in \zeta_0$. Then, the following properties hold:

- (1) If σ_n and γ_n are r -SVNRIC, then $\sigma_n \wedge \gamma_n$ is r -SVNRIC.
- (2) If σ_n and γ_n are r -SVNRIO, then $\sigma_n \vee \gamma_n$ is r -SVNRIO.

Proof. Let σ_n and γ_n be r -SVNRIC. Then, $\tilde{\tau}^{\tilde{\rho}}([\sigma_n]^c) \geq r, \tilde{\tau}^{\tilde{q}}([\sigma_n]^c) \leq 1 - r, \tilde{\tau}^{\tilde{\eta}}([\sigma_n]^c) \leq 1 - r$ and $\tilde{\tau}^{\tilde{\rho}}([\gamma_n]^c) \geq r, \tilde{\tau}^{\tilde{q}}([\gamma_n]^c) \leq 1 - r, \tilde{\tau}^{\tilde{\eta}}([\gamma_n]^c) \leq 1 - r$, by Theorem 7, we have

$$\tilde{\tau}^{\tilde{\rho}}([\text{int}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\sigma_n, r)]^c) \geq r, \quad \tilde{\tau}^{\tilde{q}}([\text{int}_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\sigma_n, r)]^c) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}([\text{int}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)]^c) \leq 1 - r,$$

and

$$\tilde{\tau}^{\tilde{\rho}}([\text{int}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\gamma_n, r)]^c) \geq r, \quad \tilde{\tau}^{\tilde{q}}([\text{int}_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\gamma_n, r)]^c) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}([\text{int}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\gamma_n, r)]^c) \leq 1 - r.$$

This implies that

$$\begin{aligned} \sigma_n \wedge \gamma_n &= C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r), r) \wedge C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\gamma_n, r), r) \\ &= \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r) \wedge \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\gamma_n, r) = \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n \wedge \gamma_n, r) \\ &\leq C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n \wedge \gamma_n, r), r). \end{aligned}$$

On the other hand,

$$\begin{aligned} C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n \wedge \gamma_n, r), r) &= C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r) \wedge \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\gamma_n, r), r) \\ &\leq C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r), r) \wedge C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\gamma_n, r), r) \\ &= \sigma_n \wedge \gamma_n. \end{aligned}$$

Thus, $C_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n \wedge \gamma_n, r), r) = \sigma_n \wedge \gamma_n$. Therefore, $\sigma_n \wedge \gamma_n$ is an r -SVNRIC.

(2) The proof is similar to that of (1). \square

Theorem 12. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS and $r \in \zeta_0$. Then, the following properties are equivalent:

- (1) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is \mathcal{L} -SVNE-disconnected,
- (2) $\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\sigma_n, r)) \geq r, \tilde{\tau}^{\tilde{q}}(\text{CI}_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r, \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r$, for every r -SVNSIO $\sigma_n \in \zeta^{\tilde{X}}$,
- (3) $\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\sigma_n, r)) \geq r, \tilde{\tau}^{\tilde{q}}(\text{CI}_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r, \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r$, for every r -SVNPIO $\sigma_n \in \zeta^{\tilde{X}}$,
- (4) $\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\sigma_n, r)) \geq r, \tilde{\tau}^{\tilde{q}}(\text{CI}_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r, \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r$, for every r -SVNRIO $\sigma_n \in \zeta^{\tilde{X}}$.

Proof. (1) \Rightarrow (2) and (1) \Rightarrow (3). Let σ_n be an r -SVNSIO (r -SVNPIO). Then, σ_n is r -SVNS β IO, and, by Theorem 7, we have,

$$\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\sigma_n, r)) \geq r, \quad \tilde{\tau}^{\tilde{q}}(\text{CI}_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r.$$

(2) \Rightarrow (4) and (3) \Rightarrow (4). Let σ_n be an r -SVNRIO. Then, σ_n is r -SVNPIO and r -SVNSIO. Thus,

$$\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\sigma_n, r)) \geq r, \quad \tilde{\tau}^{\tilde{q}}(\text{CI}_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r, \quad \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)) \leq 1 - r.$$

(4) \Rightarrow (1). Suppose that

$$\tilde{\tau}^{\tilde{\rho}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\tilde{L}}(\sigma_n, r), r)) \geq r, \quad \tilde{\tau}^{\tilde{q}}(\text{int}_{\tilde{\tau}^{\tilde{q}}}(\text{CI}_{\tilde{\tau}^{\tilde{q}}}^{\tilde{L}}(\sigma_n, r), r)) \geq r, \quad \tilde{\tau}^{\tilde{\eta}}(\text{int}_{\tilde{\tau}^{\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r), r)) \geq r.$$

Then, by (4), we have

$$\begin{aligned} \tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r), r), r)) \geq r, \quad \tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r), r), r)) \geq r, \\ \tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\mathcal{L}}(\text{int}_{\tilde{\tau}^{\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r), r)) \geq r. \end{aligned}$$

Hence,

$$\begin{aligned} \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r) &\leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}^{\mathcal{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r), r) \\ &= \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}^{\mathcal{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r), r), r) \\ &= \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r) \leq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r). \end{aligned}$$

Thus, $\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \geq r$, $\tilde{\tau}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}(\text{CI}_{\tilde{\tau}^{\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r)) \leq 1 - r$; hence, $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\eta}})$ is an \mathcal{L} -SVNE-disconnected. \square

Definition 16. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\eta}})$ be an SVNITS. Then, σ_n is said to be an r -SVN \mathcal{L} SO if $\sigma_n \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}^{\mathcal{L}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}^{\mathcal{L}}(\sigma_n, r), r)$.

Definition 17. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\eta}})$ be an SVNITS for each $r \in \zeta_0$, $\sigma_n \in \zeta^{\tilde{X}}$ and $x_{s,t,p} \in \text{Pt}(\zeta^{\tilde{X}})$. Then, $x_{s,t,p}$ is called an r -SVN $\delta\mathcal{I}$ -cluster point of σ_n if, for every $\gamma_n \in Q_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(x_{s,t,p}, r)$, we have $\sigma_n q \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}^{\mathcal{L}}(\gamma_n, r), r)$.

Definition 18. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\eta}})$ be an SVNITS for each $r \in \zeta_0$, $\sigma_n \in \zeta^{\tilde{X}}$ and $x_{s,t,p} \in \text{Pt}(\zeta^{\tilde{X}})$. Then, the single-valued neutrosophic $\delta\mathcal{I}$ -closure operator is a mapping $\text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}} : \zeta^{\tilde{X}} \times \zeta_0 \rightarrow \zeta^{\tilde{X}}$ that is defined as: $\text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r) = \bigvee \{x_{s,t,p} \in \text{Pt}(\zeta^{\tilde{X}}) \text{ is } r\text{-SVN}\delta\mathcal{I}\text{-cluster point of } \sigma_n\}$.

Lemma 3. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\eta}})$ be an SVNITS. Then, σ_n is r -SVN \mathcal{L} SO iff $\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r) = \text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)$.

Proof. Obvious. \square

Lemma 4. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\eta}})$ be an SVNITS for each $\sigma_n \in \zeta^{\tilde{X}}$ and $r \in \zeta_0$. Then, $\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r) \leq \text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)$.

Proof. Obvious. \square

Lemma 5. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\eta}})$ be an SVNITS and σ_n be an r -SVN \mathcal{L} SO. Then, $\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r) = \text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)$.

Proof. We show that $\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r) \leq \text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)$. Suppose that $\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r) \not\leq \text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)$; then, there exist $v \in \tilde{X}$ and $s, t, p \in \zeta_0$ such that

$$\begin{aligned} \tilde{\rho}_{\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v) < s \leq \tilde{\rho}_{\text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v), \quad \tilde{\rho}_{\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v) \geq t > \tilde{\rho}_{\text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v), \quad (1) \\ \tilde{\eta}_{\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v) \geq p > \tilde{\eta}_{\text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v). \end{aligned}$$

By the definition of $\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}$, there exists $\tilde{\tau}^{\tilde{\rho}}([\gamma_n]^c) \geq r$, $\tilde{\tau}^{\tilde{\rho}}([\gamma_n]^c) \leq 1 - r$, $\tilde{\tau}^{\tilde{\eta}}([\gamma_n]^c) \leq 1 - r$ with $\sigma_n \leq \gamma_n$ such that

$$\begin{aligned} \tilde{\rho}_{\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v) \leq \tilde{\rho}_{\gamma_n}(v) < s < \tilde{\rho}_{\text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v), \quad \tilde{\rho}_{\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v) \geq \tilde{\rho}_{\gamma_n}(v) > t > \tilde{\rho}_{\text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v), \\ \tilde{\eta}_{\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v) \geq \tilde{\rho}_{\gamma_n}(v) > p > \tilde{\eta}_{\text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\eta}}}(\sigma_n, r)}(v). \end{aligned}$$

Then, $[\gamma_n]^c \in Q_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(x_{s,t,p}, r)$ and

$$\begin{aligned} [\sigma_n]^c \geq [\gamma_n]^c &\Rightarrow \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}([\sigma_n]^c, r) \geq \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}([\gamma_n]^c, r) \\ &\Rightarrow \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}([\sigma_n]^c, r) \geq \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}([\gamma_n]^c, r) \\ &\Rightarrow [\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)]^c \geq [\gamma_n]^c. \end{aligned}$$

Thus, $\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \bar{q} [\gamma_n]^c$. Hence, $\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}([\gamma_n]^c, r), r) \bar{q} \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r)$. Since σ_n is an r -SVNLSO, we have $\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\gamma_n, r), r) \bar{q} \sigma_n$. So, $x_{s,t,p}$ is not an r -SVN $\delta\mathcal{I}$ -cluster point of σ_n . It is a contradiction for equation 3. Thus, $\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n, r) \geq \text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n, r)$. By Lemma 4, we have $\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n, r) = \text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n, r)$. \square

Theorem 13. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ be an SVNITS. Then, the following properties are equivalent:

- (1) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is \mathcal{L} -SVNE-disconnected,
- (2) If σ_n is r -SVNS β IO and γ_n is r -SVNLSO, then $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \wedge \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n, r) \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n \wedge \gamma_n)$,
- (3) If σ_n is r -SVNSIO and γ_n is r -SVNLSO, then $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \wedge \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n, r) \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n \wedge \gamma_n)$,
- (4) $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \bar{q} \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n, r)$, for every r -SVNSIO set $\sigma_n \in \xi^{\tilde{X}}$ and every r -SVNLSO $\gamma_n \in \xi^{\tilde{X}}$ with $\sigma_n \bar{q} \gamma_n$,
- (5) If σ_n is an r -SVNPIO and γ_n is an r -SVNLSO, then $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \wedge \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n, r) \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n \wedge \gamma_n)$.

Proof. (1) \Rightarrow (2): Let σ_n be an r -SVNS β IO and γ_n be an r -SVNLSO, by Theorem 7, $\tilde{\tau}^{\tilde{\rho}}$ ($\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)$) $\geq r$, $\tilde{\tau}^{\tilde{\rho}}$ ($\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)$) $\leq 1 - r$, $\tilde{\tau}^{\tilde{\rho}}$ ($\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)$) $\leq 1 - r$. Then,

$$\begin{aligned} \text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \wedge \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n, r) &\leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\gamma_n, r), r) \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\gamma_n, r) \wedge \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\gamma_n, r), r) \\ &\leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}[\gamma_n \wedge \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\gamma_n, r), r], r) \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}[\gamma_n \wedge \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\gamma_n, r), r], r) \\ &\leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n \wedge \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\gamma_n, r), r) \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n \wedge \gamma_n, r). \end{aligned}$$

Hence, $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \wedge \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n, r) \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n \wedge \gamma_n)$.

(2) \Rightarrow (3): It follows from the fact that every r -SVNSIO set is an r -SVNS β IO.

(3) \Rightarrow (4): Clear.

(4) \Rightarrow (1): Let σ_n be an r -SVNSIO. Since $[\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)]^c \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}([\sigma_n]^c, r), r)$ we have, $[\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)]^c$ is an r -SVNLSO. Then, by (4), $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \bar{q} \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}([\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)]^c, r)$. Thus, $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \leq [\text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)]^c, r) = \text{int}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r), r)$. Therefore, $\tilde{\tau}^{\tilde{\rho}}$ ($\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)$) $\geq r$, $\tilde{\tau}^{\tilde{\rho}}$ ($\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)$) $\leq 1 - r$, $\tilde{\tau}^{\tilde{\rho}}$ ($\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r)$) $\leq 1 - r$. Thus, by Theorem 12, $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is \mathcal{L} -SVNE-disconnected.

(2) \Rightarrow (5): It follows from the fact that every r -SVNPIO is an r -SVNS β IO. \square

Corollary 1. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ be an SVNITS. Then, the following properties are equivalent:

- (1) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is \mathcal{L} -SVNE-disconnected.
- (2) If σ_n is an r -SVNS β IO and γ_n is an r -SVNLSO, then $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \wedge \text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n, r) \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n \wedge \gamma_n)$.
- (3) If σ_n is an r -SVNSIO and γ_n is an r -SVNLSO, then $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \wedge \text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n, r) \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n \wedge \gamma_n)$.
- (4) If σ_n is an r -SVNPIO and γ_n is an r -SVNLSO, then $\text{CI}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\rho}}(\sigma_n, r) \wedge \text{C}_{\delta\mathcal{I}\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\gamma_n, r) \leq \text{C}_{\tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}(\sigma_n \wedge \gamma_n)$.

Proof. It follows directly from Lemma 3 and 5. \square

5. Some Types of Separation Axioms

In this section, some kinds of separation axioms, namely r -single valued neutrosophic ideal- R_i (r -SVNIR $_i$, for short), where $i = \{0, 1, 2, 3\}$, and r -single valued neutrosophic ideal- T_j (r -SVNIT $_j$, for short), where $j = \{1, 2, 2\frac{1}{2}, 3, 4\}$, in the sense of Šostak are defined. Some of their characterizations, fundamental properties, and the relations between these notions have been studied.

Definition 19. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS and $r \in \zeta_0$. Then, \tilde{X} is called:

- (1) r -SVNIR $_0$ iff $x_{s,t,p}\bar{q}CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(y_{s_1,t_1,p_1}, r)$ implies $y_{s_1,t_1,p_1}\bar{q}CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(x_{s,t,p}, r)$ for any $x_{s,t,p} \neq y_{s_1,t_1,p_1}$.
- (2) r -SVNIR $_1$ iff $x_{s,t,p}\bar{q}CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(y_{s_1,t_1,p_1}, r)$ implies that there exist r -SVNŁO sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n, y_{s_1,t_1,p_1} \in \gamma_n$ and $\sigma_n\bar{q}\gamma_n$.
- (3) r -SVNIR $_2$ iff $x_{s,t,p}\bar{q}\zeta_n = CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\zeta_n, r)$ implies there exist r -SVNŁO sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n, \zeta_n \leq \gamma_n$ and $\sigma_n\bar{q}\gamma_n$.
- (4) r -SVNIR $_3$ iff $[\zeta_n]_1 = CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}([\zeta_n]_1, r)\bar{q}[\zeta_n]_2 = CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}([\zeta_n]_2, r)$ implies that there exist r -SVNŁO sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $[\zeta_n]_1 \leq \sigma_n, [\zeta_n]_2 \leq \gamma_n$ and $\sigma_n\bar{q}\gamma_n$.
- (5) r -SVNIT $_1$ iff $x_{s,t,p}\bar{q}y_{s_1,t_1,p_1}$ implies that there exists r -SVNŁO $\sigma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n$ and $y_{s_1,t_1,p_1}\bar{q}\sigma_n$.
- (6) r -SVNIT $_2$ iff $x_{s,t,p}\bar{q}y_{s_1,t_1,p_1}$ implies that there exist r -SVNŁO sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n, y_{s_1,t_1,p_1} \in \gamma_n$ and $\sigma_n\bar{q}\gamma_n$.
- (7) r -SVNIT $_{2\frac{1}{2}}$ iff $x_{s,t,p}\bar{q}y_{s_1,t_1,p_1}$ implies that there exist r -SVNŁO sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n, y_{s_1,t_1,p_1} \in \gamma_n$ and $CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)\bar{q}CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\gamma_n, r)$.
- (8) r -SVNIT $_3$ iff it is r -SVNITR $_2$ and r -SVNIT $_1$.
- (9) r -SVNIT $_4$ iff it is r -SVNITR $_3$ and r -SVNIT $_1$.

Theorem 14. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS and $r \in \zeta_0$. Then, the following statements are equivalent:

- (1) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is r -SVNIR $_0$.
- (2) If $x_{s,t,p}\bar{q}\sigma_n = CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)$, then there exists r -SVNŁO $\gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p}\bar{q}\gamma_n$ and $\sigma_n \leq \gamma_n$.
- (3) If $x_{s,t,p}\bar{q}\sigma_n = CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)$, then $CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(x_{s,t,p}, r)\bar{q}\sigma_n = CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)$.
- (4) If $x_{s,t,p}\bar{q}CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(y_{s_1,t_1,p_1}, r)$, then $CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(x_{s,t,p}, r)\bar{q}CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(y_{s_1,t_1,p_1}, r)$.

Proof. (1) \Rightarrow (2): Let $x_{s,t,p}\bar{q}\sigma_n = CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)$. Then,

$$s + \tilde{\rho}\sigma_n(v) < 1, \quad t + \tilde{q}\sigma_n(v) \geq 1, \quad p + \tilde{\eta}\sigma_n(v) \geq 1,$$

for every $y_{s_1,t_1,p_1} \in \sigma_n$, we have $s_1 < \tilde{\rho}\sigma_n(v), t_1 \geq \tilde{q}\sigma_n(v)$ and $p_1 \geq \tilde{\eta}\sigma_n(v)$. Thus, $x_{s,t,p}\bar{q}CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(y_{s_1,t_1,p_1}, r)$. Since $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is an r -SVNIR $_0$, we obtain $y_{s_1,t_1,p_1}\bar{q}CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(x_{s,t,p}, r)$. By Lemma 2(2), there exists an r -SVNŁO $\zeta_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p}\bar{q}\zeta_n$ and $y_{s_1,t_1,p_1} \leq \zeta_n$. Let

$$\gamma_n = \bigvee_{y_{s_1,t_1,p_1} \in \sigma_n} \{\zeta_n : x_{s,t,p}\bar{q}\zeta_n, y_{s_1,t_1,p_1} \in \zeta_n\}.$$

From Lemma 1(1), γ_n is an r -SVNŁO. Then, $x_{s,t,p}\bar{q}\gamma_n, \sigma_n \leq \gamma_n$.

(2) \Rightarrow (3): Let $x_{s,t,p}\bar{q}\sigma_n = CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)$. Then, there exists an r -SVNŁO $\gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p}\bar{q}\gamma_n$ and $\sigma_n \leq \gamma_n$. Since for every $v \in \tilde{X}$,

$$s < 1 - \tilde{\rho}\gamma_n(v), \quad t \geq 1 - \tilde{q}\gamma_n(v), \quad p \geq 1 - \tilde{\eta}\gamma_n(v),$$

we obtain

$$CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(x_{s,t,p}, r) \leq CI_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}([\gamma_n]^c, r) = [\gamma_n]^c \leq [\sigma_n]^c.$$

Therefore, $\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(x_{s,t,p}, r)\bar{q}\sigma_n = \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(\sigma_n, r)$.

(3) \Rightarrow (4): Let $x_{s,t,p}\bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r)$. Then, $x_{s,t,p}\bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r) = \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r), r)$. By (3), $s_1, t_1, p_1(x_{s,t,p}, r)\bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r)$.

(4) \Rightarrow (1): Clear. \square

Theorem 15. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ be an SVNITS and $r \in \zeta_0$. Then, if \tilde{X} is

(1) $[r\text{-SVNIR}_3 \text{ and } r\text{-SVNIR}_0] \Rightarrow^{(a)} r\text{-SVNIR}_2 \Rightarrow^{(b)} r\text{-SVNIR}_1 \Rightarrow^{(c)} r\text{-SVNIR}_0$.

(2) $r\text{-SVNIT}_2 \Rightarrow r\text{-SVNIR}_1$.

(3) $r\text{-SVNIT}_3 \Rightarrow r\text{-SVNIR}_2$.

(4) $r\text{-SVNIT}_4 \Rightarrow r\text{-SVNIR}_3$.

(5) $r\text{-SVNIT}_4 \Rightarrow^{(a)} r\text{-SVNIT}_3 \Rightarrow^{(b)} r\text{-SVNIT}_{2\frac{1}{2}} \Rightarrow^{(c)} r\text{-SVNIT}_2 \Rightarrow^{(d)} r\text{-SVNIT}_1$.

Proof. (1a). Let $x_{s,t,p}\bar{q}\zeta_n = \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(\zeta_n, r)$, by Theorem 14(3), $\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(x_{s,t,p}, r)\bar{q}\zeta_n = \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(\zeta_n, r)$. Since $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is $r\text{-SVNIR}_3$ and $\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(x_{s,t,p}, r) = \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(x_{s,t,p}, r), r)$, there exist $r\text{-SVN}\mathcal{L}\mathcal{O}$ sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(x_{s,t,p}, r) \leq \sigma_n, \zeta_n \leq \gamma_n$ and $\sigma_n\bar{q}\gamma_n$. Hence, $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is $r\text{-SVNIR}_2$.

(1b). For each $x_{s,t,p}\bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r)$, by $r\text{-SVNIR}_2$ of \tilde{X} , there exist $r\text{-SVN}\mathcal{L}\mathcal{O}$ sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n, y_{s_1,t_1,p_1}, r \in \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r) \leq \gamma_n$ and $\sigma_n\bar{q}\gamma_n$. Thus, $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is $r\text{-SVNIR}_1$.

(1c). Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ be $r\text{-SVNIR}_1$. Then, for every $x_{s,t,p}\bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r, r)$ and $x_{s,t,p} \neq y_{s_1,t_1,p_1}$, there exist $r\text{-SVN}\mathcal{L}\mathcal{O}$ sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n, y_{s_1,t_1,p_1} \in \gamma_n$ and $\sigma_n\bar{q}\gamma_n$. Hence, $x_{s,t,p} \in \sigma_n \leq [\gamma_n]^c$. Since γ_n is an $r\text{-SVN}\mathcal{L}\mathcal{O}$ set, we obtain $\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(x_{s,t,p}, r) \leq \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}([\gamma_n]^c, r) = [\gamma_n]^c \leq [y_{s_1,t_1,p_1}]^c$. Thus, $y_{s_1,t_1,p_1}\bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(x_{s,t,p}, r)$ and $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is $r\text{-SVNIR}_0$.

(2). Let $x_{s,t,p}\bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_2,p_1}, r)$. Then, $x_{s,t,p}\bar{q}y_{s_1,t_1,p_1}$. By $r\text{-SVNIT}_2$ of \tilde{X} , there exist $r\text{-SVN}\mathcal{L}\mathcal{O}$ sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n, y_{s_1,t_1,p_1} \in \gamma_n$ and $\sigma_n\bar{q}\gamma_n$. Hence, $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is $r\text{-SVNIR}_1$.

(3) and (4) The proofs are direct consequence of (2).

(5a). The proof is direct consequence of (1).

(5b). For each $x_{s,t,p}\bar{q}y_{s_1,t_1,p_1}$, since \tilde{X} is both $r\text{-SVNIR}_2$ and $r\text{-SVNIT}_1$, then, there exists an $r\text{-SVN}\mathcal{L}\mathcal{O}$ set $\zeta_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \zeta_n$ and $y_{s_1,t_1,p_1}\bar{q}\zeta_n$. Then,

$$x_t \in \zeta_n = \text{int}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(\zeta_n, r) \leq \text{int}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}([y_{s_1,t_1,p_1}]^c, r) = [\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r)]^c.$$

Hence, $x_{s,t,p}\bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r)$. By $r\text{-SVNIR}_2$ of \tilde{X} , there exist $r\text{-SVN}\mathcal{L}\mathcal{O}$ sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n, \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r) \leq \gamma_n$ and $\sigma_n\bar{q}\gamma_n$. Thus, $\sigma_n \leq [\gamma_n]^c$, so

$$\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(\sigma_n, r) \leq \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}([\gamma_n]^c, r) = [\gamma_n]^c \leq [\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r)]^c.$$

It implies $\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(\sigma_n, r)\bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r)$ with $x_{s,t,p} \in \sigma_n$ and $y_{s_1,t_1,p_1} \in \text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(y_{s_1,t_1,p_1}, r)$. Thus, $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is $r\text{-SVNIT}_{2\frac{1}{2}}$.

(5c). Let $x_{s,t,p}\bar{q}y_{s_1,t_1,p_1}$. Then, by $r\text{-SVNIT}_{2\frac{1}{2}}$ of \tilde{X} , there exist $r\text{-SVN}\mathcal{L}\mathcal{O}$ sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n, y_{s_1,t_1,p_1} \in \gamma_n$ and $\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(\sigma_n, r)\bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}}^{\tilde{\epsilon}}(\gamma_n, r)$, which implies that $\sigma_n\bar{q}\gamma_n$. Thus, $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is $r\text{-SVNIT}_2$.

(5d). Similar to the proof of (5c). \square

Theorem 16. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ be an SVNITS and $r \in \zeta_0$. Then, the following statements are equivalent:

(1) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}}, \tilde{\mathcal{I}}^{\tilde{\rho}\tilde{\sigma}\tilde{\eta}})$ is $r\text{-SVNIR}_2$.

- (2) If $x_{s,t,p} \in \sigma_n$ and σ_n is r -SVN \mathcal{E} O set, then there exists r -SVN \mathcal{E} O set $\gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \gamma_n \leq \text{CI}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\gamma_n, r) \leq \sigma_n$.
- (3) If $x_{s,t,p} \bar{q}\sigma_n = \text{CI}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\sigma_n, r)$, then there exists r -SVN \mathcal{E} O set $[\gamma_n]_j \in \zeta^{\tilde{X}}$, $j = \{1, 2\}$ such that $x_{s,t,p} \in [\gamma_n]_1$, $\sigma_n \leq [\gamma_n]_2$ and $\text{CI}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}([\gamma_n]_1, r) \bar{q}\text{CI}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}([\gamma_n]_2, r)$.

Proof. Similar to the proof of Theorem 14. \square

Theorem 17. Let $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an SVNITS and $r \in \zeta_0$. Then, the following statements are equivalent:

- (1) $(\tilde{X}, \tilde{\tau}^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is r -SVNIR $_3$.
- (2) If $[\sigma_n]_1 \bar{q}[\sigma_n]_2$ and $[\sigma_n]_1, [\sigma_n]_2$ are r -SVN \mathcal{E} C sets, then there exists r -SVN \mathcal{E} O set $\gamma_n \in \zeta^{\tilde{X}}$ such that $[\sigma_n]_1 \leq \gamma_n$ and $\text{CI}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\gamma_n, r) \leq [\sigma_n]_2$.
- (3) For any $[\sigma_n]_1 \leq [\sigma_n]_2$, where $[\sigma_n]_1$ is an r -SVN \mathcal{E} O set, and $[\sigma_n]_2$ is an r -SVN \mathcal{E} C set, then, there exists an r -SVN \mathcal{E} O set $\gamma_n \in \zeta^{\tilde{X}}$ such that $[\sigma_n]_1 \leq \gamma_n \leq \text{CI}_{\tau^{\tilde{\rho}\tilde{q}\tilde{\eta}}}^{\tilde{L}}(\gamma_n, r) \leq [\sigma_n]_2$.

Proof. Similar to the proof of Theorem 15. \square

Theorem 18. Let $f : (\tilde{X}, \tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}) \rightarrow (\tilde{Y}, \tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be a \mathcal{L} -SVNI-irresolute, bijective, \mathcal{L} -SVNI-irresolute open mapping and $(\tilde{X}, \tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is r -SVNIR $_2$. Then, $(\tilde{Y}, \tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is r -SVNIR $_2$.

Proof. Let $y_{s,t,p} \bar{q}\zeta_n = \text{CI}^*(\zeta_n, r)$. Then, by Definition 11, ζ_n is an r -SVN \mathcal{E} C set in \tilde{Y} . By Theorem 3(2), $f^{-1}(\zeta_n)$ is an r -SVN \mathcal{E} C set in \tilde{X} . Put $y_{s,t,p} = f(x_{s,t,p})$. Then, $x_{s,t,p} \bar{q}f^{-1}(\zeta_n)$. By r -SVNIR $_2$ of \tilde{X} , there exist r -SVN \mathcal{E} O sets $\sigma_n, \gamma_n \in \zeta^{\tilde{X}}$ such that $x_{s,t,p} \in \sigma_n$, $f^{-1}(\zeta_n) \leq \gamma_n$ and $\sigma_n \bar{q}\gamma_n$. Since f is bijective and \mathcal{L} -SVNI-irresolute open, $y_{s,t,p} \in f(\sigma_n)$, $\zeta_n \leq f(f^{-1}(\zeta_n)) \leq f(\gamma_n)$ and $f(\sigma_n) \bar{q}f(\gamma_n)$. Thus, $(\tilde{Y}, \tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is r -SVNIR $_2$. \square

Theorem 19. Let $f : (\tilde{X}, \tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}) \rightarrow (\tilde{Y}, \tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an \mathcal{L} -SVNI-irresolute, bijective, \mathcal{L} -SVNI-irresolute open mapping and $(\tilde{X}, \tilde{\tau}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_1^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ be an r -SVNIR $_3$. Then, $(\tilde{Y}, \tilde{\tau}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}}, \mathcal{I}_2^{\tilde{\rho}\tilde{q}\tilde{\eta}})$ is r -SVNIR $_3$.

Proof. Similar to the proof of Theorem 18. \square

6. Conclusions

In summary, we have introduced the definition of the r -single valued neutrosophic \mathcal{L} -closed and r -single valued neutrosophic \mathcal{L} -open sets over single valued neutrosophic ideal topology space in Šostak's sense. Many consequences have been arisen up to show that how far topological structures are preserved by these r -single valued neutrosophic \mathcal{L} -closed. We also have provided some counterexamples where such properties fail to be preserved. The most important contribution to this area of research is that we have introduced the notion of \mathcal{L} -single valued neutrosophic irresolute mapping, \mathcal{L} -single valued neutrosophic extremally disconnected spaces, \mathcal{L} -single valued neutrosophic normal spaces and that we defined some kinds of separation axioms, namely r -SVNIR $_i$, where $i = \{0, 1, 2, 3\}$, and r -SVNIT $_j$, where $j = \{1, 2, 2\frac{1}{2}, 3, 4\}$, in the sense of Šostak. Some of their characterizations, fundamental properties, and the relations between these notions have been studied.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Acknowledgments: The author would like to express his sincere thanks to Majmaah University for supporting this work. The author is also grateful to the reviewers for their valuable comments and suggestions which led to the improvement of this research.

Conflicts of Interest: The author declares that there is no conflict of interest regarding the publication of this manuscript.

Discussion for Further Works: The theory in this article can be extended in the following natural ways. One may study the properties of neutrosophic metric topological spaces using the concepts defined through this paper.

References

1. Smarandache, F. *Unifying A Field in Logics, Neutrosophy: Neutrosophic Probability; Set and Logic*; American Research Press: Rehoboth, NM, USA, 1999.
2. Hur, K.; Lim, P.K.; Lee, J.G.; Kim, J. The category of neutrosophic sets. *Neutrosophic Sets Syst.* **2016**, *14*, 12–20.
3. Hur, K.; Lim, P.K.; Lee, J.G.; Kim, J. The category of neutrosophic crisp sets. *Ann. Fuzzy Math. Inform.* **2017**, *14*, 43–54.
4. Salama, A.A.; Alblowi, S.A. Neutrosophic set and neutrosophic topological spaces. *IOSR J. Math.* **2012**, *3*, 31–35.
5. Salama, A.A.; Smarandache, F. *Neutrosophic Crisp Set Theory*; Educational Publisher: Columbus, OH, USA, 2015.
6. Salama, A.A.; Smarandache, F.; Kroumov, V. Neutrosophic crisp sets and neutrosophic crisp topological spaces. *Neutrosophic Sets Syst.* **2014**, *2*, 25–30.
7. Chang, C.L. Fuzzy topological spaces. *J. Math. Anal. Appl.* **1968**, *24*, 182–190.
8. Šostak, A. On a fuzzy topological structure. In *Circolo Matematico di Palermo, Palermo; Rendiconti del Circolo Matematico di Palermo, Proceedings of the 13th Winter School on Abstract Analysis, Section of Topology, Srni, Czech Republic, 5–12 January 1985*; Circolo Matematico di Palermo: Palermo, Italy, 1985; pp. 89–103.
9. Liang, C.; Yan, C. Base and subbase in intuitionistic *I*-fuzzy topological spaces. *Hacet. J. Math. Stat.* **2014**, *43*, 231–247.
10. Yan, C.H.; Wang, X.K. Intuitionistic *I*-fuzzy topological spaces. *Czechoslov. Math. J.* **2010**, *60*, 233–252.
11. Wang, H.; Smarandache, F.; Zhang, Y.Q.; Sunderraman, R. Single valued neutrosophic sets. *Multispace Multistruct.* **2010**, *4*, 410–413.
12. El-Gayyar, M. Smooth neutrosophic topological spaces. *Neutrosophic Sets Syst.* **2016**, *65*, 65–72.
13. Kim, J.; Smarandache, F.; Lee, J.G.; Hur, K. Ordinary Single Valued Neutrosophic Topological Spaces. *Symmetry* **2019**, *11*, 1075.
14. Saber, Y.; Alsharari, F.; Smarandache, F. On single-valued neutrosophic ideals in Šostak's sense. *Symmetry* **2020**, *12*, 193.
15. Saber, Y.; Alsharari, F.; Smarandache, F. Connectedness and Stratification of Single-Valued Neutrosophic Topological Spaces. *Symmetry* **2020**, *12*, 1464.
16. Abdel-Basset, M.; Ali, M.; Atef, A. Uncertainty assessments of linear time-cost tradeoffs using neutrosophic set. *Comput. Ind. Eng.* **2020**, *141*, 106–286.
17. Abdel-Basset, M.; Ali, M.; Atef, A. Resource levelling problem in construction projects under neutrosophic environment. *J. Supercomput.* **2020**, *76*, 964–988.
18. Abdel-Basset, M.; Gamal, A.; Son, L. H.; Smarandache, F. A Bipolar Neutrosophic Multi Criteria Decision Making Framework for Professional Selection. *Appl. Sci.* **2020**, *10*, 12–20.
19. Ubeda, T.; Egenhofer, M. J. Topological error correcting in GIS. *Adv. Spat. Databases* **2005**, 281–297.
20. Shang, Y. Average consensus in multi-agent systems with uncertain topologies and multiple time-varying delays. *Linear Algebra Its Appl.* **2014**, *459*, 411–429.
21. Smarandache, F. *A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability and Statistics*, 6th ed.; InfoLearnQuest: Ann Arbor, MI, USA, 2007.
22. Ye, J. A multicriteria decision-making method using aggregation operators for simplified neutrosophic sets. *J. Intell. Fuzzy Syst.* **2014**, *26*, 2450–2466.
23. Yang, H.L.; Guo, Z.L.; Liao, X. On single valued neutrosophic relations. *J. Intell. Fuzzy Syst.* **2016**, *30*, 1045–1056.
24. Alsharari, F. Decomposition of single-valued neutrosophic ideal continuity via fuzzy idealization. *Neutrosophic Sets Syst.* **2020**, *38*, 145–163.