

Short Note

4-Bromo-3-methoxy-1-phenyl-1*H*-pyrazole

Neringa Kleizienė¹, Eglė Arbačiauskienė^{1,*}, Wolfgang Holzer² and Algirdas Šačkus¹

¹ Institute of Synthetic Chemistry, Kaunas University of Technology, Radvilenu pl. 19, LT-50254 Kaunas, Lithuania

² Department of Drug and Natural Product Synthesis, Faculty of Life Sciences, University of Vienna, Althanstrasse 14, A-1090 Vienna, Austria

* Author to whom correspondence should be addressed; E-Mail: eglnedz@ktu.lt.


Received: 29 October 2009 / Accepted: 4 November 2009 / Published: 5 November 2009

Abstract: The title compound was prepared by treatment of 4-bromo-1-phenyl-1*H*-pyrazol-3-ol with sodium hydride/methyl iodide in good yield. Detailed spectroscopic data (¹H NMR, ¹³C NMR, ¹⁵N NMR, IR, MS) are presented.

Keywords: pyrazole; 1*H*-pyrazol-3-ol; methylation; NMR

Bromo(hetero)arenes are valuable starting materials for further functionalization, for instance *via* metalation reactions (halogen-metal exchange) [1] or transition-metal-catalyzed cross coupling reactions [2–4]. With an OH- or OR function in *ortho*-position to the bromo atom, such (hetero)arenes can be considered as potential precursors for condensed systems involving a ring-oxygen atom (condensed furans, pyranes) [5]. However, for some of the above mentioned functionalization reactions, free hydroxy groups are unfavourable, for instance in metalation reactions [6] or in Suzuki-type cross couplings [7]. In such cases, the OH group has to be masked. In this regard, a possible option is to transform the OH into the very stable OMe group. The O–Me bond then can be cleaved in the course of the cyclization reaction into the *O*-containing ring system employing, for instance, pyridine hydrochloride as the reagent [8].

We here present the synthesis of 4-bromo-3-methoxy-1-phenyl-1*H*-pyrazole (**2**), with a protected OH-group in position 3 and a bromo substituent in position 4 of the pyrazole nucleus, which can be considered as a desirable starting compound for further transformation. The preparation of **2** was carried out according to a standard method for *O*-alkylation, namely by reaction of 4-bromo-1-phenyl-1*H*-pyrazol-3-ol (**1**) [9] with methyl iodide in alkaline medium, which afforded the title compound in 88% yield after column chromatography (Scheme 1).



Although compound **2** is reported in a patent application, neither an experimental procedure for its preparation nor spectroscopic/physical data, was provided [10]. Thus, a detailed characterization of **2** including IR, MS and NMR (^1H , ^{13}C , ^{15}N) spectral data as well as microanalytical data is given in the Experimental.

Experimental

The melting point was determined on a Mel-Temp (Capillary Melting point Apparatus) and is uncorrected. Mass spectrum was recorded on a Waters ZQ 2000 instrument (APCI+, 20 V) and IR spectrum on a Perkin Elmer Spectrum GX FT-IR System instrument (KBr-disc). The elemental analysis was performed with an Exeter Analytical CE-440 Elemental Analyzer. All NMR spectra were recorded from CDCl_3 solutions on a Bruker Avance 500 instrument with a 'directly' detecting broadband observe probe (BBFO) at 298 K (500.13 MHz for ^1H , 125.76 MHz for ^{13}C , 50.68 MHz for ^{15}N). The centre of the solvent signal was used as an internal standard which was related to TMS with $\delta = 7.26$ ppm (^1H in CDCl_3) and $\delta = 77.0$ ppm (^{13}C in CDCl_3). The digital resolutions were 0.2 Hz/data point in the ^1H and 0.4 Hz/data point in the ^1H -coupled ^{13}C -NMR spectra (gated decoupling). The ^{15}N NMR spectrum (gradient-selected ^{15}N , ^1H -HMBC) was referenced against external nitromethane.

4-Bromo-3-methoxy-1-phenyl-1H-pyrazole (**2**)

A solution of 4-bromo-1-phenyl-1H-pyrazol-3-ol (**1**) (3.825 g, 16.0 mmol) in dry DMF (20 mL) was cooled to 0 °C under an inert atmosphere and NaH (60% dispersion in mineral oil, 640 mg, 16.0 mmol) was added portionwise. After stirring for 15 min, methyl iodide (1.2 mL, 19.2 mmol) was added dropwise at 0 °C. The mixture was warmed to r.t. and stirred at 60 °C for 1 h (TLC control, eluent: ethyl acetate–*n*-hexane, 1:10). Then, 10 mL of water were added and the mixture was extracted with ethyl acetate. The organic layers were combined, washed with brine and dried over anhydrous Na_2SO_4 and filtered. The solvent was evaporated under reduced pressure and the residue was purified by column chromatography (silica gel, eluent: ethyl acetate–*n*-hexane, 1:20) to give pure **2** as colorless crystals, m.p. 24.5–25.5 °C. Yield: 3.56 g (88%).

IR (KBr) ν (cm^{-1}): 2947, 1554, 1502, 1099, 749, 686.

MS (EI, 70 eV): (m/z , %) 253/255 (M^+ , 100/99), 175 ($[\text{M}-\text{Br}+\text{H}]^+$, 47), 174 ($[\text{M}-\text{Br}]^+$, 75).

^1H NMR (CDCl_3): δ (ppm) 4.05 (s, 3H, Me), 7.23 (m, 1H, Ph H-4), 7.42 (m, 2H, Ph H-3,5), 7.56 (m, 2H, Ph H-2,6), 7.78 (s, 1H, H-5).

^{13}C NMR (CDCl_3): δ (ppm) 56.8 (OCH_3 , $^1J = 146.2$ Hz), 82.0 (C-4, $^2J(\text{C4,H5}) = 5.2$ Hz), 117.6 (Ph C-2,6), 125.8 (Ph C-4), 127.8 (C-5, $^1J = 191.7$ Hz), 129.4 (Ph C-3,5), 139.7 (Ph C-1), 161.2 (C-3, $^3J(\text{C3,H5}) = 8.8$ Hz, $^3J(\text{C3,OCH}_3) = 3.7$ Hz).

^{15}N NMR (CDCl_3): δ (ppm) -187.8 (N-1), -119.9 (N-2).

Anal. Calcd for $\text{C}_5\text{H}_9\text{BrN}_2\text{O}$: C, 47.46%; H, 3.58%; N, 11.07%. Found: C, 47.08%; H, 3.42%; N 10.72%.

References and Notes

1. Inoue, A.; Kitagawa, K.; Shinokubo, H.; Oshima, K. Selective halogen-magnesium exchange reaction via organomagnesium ate complex. *J. Org. Chem.* **2001**, *66*, 4333–4339 and references cited herein.
2. Sonogashira, K.; Tohda, Y.; Hagihara, N. Convenient synthesis of acetylenes. Catalytic substitutions of acetylenic hydrogen with bromo alkenes, iodo arenes, and bromopyridines. *Tetrahedron Lett.* **1975**, *16*, 4467–4470.
3. Miyaura, N.; Suzuki, A. Stereoselective synthesis of arylated (*E*)-alkenes by the reaction of alk-1-enylboranes with aryl halides in the presence of palladium catalyst. *J. Chem. Soc. Chem. Commun.* **1979**, *19*, 866–867.
4. Miyaura, N.; Suzuki, A. Palladium-catalyzed cross-coupling reactions of organoboron compounds. *Chem. Rev.* **1995**, *95*, 2457–2583.
5. Ellis, G.P. *Synthesis of Fused Heterocycles*; John Wiley & Sons: New York, NY, USA, 1987; pp. 477–484.
6. Riego, E.; Bayó, N.; Cuevas, C.; Albericio, F.; Álvarez, M. A new approach to 3-hydroxyquinoline-2-carboxylic acid. *Tetrahedron* **2005**, *61*, 1407–1411.
7. Arbačiauskienė, E.; Vilkauskaitė, G.; Eller, G.A.; Holzer, W.; Šačkus, A. Pd-catalyzed cross-coupling reactions of halogenated 1-phenylpyrazol-3-ols and related triflates. *Tetrahedron* **2009**, *65*, 7817–7824 and the references cited herein.
8. Ellis, G.P. *Synthesis of Fused Heterocycles*; John Wiley & Sons: New York, NY, USA, 1987; pp. 128, 190, 352.
9. O'Brien, D.F.; Gates, J.W., Jr. Some reactions of 3-hydroxy-1-phenylpyrazole. *J. Org. Chem.* **1966**, *31*, 1538–1542.
10. Koenig, H.; Goetz, N.; Klein, U.; Eller, K. Process for producing *N*-substituted 3-hydroxypyrazoles *WO 9703969*, 1997; *Chem. Abstr.* **1997**, *126*, 199566.