

Article

A Reliable Algorithm for a Local Fractional Tricomi Equation Arising in Fractal Transonic Flow

Jagdev Singh ^{1,*}, Devendra Kumar ² and Juan J. Nieto ^{3,4}

¹ Department of Mathematics, Jagan Nath University, Jaipur 303901, India

² Department of Mathematics, JECRC University, Jaipur 303905, India; devendra.maths@gmail.com

³ Departamento de Análise Matemática, Faculdade de Matemáticas, Universidade de Santiago de Compostela, Santiago de Compostela 15782, Spain; juanjose.nieto.roig@usc.es

⁴ Faculty of Science, King Abdulaziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia

* Correspondence: jagdevsinghrathore@gmail.com; Tel.: +91-946-090-5224

Academic Editor: Carlo Cattani

Received: 13 April 2016; Accepted: 23 May 2016; Published: 25 May 2016

Abstract: The pivotal proposal of this work is to present a reliable algorithm based on the local fractional homotopy perturbation Sumudu transform technique for solving a local fractional Tricomi equation occurring in fractal transonic flow. The proposed technique provides the results without any transformation of the equation into discrete counterparts or imposing restrictive assumptions and is completely free of round-off errors. The results of the scheme show that the approach is straightforward to apply and computationally very user-friendly and accurate.

Keywords: Tricomi equation; fractal transonic flow; local fractional derivative; homotopy perturbation method; local fractional Sumudu transform method

1. Introduction

Partial differential equations of mixed type with boundary conditions have played an important role in describing real world problems such as in elementary research conducted by Tricomi [1]. Mixed type partial differential equations [2,3] are used to investigate transonic flow and they produce special boundary value problems, known as Tricomi and Frankl problems [1,4]. Transonic flows include a change from the subsonic to the hypersonic region [4,5] via the sonic curve. Consequently, transonic flows are very attractive phenomena occurring in aeronautics and hydraulics. The familiar mixed type partial differential equation is known as a Tricomi equation, $yu_{xx} + u_{yy} = 0$, because of Tricomi, who found this mathematical model, for the function $u = u(x, y)$ of two variables x and y . It acts as a basis for the mathematical modelling of the transonic flows, since it is of elliptic and hyperbolic type, where the coefficient y of the second order partial differential coefficient of the required function $u = u(x, y)$ with respect to x changes sign. This mathematical equation is also parabolic at the points where y vanishes.

The Tricomi equation [1] is a mixed type of linear partial differential equation of the second order, which has been used to narrate the theory of plane transonic flow [6–9]. The Tricomi equation was used to recount differentiable problems for the theory of plane transonic flow. However, for the fractal theory of plane transonic flow with non-differentiable expressions, the Tricomi equation is not registered to report them. Recently, local fractional calculus [10] was tried for non-differentiable problems, for instance fractal heat conduction [10,11], damped and dissipative wave equations in fractal strings [12], local fractional Laplace equations [13], the Helmholtz equation associated with local fractional derivatives [14], the wave equation on Cantor sets [15], Navier–Stokes equations on Cantor sets [16], local fractional Schrödinger equations [17], Korteweg–de Vries equations involving

local fractional derivative [18], etc. In recent times, the local fractional model of the Tricomi equation in fractal transonic flows was recommended in the form [19,20]

$$\frac{y^\beta}{\Gamma(1+\beta)} \frac{\partial^{2\beta} u(x,y)}{\partial x^{2\beta}} + \frac{\partial^{2\beta} u(x,y)}{\partial y^{2\beta}} = 0, \quad (1)$$

where the size $u(x,y)$ is the non-differentiable function, and the local fractional operator indicates [10]

$$\frac{\partial^\beta u(x,t)}{\partial t^\beta} = \frac{\Delta^\beta (u(x,t) - u(x,t_0))}{(t - t_0)}, \quad (2)$$

where

$$\Delta^\beta (u(x,t) - u(x,t_0)) \cong \Gamma(1+\beta) [u(x,t) - u(x,t_0)] \quad (3)$$

The Tricomi equation finds its application in modeling transonic flow [21–23]. Inspired and motivated by the ongoing research in this area and wide applications of local fractional differential equations, we propose the local fractional homotopy perturbation Sumudu transform method (LFHPSTM) to solve the local fractional model of the Tricomi equation appearing in fractal transonic flow pertaining to the local fractional derivative boundary value conditions. The LFHPSTM is a conjunction of the classical homotopy perturbation method (HPM) [24–26] and the local fractional Sumudu transform technique. The formation of this article is as developed. In Section 2, the local fractional integrals and derivatives are initiated. In Section 3, the local fractional homotopy perturbation Sumudu transform method is proposed. In Section 4, the non-differentiable numerical solutions for local fractional Tricomi equation along the local fractional derivative boundary value conditions are specified. Finally, in Section 5, the conclusions are discussed.

2. Local Fractional Integrals and Derivatives

In this section, we review the basic theory of local fractional calculus, which is applied in this research article.

Definition 1 ([10–20]). For the relation $|x - x_0| < \delta$, when $\varepsilon, \delta > 0$ and $\varepsilon > R$, we permit the function $f(x) \in C_\beta(a, b)$, while

$$|f(x) - f(x_0)| < \varepsilon^\beta, 0 < \beta \leq 1, \quad (4)$$

exists.

Definition 2 ([10–20]). Consider the interval $[a, b]$ and (t_j, t_{j+1}) , $j = 0, \dots, N-1$, $t_0 = a$, and $t_N = b$ with $\Delta t_j = t_{j+1} - t_j$, $\Delta t = \max\{\Delta t_0, \Delta t_1, \Delta t_2, \dots\}$ a partition of this interval. Then, the local fractional integral of $f(x)$ is explained as

$${}_a I_b^{(\beta)} f(x) = \frac{1}{\Gamma(1+\beta)} \int_a^b f(t) (dt)^\beta = \frac{1}{\Gamma(1+\beta)} \lim_{\Delta t \rightarrow 0} \sum_{j=0}^{j=N-1} f(t_j) (\Delta t_j)^\beta, \quad (5)$$

Definition 3 ([10–20]). Suppose the function $f(x)$ fulfill state in Equation (4), then the inverse formula of Equation (5) is defined as follows:

$$\frac{d^\beta f(x_0)}{dx^\beta} = D_x^{(\beta)} f(x_0) = \frac{\Delta^\beta (f(x) - f(x_0))}{(x - x_0)^\beta}, \quad (6)$$

where

$$\Delta^\beta (f(x) - f(x_0)) \cong \Gamma(1+\beta) [f(x) - f(x_0)]. \quad (7)$$

The formula of the local fractional derivative, employed in this paper, is given as follows [10]:

$$\frac{d^\beta}{dx^\beta} \frac{x^{n\beta}}{\Gamma(1+n\beta)} = \frac{x^{(n-1)\beta}}{\Gamma(1+(n-1)\beta)}, n \in N. \tag{8}$$

3. Local Fractional Sumudu Transform

The Sumudu transform was initially proposed and developed by Watugala [27] and some of its important properties were discovered and investigated by Belgacem *et al.* [28] and Belgacem and Karaballi [29]. Katatbeh and Belgacem [30] employed the Sumudu transform to solve fractional differential equations. Gupta *et al.* [31] used the Sumudu transform to solve generalized fractional kinetic equations. Belgacem [32] investigated the applications of the Sumudu transform to Bessel functions and equations. Belgacem [33] introduced and analyzed deeper Sumudu properties. Bulut *et al.* [34] obtained the analytical solutions of some fractional ordinary differential equations by using the Sumudu transform technique. The Sumudu transform method is also coupled with HPM to investigate the fractional biological population model [35]. The local fractional Sumudu transform of a function $f(x)$ is first introduced and defined by Srivastava *et al.* [36] in the following manner:

$$\begin{aligned} LFS_\beta \{f(x)\} &= F_\beta(z) \\ &= \frac{1}{\Gamma(1+\beta)} \int_0^\infty E_\beta(-z^{-\beta} x^\beta) \frac{f(x)}{z^\beta} (dx)^\beta, 0 < \beta \leq 1 \end{aligned} \tag{9}$$

and the inverse formula is expressed as follows

$$LFS_\beta^{-1} \{F_\beta(z)\} = f(x), 0 < \beta \leq 1. \tag{10}$$

4. Local Fractional Homotopy Perturbation Sumudu Transform Method

In order to establish the basic idea of the LFHPSTM, we assume the following linear differential equation with a local fractional derivative

$$L_\beta u(x, t) + R_\beta u(x, t) = h(x, t), \tag{11}$$

where L_β denotes the linear local fractional differential operator, R_β is the remaining linear operator and $h(x, t)$ is a source term.

Using the local Sumudu transform on Equation (11) yields

$$\begin{aligned} U_\beta(x, z) &= u(x, 0) + z^\beta u^\beta(x, 0) + z^{2\beta} u^{2\beta}(x, 0) + \dots \\ &+ z^{(k-1)\beta} u^{(k-1)\beta}(x, 0) - z^{k\beta} LFS_\beta [R_\beta u(x, t)] + z^{k\beta} LFS_\beta [h(x, t)]. \end{aligned} \tag{12}$$

Applying the inverse of the local fractional Sumudu transform on Equation (12), we have the following result

$$\begin{aligned} u(x, t) &= u(x, 0) + \frac{t^\beta}{\Gamma(1+\beta)} u^\beta(x, 0) + \frac{t^{2\beta}}{\Gamma(1+2\beta)} u^{2\beta}(x, 0) + \dots \\ &+ \frac{t^{(k-1)\beta}}{\Gamma(1+(k-1)\beta)} u^{(k-1)\beta}(x, 0) - LFS_\beta^{-1} \left[z^{k\beta} LFS_\beta [R_\beta u(x, t)] \right] \\ &+ LFS_\beta^{-1} \left[z^{k\beta} LFS_\beta [h(x, t)] \right]. \end{aligned} \tag{13}$$

Now we use the HPM [24–26]

$$u(x, t) = \sum_{n=0}^\infty p^n u_n(x, t). \tag{14}$$

Substituting Equation (14) in Equation (13), we get the following result:

$$\begin{aligned} \sum_{n=0}^{\infty} p^n u_n(x, t) = & u(x, 0) + \frac{t^\beta}{\Gamma(1+\beta)} u^\beta(x, 0) + \frac{t^{2\beta}}{\Gamma(1+2\beta)} u^{2\beta}(x, 0) + \dots \\ & + \frac{t^{(k-1)\beta}}{\Gamma(1+(k-1)\beta)} u^{(k-1)\beta}(x, 0) - p LFS_\beta^{-1} \left[z^{k\beta} LFS_\beta \left[R_\beta \sum_{n=0}^{\infty} p^n u_n(x, t) \right] \right] \\ & + LFS_\beta^{-1} \left[z^{k\beta} LFS_\beta [h(x, t)] \right]. \end{aligned} \tag{15}$$

which is a mixture of the local fractional Sumudu transform technique and HPM. Comparing the coefficients of like powers of p , we get

$$\begin{aligned} p^0 : u_0(x, t) &= u(x, 0) + \frac{t^\beta}{\Gamma(1+\beta)} u^\beta(x, 0) + \frac{t^{2\beta}}{\Gamma(1+2\beta)} u^{2\beta}(x, 0) + \dots \\ &\quad + \frac{t^{(k-1)\beta}}{\Gamma(1+(k-1)\beta)} u^{(k-1)\beta}(x, 0) + LFS_\beta^{-1} \left[z^{k\beta} LFS_\beta [h(x, t)] \right], \\ p^1 : u_1(x, t) &= -LFS_\beta^{-1} \left[z^{k\beta} LFS_\beta [R_\beta u_0(x, t)] \right], \\ p^2 : u_2(x, t) &= -LFS_\beta^{-1} \left[z^{k\beta} LFS_\beta [R_\beta u_1(x, t)] \right], \\ &\vdots \end{aligned} \tag{16}$$

Therefore, the solution of Equation (11) is given by

$$u(x, t) = \lim_{N \rightarrow \infty} \sum_{n=0}^N u_n(x, t) \tag{17}$$

5. Nondifferential Solutions for the Local Fractional Tricomi Equation

In this section, we present the nondifferential solutions for the Tricomi equation pertaining to the local fractional derivative occurring in fractal transonic flow with local fractional derivative boundary value conditions.

Example 1. Firstly, we investigate the following local fractional Tricomi equation

$$\frac{y^\beta}{\Gamma(1+\beta)} \frac{\partial^{2\beta} u(x, y)}{\partial x^{2\beta}} + \frac{\partial^{2\beta} u(x, y)}{\partial y^{2\beta}} = 0 \tag{18}$$

subject to the initial-boundary value conditions

$$\begin{aligned} u(x, 0) &= 0, \\ u(h, l) &= 0, \\ u(x, 0) &= \frac{x^{2\beta}}{\Gamma(1+2\beta)}, \\ \frac{\partial^\beta u(x, 0)}{\partial y^\beta} &= 0. \end{aligned} \tag{19}$$

Applying the local fractional Sumudu transform on Equation (18), we get

$$U_\beta(x, z) = u(x, 0) + z^\beta u^\beta(x, 0) + z^{2\beta} LFS_\beta \left[-\frac{y^\beta}{\Gamma(1+\beta)} \frac{\partial^{2\beta} u(x, y)}{\partial x^{2\beta}} \right] \tag{20}$$

which implies

$$U_\beta(x, z) = \frac{x^{2\beta}}{\Gamma(1+2\beta)} + z^{2\beta} LFS_\beta \left[-\frac{y^\beta}{\Gamma(1+\beta)} \frac{\partial^{2\beta} u(x, y)}{\partial x^{2\beta}} \right] \tag{21}$$

Applying the inverse local fractional Sumudu transform to Equation (21) gives

$$u(x, y) = \frac{x^{2\beta}}{\Gamma(1 + 2\beta)} + LFS_{\beta}^{-1} \left[z^{2\beta} LFS_{\beta} \left[-\frac{y^{\beta}}{\Gamma(1 + \beta)} \frac{\partial^{2\beta} u(x, y)}{\partial x^{2\beta}} \right] \right] \tag{22}$$

Now using HPM [24–26] we get

$$\sum_{n=0}^{\infty} p^n u_n(x, y) = \frac{x^{2\beta}}{\Gamma(1 + 2\beta)} + p LFS_{\beta}^{-1} \left[z^{2\beta} LFS_{\beta} \left[-\frac{y^{\beta}}{\Gamma(1 + \beta)} \frac{\partial^{2\beta} \left(\sum_{n=0}^{\infty} p^n u_n(x, y) \right)}{\partial x^{2\beta}} \right] \right] \tag{23}$$

Comparing the like powers of p , we get the following components of the series solution

$$\begin{aligned} p^0 : u_0(x, y) &= \frac{x^{2\beta}}{\Gamma(1+2\beta)}, \\ p^1 : u_1(x, y) &= -\frac{y^{3\beta}}{\Gamma(1+3\beta)}, \\ p^2 : u_2(x, y) &= 0, \\ &\vdots \end{aligned} \tag{24}$$

Finally, we get the exact solution of Equation (18) with the local fractional derivative boundary value conditions Equation (19), namely

$$\begin{aligned} u(x, y) &= \lim_{N \rightarrow \infty} \sum_{n=0}^N u_n(x, y) \\ &= \frac{x^{2\beta}}{\Gamma(1+2\beta)} - \frac{y^{3\beta}}{\Gamma(1+3\beta)} \end{aligned} \tag{25}$$

Example 2. Next, we study the local fractional Tricomi equation of the form

$$\frac{y^{\beta}}{\Gamma(1 + \beta)} \frac{\partial^{2\beta} u(x, y)}{\partial x^{2\beta}} + \frac{\partial^{2\beta} u(x, y)}{\partial y^{2\beta}} = 0 \tag{26}$$

and the initial conditions are presented as

$$\begin{aligned} u(x, 0) &= \frac{x^{\beta}}{\Gamma(1+\beta)}, \\ \frac{\partial^{\beta} u(x, 0)}{\partial y^{\beta}} &= \frac{x^{\beta}}{\Gamma(1+\beta)}, \end{aligned} \tag{27}$$

Applying the local fractional Sumudu transform to Equation (26), we get

$$U_{\beta}(x, z) = u(x, 0) + z^{\beta} u^{\beta}(x, 0) + z^{2\beta} LFS_{\beta} \left[-\frac{y^{\beta}}{\Gamma(1 + \beta)} \frac{\partial^{2\beta} u(x, y)}{\partial x^{2\beta}} \right] \tag{28}$$

which gives

$$U_{\beta}(x, z) = \frac{x^{\beta}}{\Gamma(1 + \beta)} + z^{\beta} \frac{x^{\beta}}{\Gamma(1 + \beta)} + z^{2\beta} LFS_{\beta} \left[-\frac{y^{\beta}}{\Gamma(1 + \beta)} \frac{\partial^{2\beta} u(x, y)}{\partial x^{2\beta}} \right] \tag{29}$$

Applying the inverse local fractional Sumudu transform to Equation (29), we have

$$u(x, y) = \frac{x^{\beta}}{\Gamma(1 + \beta)} \left[1 + \frac{y^{\beta}}{\Gamma(1 + \beta)} \right] + LFS_{\beta}^{-1} \left[z^{2\beta} LFS_{\beta} \left[-\frac{y^{\beta}}{\Gamma(1 + \beta)} \frac{\partial^{2\beta} u(x, y)}{\partial x^{2\beta}} \right] \right] \tag{30}$$

Now using HPM [24–26], we get

$$\sum_{n=0}^{\infty} p^n u_n(x, y) = \frac{x^\beta}{\Gamma(1+\beta)} \left[1 + \frac{y^\beta}{\Gamma(1+\beta)} \right] + p LFS_\beta^{-1} \left[z^{2\beta} LFS_\beta \left[-\frac{y^\beta}{\Gamma(1+\beta)} \frac{\partial^{2\beta} \left(\sum_{n=0}^{\infty} p^n u_n(x, y) \right)}{\partial x^{2\beta}} \right] \right] \quad (31)$$

Comparing the coefficients of like powers of p provides

$$\begin{aligned} p^0 : u_0(x, y) &= \frac{x^\beta}{\Gamma(1+\beta)} \left[1 + \frac{y^\beta}{\Gamma(1+\beta)} \right], \\ p^1 : u_1(x, y) &= 0, \\ p^2 : u_2(x, y) &= 0, \\ &\vdots \end{aligned} \quad (32)$$

Hence, we get the exact solution with non-differential term, as follows:

$$u(x, y) = \lim_{N \rightarrow \infty} \sum_{n=0}^N u_n(x, y) = \frac{x^\beta}{\Gamma(1+\beta)} \left[1 + \frac{y^\beta}{\Gamma(1+\beta)} \right] \quad (33)$$

6. Conclusions

In the present paper, the local fractional Tricomi equation with its applications in fractal transonic flow is discussed by using the local fractional homotopy perturbation Sumudu transform technique. We obtain the solution with non-differential terms by applying this approach. The results show that the proposed technique is very efficient and can be used to solve various kinds of local fractional differential equations. Hence, the introduced method is a powerful tool for solving local fractional linear equations of physical importance.

Acknowledgments: The authors are very grateful to the referees for their invaluable suggestions and comments for the improvement of this paper.

Author Contributions: Jagdev Singh, Devendra Kumar and Juan J. Nieto contributed equally to this work. All authors have read and approved the final manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Tricomi, F.G. Sulle Equazioni Lineari alle derivate Parziali di 20 Ordine, di Tipo Misto, Atti Accad. Nazionale dei Lincei **1923**, *14*, 133–247. (In Italian)
2. Chen, D.; Wen, G. Initial-Oblique Derivative Problem for Nonlinear Parabolic Equations in High Dimensional Domains. *Int. J. Appl. Math. Stat.* **2007**, *8*, 8–19.
3. Rassias, J.M. Mixed Type Partial Differential Equations With Initial and Boundary Values in Fluid Mechanics. *Int. J. Appl. Math. Stat.* **2008**, *13*, 77–107.
4. Frankl, F.I.; On, S.A. On the problems of Chaplygin for mixed subsonic and supersonic flows. *Izvestiya Rossiiskoi Akademii Nauk. Seriya Matematicheskaya* **1945**, *9*, 121–143.
5. Guderley, G. On the presence of shocks in mixed subsonic-supersonic flow patterns. In *Advanced in Applied Mechanics*; Academic Press: New York, NY, USA, 1953; Volume 3, pp. 145–184.
6. Lupo, D.; Payne, K.R. A dual variational approach to a class of nonlocal semilinear Tricomi problems. *Nonlinear Differ. Equ. Appl.* **1999**, *6*, 247–266. [[CrossRef](#)]

7. Yagdjian, K. Global existence for the n -dimensional semilinear Tricomi-type equations. *Commun. Partial Differ. Equ.* **2006**, *31*, 907–944. [[CrossRef](#)]
8. Yagdjian, K. A note on the fundamental solution for the Tricomi-type equation in the hyperbolic domain. *J. Differ. Equ.* **2004**, *206*, 227–252. [[CrossRef](#)]
9. Rassias, J.M. Uniqueness of quasi-regular solutions for abi-parabolic elliptic bi-hyperbolic Tricomi problem. *Complex Var. Theory Appl.* **2002**, *47*, 707–718. [[CrossRef](#)]
10. Yang, X.J. *Advanced Local Fractional Calculus and Its Applications*; World Science: New York, NY, USA, 2012.
11. Yang, A.M.; Zhang, Y.Z.; Long, Y. The Yang-Fourier transforms to heat-conduction in a semi-infinite fractal bar. *Therm. Sci.* **2013**, *17*, 707–713. [[CrossRef](#)]
12. Su, W.H.; Baleanu, D.; Yang, X.J.; Jafari, H. Damped wave equation and dissipative wave equation in fractal strings within the local fractional variational iteration method. *Fixed Point Theory Appl.* **2013**, *2013*, 89. [[CrossRef](#)]
13. Yang, X.J.; Yang, Y.J.; Baleanu, D. A local fractional variational iteration method for Laplace equation within local fractional operator. *Abstr. Appl. Anal.* **2013**, *2013*, 202650. [[CrossRef](#)]
14. Yang, A.M.; Chen, Z.; Srivastava, H.M.; Yang, X.J. Application of the local fractional series expansion method and the variation iteration method to the Helmholtz equation involving local fractional derivative operators. *Abstr. Appl. Anal.* **2013**, *2013*, 259125. [[CrossRef](#)]
15. Baleanu, D.; Machado, J.A.T.; Cattani, C.; Baleanu, M.C.; Yang, X.J. Local fractional variational iteration and decomposition methods for wave equation on Cantor sets within local fractional operators. *Abstr. Appl. Anal.* **2014**, *2014*, 535048. [[CrossRef](#)]
16. Yang, X.J.; Srivastava, H.M.; He, J.H.; Baleanu, D. Cantor-type cylindrical-coordinate method for differential equations with local fractional derivatives. *Phys. Lett. A* **2013**, *377*, 1696–1700. [[CrossRef](#)]
17. Zhao, Y.; Cheng, D.F.; Yang, X.J. Approximation solutions for local fractional Schrödinger equation in the one dimensional Cantorian system. *Adv. Math. Phys.* **2013**, *2013*, 291386. [[CrossRef](#)]
18. Yang, X.J.; Hristov, J.; Srivastava, H.M.; Ahmad, B. Modelling Fractal Waves on Shallow Water Surfaces via Local Fractional Korteweg–de Vries Equation. *Abstr. Appl. Anal.* **2014**, *2014*, 278672. [[CrossRef](#)]
19. Yang, A.M.; Zhang, Y.Z.; Zhang, X.L. The Nondifferential Solution for Local Fractional Tricomi Equation Arising in Fractal Transonic Flow by Local Fractional Variational Iteration Method. *Adv. Math. Phys.* **2014**, *2014*, 983254. [[CrossRef](#)]
20. Niu, X.F.; Zhang, C.L.; Li, Z.B.; Zhao, Y. Local Fractional Derivative Boundary Value Problems for Tricomi Equation Arising in Fractal Transonic Flow. *Abstr. Appl. Anal.* **2014**, *2014*, 872318. [[CrossRef](#)]
21. Chen, G.Q.; Feldman, M. Shock Reflection-Diffraction and von Neumann’s Con-jectures. In *Annals of Mathematics Studies Series*; Princeton University Press: Princeton, NJ, USA, 2015.
22. Morawetz, C.M. Mixed equations and transonic flow. *J. Hyperbolic Differ. Equ.* **2004**, *1*, 1–26. [[CrossRef](#)]
23. Tricomi, F. *On Second-Order Linear Partial Differential Equations of Mixed Type*; Moscow: Leningrad, Russia, 1947. (In Russian)
24. He, J.H. Homotopy perturbation method: A new nonlinear analytical technique. *Appl. Math. Comput.* **2003**, *135*, 73–79. [[CrossRef](#)]
25. He, J.H. Comparison of homotopy perturbation method and homotopy analysis method. *Appl. Math. Comput.* **2004**, *156*, 527–539. [[CrossRef](#)]
26. He, J.H. The homotopy perturbation method for nonlinear oscillators with discontinuities. *Appl. Math. Comput.* **2004**, *151*, 287–292. [[CrossRef](#)]
27. Watugala, G.K. Sumudu transform—A new integral transform to solve differential equations and control engineering problems. *Int. J. Math. Educ. Sci. Technol.* **1993**, *24*, 35–43. [[CrossRef](#)]
28. Belgacem, F.B.M.; Karaballi, A.A.; Kalla, S.L. Analytical investigations of the Sumudu transform and applications to integral production equations. *Math. Probl. Eng.* **2003**, *3*, 103–118. [[CrossRef](#)]
29. Belgacem, F.B.M.; Karaballi, A.A. Sumudu Transform Fundamental Properties Investigations and Applications. *J. Appl. Math. Stoch. Anal.* **2006**, *2006*, 91083. [[CrossRef](#)]
30. Katatbeh, Q.K.; Belgacem, F.B.M. Applications of the Sumudu transform to fractional differential equations. *Nonlinear Stud.* **2011**, *18*, 99–112.
31. Gupta, V.G.; Sharma, B.; Belgacem, F.B.M. On the solutions of generalized fractional kinetic equations. *Appl. Math. Sci.* **2011**, *5*, 899–910. [[CrossRef](#)]

32. Belgacem, F.B.M. Applications with the Sumudu transform to Bessel functions and equations. *Appl. Math. Sci.* **2010**, *4*, 3665–3686.
33. Belgacem, F.B.M. Introducing and analyzing deeper Sumudu properties. *Nonlinear Stud.* **2006**, *13*, 23–42.
34. Bulut, H.; Baskonus, H.M.; Belgacem, F.B.M. The analytical solutions of some fractional ordinary differential equations by Sumudu transform method. *Abstr. Appl. Anal.* **2013**, *2013*, 203875. [[CrossRef](#)]
35. Singh, J.; Kumar, D.; Kilicman, A. Numerical solutions of nonlinear fractional partial differential equations arising in spatial diffusion of biological populations. *Abstr. Appl. Anal.* **2014**, *2014*, 535793. [[CrossRef](#)]
36. Srivastava, H.M.; Golmankhaneh, A.K.; Baleanu, D.; Yang, X.J. Local Fractional Sumudu Transform with Application to IVPs on Cantor Sets. *Abstr. Appl. Anal.* **2014**, *2014*, 620529. [[CrossRef](#)]



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).