

Article

Dust Deficiency in the Interacting Galaxy NGC 3077

Jairo Armijos-Abendaño ^{1,*}, Ericson López ¹, Mario Llerena ¹, Franklin Aldás ¹
and Crispin Logan ²

¹ Observatorio Astronómico de Quito, Escuela Politécnica Nacional, Av. Gran Colombia S/N, Quito 170403, Ecuador; ericson.lopez@epn.edu.ec (E.L.); mario.llerena01@epn.edu.ec (M.L.); franklin.aldas@epn.edu.ec (F.A.)

² H.H. Wills Physics Laboratory, University of Bristol, Tyndall Ave, Bristol BS8 1TL, UK; crispin.logan@bristol.ac.uk

* Correspondence: jairo.armijos@epn.edu.ec

Academic Editor: Emilio Elizalde

Received: 24 July 2017; Accepted: 13 September 2017; Published: date

Abstract: Using 70 μm observations taken with the PACS instrument of the Herschel space telescope, the dust content of the nearby and interacting spiral galaxy NGC 3077 has been compared with the dust content of the isolated galaxies such as NGC 2841, NGC 3184 and NGC 3351. The dust content has allowed us to derive dust-to-gas ratios for the four spiral galaxies of our sample. We find that NGC 2841, NGC 3184 and NGC 3351 have dust masses of $6.5\text{--}9.1 \times 10^7 M_{\odot}$, which are a factor of ~ 10 higher than the value found for NGC 3077. This result shows that NGC 3077 is a dust deficient galaxy, as was expected, because this galaxy is affected by tidal interactions with its neighboring galaxies M81 and M82. NGC 3077 reveals a dust-to-gas ratio of 17.5%, much higher than the average ratio of 1.8% of the isolated galaxies, evidencing that NGC 3077 is also deficient in $\text{H}_2 + \text{HI}$ gas. Therefore, it seems that, in this galaxy, gas has been stripped more efficiently than dust.

Keywords: spiral galaxies; dust mass; dust-to-gas ratio

1. Introduction

Several studies have shown that galaxies located in high density environments lose atomic neutral hydrogen (HI) due to tidal interactions [1], simultaneous ram pressure and tidal interactions [2], among other mechanisms such as viscous stripping [3] and thermal evaporation [4]. Galaxies in high density environments have less HI content than isolated galaxies [5,6]. There are few works devoted to the study of environmental effects of spiral galaxies on dust content; as examples, we have the studies of [7,8]. Not only HI gas but also dust is stripped in spiral galaxies located in a cluster environment [7,8], where HI gas is stripped more efficiently than dust [7]. The authors of [9] have studied the emission of dust from a large sample of nearby galaxies, including the galaxies addressed in this paper. Their work focused on the dust-to-gas ratios derived from maps of dust mass surface density, obtained from pixel-by-pixel modeling of infrared data.

To investigate possible effects of the environment on the dust of a nearby spiral galaxy, we study the dust content and the dust-to-gas ratio of the galaxy NGC 3077, that is part of a galaxy triplet [10] and therefore affected by tidal interactions. For comparison purposes, we include in our sample the spiral galaxies NGC 2841, NGC 3184 and NGC 3351, which we have considered as isolated galaxies. The positions, morphology and distances of the galaxies of our sample are given in Table 1. In a recent work, dedicated to the study of the environmental effects on dust of nearby galaxies [7], our sources have not been considered.

Table 1. Galaxy sample, morphology and positions.

| Galaxy Name | RA ¹ (hh:mm:ss.s) | DEC ¹ (dd:mm:ss.s) | Morphology ¹ | Distance ¹ (Mpc) |
|-------------|---------------------------------|----------------------------------|-------------------------|--------------------------------|
| NGC 2841 | 09:22:02.7 | +50:58:35.3 | SAa C | 14.6 |
| NGC 3077 | 10:03:19.1 | +68:44:02.2 | S0 C | 3.8 |
| NGC 3184 | 10:18:17.0 | +41:25:27.8 | SAc C | 11.3 |
| NGC 3351 | 10:43:57.7 | +11:42:13.0 | SBb C | 10.5 |

¹ Information taken from the SIMBAD Astronomical Database.

2. Infrared Data

As it was mentioned above, we aim to study the dust content of four nearby spiral galaxies. For this, we use 70 μm archival maps that can be downloaded from the SIMBAD Astronomical Database¹. These data were observed with the Photoconductor Array Camera and Spectrometer (PACS) instrument of the Herschel space telescope² and obtained thanks to the KINGFISH (Key Insights on Nearby Galaxies: a Far-Infrared Survey with Herschel) survey. These data were first published by [11]. The PACS maps of our four galaxies are shown in Figure 1.




Figure 1. Images at 70 μm from our galaxy sample. The white ellipse is used to derive the infrared flux density (see Section 3.1). The white line indicates the major-axis of each galaxy.

¹ <http://simbad.u-strasbg.fr/simbad/>

² Herschel is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia with an important NASA participation

3. Results and Discussion

3.1. Infrared Flux Density

To estimate the infrared flux density (S_λ) for the galaxies in our sample, we define an ellipse which encompasses almost all the infrared emission from each galaxy disk (see Figure 1). The S_λ will be used to estimate the dust mass in Section 3.2. The sizes of the ellipse used to enclose the galaxy disks are the same as those used to derive CO(2-1) luminosities by [12], who used CO luminosities to derive H₂ + HI masses. The similarity in the size of the ellipses in both studies is due to the fact that a good correlation exists in the spatial distribution of the CO(2-1) line emission and the 70 μm emission. This correlation can be tested by comparing the NGC 2841, NGC 3077, NGC 3184 and NGC 3351 maps given in Figure 1 of [12] with the maps of galaxies given in Figure 1 of this work. The derived values of S_λ are given in Table 2, where we also listed the semi-major axis, semi-minor axis and the position angle (PA) of the ellipses.

3.2. Dust Mass

Once the S_λ flux density is derived, we can estimate the dust mass (M_d) using the expression given by [13]:

$$M_d = \frac{S_\lambda d^2}{k_\lambda B_\lambda(T_d)} \quad (1)$$

where d is the distance to the source, k_λ is the dust mass absorption coefficient and B_λ is the Planck function at a given dust temperature (T_d). We use Equation 1 to estimate the M_d values, listed in Table 2 for the galaxies in our sample. For our mass estimates, we assumed a T_d of 25 K and k_λ of 48 cm² g⁻¹. This T_d is a compromise value derived from values found for a sample of nearby galaxies [14].

We found dust masses in the range of $\sim 6.5\text{--}9.1 \times 10^7 M_\odot$ for NGC 2841, NGC 3184 and NGC 3351. NGC 3077 has a dust mass that is a factor of ~ 10 lower than the dust masses of the other galaxies included in our study. NGC 3077 is the nearest galaxy in our sample and its average size (observed at 70 μm) is a factor of 4.2–4.8 smaller than those (observed at 70 μm) of NGC 2841, NGC 3184 and NGC 3351. Based on this fact, the dust mass of NGC 3077 is expected to be a factor of 4.2–4.8 lower than the other galaxies, which contrasts with what has been previously estimated. This result implies that NGC 3077 is dust deficient, which may be caused by the tidal interactions that this galaxy suffers. Dust deficiency is also observed in cluster galaxies [5,6].

Table 2. Ellipse parameters and derived physical parameters for our sample of galaxies.

| Galaxy Name | Semi-Major Axis Arcsec | Semi-Minor Axis Arcsec | PA ¹ Degrees | $S_{70 \mu\text{m}}$ Jy Arcsec ² | $M_d \times 10^7 M_\odot$ | $M_{H_2+HI}^2 \times 10^9 M_\odot$ | Dust-to-Gas Ratio % |
|-------------|------------------------|------------------------|-------------------------|---|---------------------------|------------------------------------|---------------------|
| NGC 2841 | 140 | 70 | 150 | 24.4 | 6.6 | 9.0 | 0.7 |
| NGC 3077 | 30 | 22 | 65 | 36.1 | 0.7 | 0.04 | 17.5 |
| NGC 3184 | 140 | 100 | 117 | 34.5 | 6.5 | 3.7 | 1.8 |
| NGC 3351 | 110 | 110 | 163 | 65.3 | 9.1 | 3.2 | 2.8 |

¹ Parameter taken from the SIMBAD Astronomical Database. ² This mass is taken from the work of [12].

3.3. Dust-To-Gas Ratio

As mentioned in Section 3.1, the size of the ellipse used to derive the S_λ flux density is the same as that used to derive the CO luminosity by [12], who used this parameter to find the H₂ + HI mass (M_{H_2+HI}) for the galaxies in our study. These masses are listed in Table 2. Thanks to the similarity in the size of the ellipses, we are able to derive dust-to-gas ratios for the studied galaxies; values are given in Table 2. NGC 2841, NGC 3184 and NGC 3351 show an average dust-to-gas ratio of 1.8% that is consistent with the average value of $\sim 1\%$ found in a sample of nearby star-forming galaxies [9]. On the other hand, NGC 3077 has a dust-to-gas ratio of 17.5%, which is much higher than the average

value found for the other studied galaxies. The 17.5% ratio suggests that NGC 3077 is also deficient in $H_2 + HI$ gas, in addition to being more deficient in $H_2 + HI$ gas than dust. The authors of [7] found that HI gas is stripped more efficiently than dust in their study of a sample of spiral galaxies in a cluster environment, which is consistent with the scenario observed in NGC 3077.

4. Conclusions

We studied the dust content and dust-to-gas ratios of four nearby spiral galaxies. The main conclusions of our work are the following:

- For the isolated NGC 2841, NGC 3184 and NGC 3351 galaxies, we find dust masses in the range of $6.5\text{--}9.1 \times 10^7 M_\odot$. The dust masses of these galaxies are a factor of ~ 10 higher than the dust mass found for NGC 3077 affected by the tidal interactions of galaxies M81 and M82, indicating that NGC 3077 is a dust-deficient galaxy.
- NGC 3077 shows a dust-to-gas ratio of 17.5% that is much higher than the average value of 1.8% found for NGC 2841, NGC 3184 and NGC 3351. The ratio of 17.5% suggests that NGC 3077 is also deficient in $H_2 + HI$ gas, which has been stripped more efficiently than dust in this galaxy.

Author Contributions: J. Armijos-Abendaño, E. López, M. Llerena and F. Aldás performed the data analysis. J. Armijos-Abendaño and C. Logan wrote the manuscript. All authors contributed to the discussion and interpretation of the results.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Rasmussen, J.; Ponman, T.; Verdes-Montenegro, L.; Yun, M.S.; Borthakur, S. Galaxy evolution in Hickson compact groups: The role of ram-pressure stripping and strangulation. *Mon. Not. R. Astron. Soc.* **2008**, *388*, 1245–1264.
2. Mayer, L.; Mastropietro, C.; Wadsley, J.; Stadel, J.; Moore, B. Simultaneous ram pressure and tidal stripping: how dwarf spheroidals lost their gas. *Mon. Not. R. Astron. Soc.* **2006**, *369*, 1021–1038.
3. Nulsen, P.E.J. Transport processes and the stripping of cluster galaxies. *Mon. Not. R. Astron. Soc.* **1982**, *198*, 1007–1016.
4. Cowie, L.L.; Songaila, A. Thermal evaporation of gas within galaxies by a hot intergalactic medium. *Nature* **1977**, *266*, 501–503.
5. Giovanelli, R.; Haynes, M.P. Gas deficiency in cluster galaxies: a comparison of nine clusters. *Astrophys. J.* **1985**, *292*, 404–425.
6. Solanes, J.M.; Manrique, A.; García-Gómez, C.; González-Casado, G.; Giovanelli, R.; Haynes, M.P. The HI content of spirals. II. Gas deficiency in cluster galaxies. *Astrophys. J.* **2001**, *548*, 97–113.
7. Pappalardo, C.; Bianchi, S.; Corbelli, E.; Giovanardi, C.; Hunt, L.; Bendo, G.J.; Boselli, A.; Cortese, L.; Magrini, L.; Zibetti, S.; et al. The Herschel Virgo Cluster Survey. XI. Environmental effects on molecular gas and dust in spiral disks. *Astron. Astrophys.* **2012**, *545*, A75–A91.
8. Cortese, L.; Bekki, K.; Boselli, A.; Catinella, B.; Ciesla, L.; Hughes, T.M.; Baes, M.; Bendo, G.J.; Boquien, M.; de Looze, I.; et al. The selective effect of environment on the atomic and molecular gas-to-dust ratio of nearby galaxies in the Herschel reference survey. *Mon. Not. R. Astron. Soc.* **2016**, *459*, 3574–3584.
9. Sandstrom, K.M.; Leroy, A.K.; Walter, F.; Bolatto, A.D.; Crowell, K.V.; Draine, B.T.; Wilson, C.D.; Wolfire, M.; Calzetti, D.; Kennicutt, R.C.; et al. The CO-to- H_2 Conversion Factor and Dust-to-Gas Ratio on Kiloparsec Scales in Nearby Galaxies. *Astrophys. J.* **2013**, *777*, 5, doi:10.1088/0004-637X/777/1/5.
10. Walter, F.; Weiss, A.; Martin, C.; Scoville, N. The Interacting Dwarf Galaxy NGC 3077: The Interplay of Atomic and Molecular Gas with Violent Star Formation. *Astron. J.* **2002**, *123*, 225–237.
11. Kennicutt, R.C.; Calzetti, D.; Aniano, G.; Appleton, P.; Armus, L.; Beirão, P.; Bolatto, A.D.; Brandl, B.; Crocker, A.; Croxall, K. KINGFISH-Key Insights on Nearby Galaxies: A Far-Infrared Survey with Herschel: Survey Description and Image Atlas. *Publ. Astron. Soc. Pac.* **2011**, *123*, 1347–1369.
12. López, E.; Armijos-Abendaño, J.; Llerena, M.; Aldás, F. Upper limits to magnetic fields in the outskirts of galaxies. *Galaxies* **2017**, in press.

13. Hildebrand, R.H. The Determination of Cloud Masses and Dust Characteristics from Submillimetre Thermal Emission. *Q. J. R. Astron. Soc.* **1983**, *24*, 267–282.
14. Trehella, M.; Davies, J.I.; Alton, P.B.; Bianchi, S.; Madore, B.M. ISO Long Wavelength Spectrograph Observations of Cold Dust in Galaxies. *Astrophys. J.* **2000**, *543*, 153–160.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).