

Article

## Periodic CO<sub>2</sub> Dosing Strategy for *Dunaliella salina* Batch Culture

Kezhen Ying <sup>1,\*</sup>, D. James Gilmour <sup>2,†</sup> and William B. Zimmerman <sup>1,†</sup>

<sup>1</sup> Department of Chemical and Biological Engineering, University of Sheffield, Mappin Street, Sheffield S1 3JD, UK; E-Mail: w.zimmerman@sheffield.ac.uk

<sup>2</sup> Department of Molecular Biology and Biotechnology, University of Sheffield, Firth Court, Western Bank, Sheffield S10 2TN, UK; E-Mail: d.j.gilmour@sheffield.ac.uk

† These authors contributed equally to this work.

\* Author to whom correspondence should be addressed; E-Mail: kezhen.y@outlook.com.

Academic Editor: Christopher Q. Lan

Received: 20 March 2015 / Accepted: 7 May 2015 / Published: 19 May 2015

---

**Abstract:** A periodic CO<sub>2</sub> dosing strategy for *D. salina* 19/30 batch culture is proposed. A model of periodic CO<sub>2</sub> dosing including dosing time calculation, dosing interval estimation and final chlorophyll yield prediction was established. In experiments, 5% CO<sub>2</sub>/95% N<sub>2</sub> gas was periodically dosed into *D. salina* culture. Two different gas dosing flow rates were tested. The corresponding dosing time for each flow rate was estimated via the model (10 min·d<sup>-1</sup> for 0.7 L·min<sup>-1</sup> and 36 min·d<sup>-1</sup> for 0.3 L·min<sup>-1</sup>). Daily pH measurements showed that the pH of these cultures dosed periodically was always kept between 7.5 and 9.5, which highlights that periodic gas supply can maintain a suitable range of pH for microalgal growth without expensive buffers. Notably the culture dosed for set daily intervals was seen to have similar growth to the culture supplied constantly, but with much higher CO<sub>2</sub> capture efficiency (11%–18%) compared to continuous dosing (0.25%). It shows great potential for using periodic gas supply to reduce cost, wasted gas and energy use.

**Keywords:** periodic CO<sub>2</sub> dosing; *D. salina*; dosing time; dosing interval; CO<sub>2</sub> capture efficiency

---

## 1. Introduction

Currently, one emerging application of microalgae is for the fixation of CO<sub>2</sub> [1], for it may offer a way to reduce the levels of unwanted CO<sub>2</sub> whilst also allowing the production of useful by-products from the algae such as bio-oils, chemicals, fatty acids and substances, *etc.* [2]. However, the large-scale cultivation of microalgae still faces many problems and barriers. These include the energy required to continuously mix, dewater and process the algal biomass, and the large areas of land needed for such projects [3]. CO<sub>2</sub> dosing is one of the concerns highlighted in this study. The major problem with CO<sub>2</sub> being dosed into microalgae cultures is that conventionally the bubbles used are fairly large. As Zimmerman *et al.* [4] discuss, such bubbles will reduce the interfacial area between bubbles and the liquid, and will reduce the overall mass transfer for both CO<sub>2</sub> dissolution and O<sub>2</sub> stripping. To counteract the resulting low mass transfer, large flow rates of CO<sub>2</sub> enriched gas are often continuously bubbled into the cultures. While the mixing effect on the culture is beneficial for ensuring the even distribution of light and nutrients, the intense turbulence produced by large flow rate can damage the algal cells within the culture and reduce the productivity [5]. A further disadvantage of using high flow rate, especially with low mass transfer, is that most of the gas bubbled into the culture will pass through and be wasted. Carvalho and Malcata [6] agree that when CO<sub>2</sub> is bubbled into algal cultures, the mass transfer is not particularly effective and considerable gas is wasted, adding to operational costs. Even with some high mass transfer dosing techniques (e.g., Dissolved Air Flotation systems), hypothetically, continuous dosing is still not a wise option, because when the concentration of dissolved CO<sub>2</sub> has reached an equilibrium value, the gas-liquid mass transfer process stops. Any additional input beyond this point would not increase the total amount of dissolved CO<sub>2</sub>, but cost energy and waste CO<sub>2</sub>.

An innovative CO<sub>2</sub> microbubble-dosing technology was studied by Ying *et al.* [7–9], which has proved that the microbubble-dosing technology can enable a higher CO<sub>2</sub> mass transfer and consequently lead to a higher microalgal growth rate and a greater CO<sub>2</sub> capture efficiency. Nonetheless, the operational parameters (e.g., dosing time, dosing interval and flow rate, *etc.*) for microbubble-dosing technology still need to be engineered. In this study, an optimal periodic CO<sub>2</sub> dosing strategy is proposed and a model established based on *D. salina* (19/30) cultures. The main hypothesis for this study is that by using a microbubble driven airlift bioreactor, high mass transfer can be attained in a batch culture by supplying gas periodically (with little CO<sub>2</sub> wasted and less energy cost) and achieve similar algal growth compared to when the gas is supplied continuously.

## 2. Results and Discussion

### 2.1. Model of Periodical CO<sub>2</sub> Dosing

For an optimal periodical dosing strategy, three major principles need to be followed. Firstly, despite the exclusion of buffer solution and pH auto-regulating system, the pH of the culture needs to be controlled in a suitable range by periodic CO<sub>2</sub> dosing. Secondly, the dosing time should only be long enough for CO<sub>2</sub> to reach its equilibrium concentration. Meanwhile, the equilibrium pH (corresponding to CO<sub>2</sub> equilibrium concentration) is expected to be the lower limit of the suitable pH range for the microalgal species being utilized. Thirdly, the time period without dosing (dosing

interval) should ensure that the microalgae use up the dosed CO<sub>2</sub>, whilst ensuring that the pH increase to the upper limit is within the suitable pH range. Following these three principles, a culture with periodic dosing, compared to one with continuous dosing, should have sufficient (but not excess) CO<sub>2</sub> and an optimal pH range to support optimal growth (achieving similar growth as with continuous dosing), while with minimal amount of CO<sub>2</sub> wasted and less energy input. Based on these three principles, the dosing time, dosing interval and final algal yield can be estimated.

### 2.1.1. Estimation of Dosing Time

Assuming the suitable pH range for the culture of a particular microalga species is given between A and B, of which the corresponding concentration of [CO<sub>2</sub>] can be calculated by Equation (1) [7–11].

$$[\text{CO}_2] = \frac{(10^{-\text{pH}} - 10^{-(\text{pH}-14)} + \Delta[\text{Na}^+])10^{(-2\text{pH})}}{10^{(-6.381-\text{pH})} + 2 \times 10^{(-16.758)}} \quad (\text{mol/L}) \quad (1)$$

To control the culture in a target pH range the amount of CO<sub>2</sub> needed to be transferred to the medium can simply be estimated as the difference between dissolved CO<sub>2</sub> level at pH=B and pH=A (assuming the CO<sub>2</sub> uptake rate is negligible compared with CO<sub>2</sub> gas-liquid mass transfer rate). The dosing time is thereby calculated as the amount of CO<sub>2</sub> needed to be transferred to the medium divided by the average CO<sub>2</sub> mass transfer rate, shown in Equation (2), where  $v'_{\text{MTR}}$  represents CO<sub>2</sub> average mass transfer rate (mol·L<sup>-1</sup>·min<sup>-1</sup>) and can be calculated as Equation (3). The derivation of Equation (3) was explained in Ying *et al.* [9]. Finally, assuming the pH before dosing was controlled at B which gives the initial dissolved CO<sub>2</sub> level as [CO<sub>2</sub>]<sub>pH=B</sub>, the optimal valid dosing time needed to drop the pH from B to A can then be estimated through Equation (4), which is obtained by combining Equations (2) and (3). As long as the suitable pH range for a particular type of algae is given and the K<sub>La</sub> for a certain dosing condition is known, the optimal dosing time can be estimated.

$$t_d = \frac{[\text{CO}_2]_{\text{pH=A}} - [\text{CO}_2]_{\text{pH=B}}}{v'_{\text{MTR}}} \quad (2)$$

$$v'_{\text{MTR}} = \frac{K_L a ([\text{CO}_2]^* - [\text{CO}_2]_0)}{\frac{K_L a t_d}{2} + 1} \quad (3)$$

$$t_d = \frac{[\text{CO}_2]_{\text{pH=A}} - [\text{CO}_2]_{\text{pH=B}}}{K_L a ([\text{CO}_2]^* - [\text{CO}_2]_{\text{pH=B}}) - \frac{1}{2} K_L a ([\text{CO}_2]_A - [\text{CO}_2]_{\text{pH=B}})} \quad (4)$$

From the previous study carried out by Ying *et al.* [9], it was found that K<sub>La</sub> estimation is more accurate based on the changes in [C<sub>T</sub>], especially when pH > 8.4. Therefore, Equation (4) can also be written in terms of [C<sub>T</sub>], shown as Equation (5).

$$t_d = \frac{[C_T]_{\text{pH=A}} - [C_T]_{\text{pH=B}}}{K_L a ([C_T]^* - [C_T]_{\text{pH=B}}) - \frac{1}{2} K_L a ([C_T]_A - [C_T]_{\text{pH=B}})} \quad (5)$$

## 2.1.2. Estimation of Dosing Interval

The dosing interval here is defined as the time period without gas dosing. During this period, pH increases gradually because of the uptake of CO<sub>2</sub> by microalgae, until the pH achieves the upper limit of the suitable range, then dosing needs to be started again. Therefore, the effective estimation of dosing interval is crucial for periodic dosing, either too long or too short could cause the pH to exceed the upper or lower limit of the suitable range and adversely affect the algal growth.

The simplest way to estimate the dosing interval is to divide the amount of CO<sub>2</sub> expected to be absorbed by the CO<sub>2</sub> uptake rate. However, the instantaneous growth rate differs with the concentration of the algae [12], which leads to changes in CO<sub>2</sub> uptake rate. Instead of instantaneous CO<sub>2</sub> uptake rate, the average CO<sub>2</sub> uptake rate for the whole active growth period is therefore used to simplify the estimation of dosing interval Equation (6).

$$t_i = \frac{[\text{CO}_2]_{\text{pH=A}} - [\text{CO}_2]_{\text{pH=B}}}{v'_{\text{CO}_2\text{uptake}}} \quad (6)$$

Ying *et al.* [9] reported that when the pH was less than 8.4, the changes in the amount of total carbon almost all come from the changes in dissolved CO<sub>2</sub> [13], however, when pH was more than 8.4, the changes in dissolved CO<sub>2</sub> cannot fairly represent the CO<sub>2</sub> uptake by algae, as both HCO<sub>3</sub><sup>-</sup> and CO<sub>3</sub><sup>2-</sup> would generate dissolved CO<sub>2</sub> to compensate for the consumption of CO<sub>2</sub>. In other words, the amount of CO<sub>2</sub> consumed by algae should be more than the changes in dissolved CO<sub>2</sub>, as HCO<sub>3</sub><sup>-</sup> and CO<sub>3</sub><sup>2-</sup> would also contribute to the amount of CO<sub>2</sub> consumption. Therefore, the changes in total carbon [C<sub>T</sub>] should be considered instead of the changes in [CO<sub>2</sub>]. The [C<sub>T</sub>] can be calculated by Equation (7) [9]. Equation (6) should be converted into Equation (8). The dosing interval can then be estimated as long as the average CO<sub>2</sub> uptake rate is known.

$$[C_T] = [\text{CO}_2] + [\text{HCO}_3^-] + [\text{CO}_3^{2-}] = [\text{CO}_2] \cdot (1 + 10^{\text{pH}-6.381} + 10^{2\text{pH}-16.758}) \quad (7)$$

$$t_i = \frac{\Delta[C_T]}{v'_{\text{CO}_2\text{uptake}}} = \frac{[C_T]_{\text{pH=A}} - [C_T]_{\text{pH=B}}}{v'_{\text{CO}_2\text{uptake}}} \quad (8)$$

Since the periodic dosing strategy is proposed to achieve similar algal growth to when gas is supplied continuously, the average CO<sub>2</sub> uptake rate is assumed to be the same as in the culture with continuous or excessive CO<sub>2</sub> dosing. According to the information from previous *D. salina* cultures [7], the correlations for CO<sub>2</sub> uptake rate *versus* algal biomass concentration (measured as chlorophyll content) was described by Equation (9) [7], and the relation between total chlorophyll content increase and total CO<sub>2</sub> uptake was given (based on the cultures with excessive gas dosing) as Equation (10) [7].

$$v_{\text{CO}_2\text{uptake}} = 7 \times 10^{-5} \times [\text{Chl}] \quad (9)$$

$$\Delta[\text{Chl}] = 2703.4 \times \Delta[\text{CO}_2]_{\text{uptake}} = 2703.4 \times \Delta[\text{CO}_2]_{\text{dosed}} \quad (10)$$

For the same time period, Equation (10) can be transformed into Equation (11):

$$v'_{\text{Chl}} = 2703.4 \times v'_{\text{CO}_2\text{uptake}} \quad (11)$$

The average CO<sub>2</sub> uptake rate can be fairly described as:

$$v'_{\text{CO}_2\text{uptake}} = \int_{t_{c1}}^{t_{c2}} \frac{V_{\text{CO}_2\text{uptake}}}{t_c} dt \quad (12)$$

where  $t_c$  is the selected culture time period ( $t_{c2}-t_{c1}$ ), beginning from  $t_{c1}$  (the start of log growth phase) and ending by  $t_{c2}$  (the end of log growth phase).

Assuming the chlorophyll content ( $[\text{Chl}]$ ,  $\text{mg}\cdot\text{L}^{-1}$ ) is equal to the initial concentration ( $[\text{Chl}]_0$ ,  $\text{mg}\cdot\text{L}^{-1}$ ) plus the amount of its increase ( $v'_{\text{Chl}} \times t$ ),

$$[\text{Chl}] = [\text{Chl}]_0 + v'_{\text{Chl}} \cdot t \quad (13)$$

The average  $\text{CO}_2$  uptake rate can then be obtained by solving Equation (9), Equation (11)–(13), which gives:

$$v'_{\text{CO}_2\text{uptake}} = \frac{7 \times 10^{-5} \cdot [\text{Chl}]_0}{1 - 0.0946 \cdot t_c} \quad (14)$$

The dosing interval is then given by:

$$t_i = \frac{[C_T]_{\text{pH=A}} - [C_T]_{\text{pH=B}}}{7 \times 10^{-5} \cdot [\text{Chl}]_0} \cdot (1 - 0.0946 \cdot t_c) \quad (15)$$

Theoretically the dosing interval is better to be shortened as the algae grows, which may in practice increase the complexity of the time control process. Using a constant dosing interval through the whole log growth period can simplify the operating process. By doing so, one of the major concerns is that the pH level may exceed the upper limit of a target range. However, one magnitude of difference in the concentration ( $\text{mol}\cdot\text{L}^{-1}$ ) of dissolved  $\text{CO}_2$  only changes the pH by one unit [7], while the  $\text{CO}_2$  uptake rates for *D. salina* cultures are in the range of  $10^{-4}$  to  $10^{-3}$   $\text{mol}\cdot\text{L}^{-1}\cdot\text{d}^{-1}$ , therefore, the pH value may only increase over the upper limit of a target range by no more than 0.1 unit. Plus empirically the target pH range for algal culture can be slightly narrowed down during the dosing interval calculation in order to prevent the pH value exceeding the real upper limit. The second concern is dissolved oxygen accumulation. According to the basic photosynthetic reaction equation, the  $\text{O}_2$  generation rate equals to  $\text{CO}_2$  uptake rate. Therefore,  $10^{-3}$   $\text{mol}\cdot\text{L}^{-1}\cdot\text{d}^{-1}$  of  $\text{CO}_2$  uptake rate could result in around  $32$   $\text{mg}\cdot\text{L}^{-1}$  of daily  $\text{O}_2$  accumulation (300% over saturation with respect to air), which will diminish the rate of photosynthesis [14]. Thus, the dosing interval has to be limited to no more than 1 day so that Dissolved Oxygen (DO) can be removed by  $\text{CO}_2$  dosing in time. In terms of  $\text{O}_2$  stripping, it takes less than 10 min to reduce the dissolved oxygen from 32 to  $0.03$   $\text{mg}\cdot\text{L}^{-1}$  by microbubble ( $d_{32} = 388$   $\mu\text{m}$ ) dosing of 5%  $\text{CO}_2/95\%$   $\text{N}_2$  under  $0.7$   $\text{L}\cdot\text{min}^{-1}$ , according to the previous mass transfer study [7].

Statistically, from previous *D. salina* studies [7], the active growth phase usually lasted for 8 days (due to the light limitation). According to the literature, *D. salina* can tolerate a pH range of 5.5 to 10 [15]. A target pH range 7.5–9.5 was selected in this study. The dosing interval for *D. salina* cultures was estimated via Equation (15) to give approximately 1.5 days (based on  $\text{Chl}_0 = 5$   $\text{mg}\cdot\text{L}^{-1}$  and  $t_c = 8$  days). Conservatively, the dosing interval of 1 day was suggested for best results in practice, so that DO can be removed by  $\text{CO}_2$  dosing in time.

### 2.1.3. Prediction of Final Concentration of Chlorophyll Content

Once the dosing time  $t_d$  and dosing interval  $t_i$  are known, the total dosing time ( $t_{\text{Total}}$ ) through the culture period  $t_c$  can be calculated as:  $t_d \times t_c / (t_d + t_i)$ . The total amount of CO<sub>2</sub> uptake can be predicted by Equation (16):

$$\Delta[\text{CO}_2]_{\text{uptake(Total)}} = v'_{\text{MTR}} \times t_{\text{Total}} = \frac{K_L a ([\text{CO}_2]^* - [\text{CO}_2]_0)}{\frac{K_L a \cdot t_d}{2} + 1} \times \frac{t_d \cdot t_c}{t_d + t_i} \quad (16)$$

The final chlorophyll content can be estimated by Equation (17),

$$[\text{Chl}] = [\text{Chl}]_0 + \Delta[\text{Chl}] \quad (17)$$

By combining Equation (11), Equations (16) and (17), it gives:

$$[\text{Chl}] = [\text{Chl}]_0 + 2703.4 \times \frac{K_L a ([\text{CO}_2]^* - [\text{CO}_2]_0)}{\frac{K_L a \cdot t_d}{2} + 1} \times \frac{t_d \cdot t_c}{t_d + t_i} \quad (18)$$

Equations (16) and (18) are valid only when the dosing time is in a valid range ( $v'_{\text{MTR}} \times t_d \leq [\text{CO}_2]^*$ ).


## 2.2. Experimental Results

A set of *D. salina* batch cultures were carried out to test the hypothesis that sufficient CO<sub>2</sub> can be attained in a culture by “micro-bubbling” gas periodically (with little CO<sub>2</sub> wasted and less energy cost) whilst similar algal growth can be achieved compared to when the gas is supplied continuously, but with higher CO<sub>2</sub> capture efficiency.

Figure 1 shows the daily chlorophyll content of *D. salina* cultures supplied continuously and periodically with 5% CO<sub>2</sub>. As can be seen, the growth of these cultures, indicated by their chlorophyll content, appears to be fairly similar. This strongly supports the idea that *D. salina* growth is proportional to the total amount of CO<sub>2</sub> that has been effectively transferred from gas phase into liquid phase, while extra CO<sub>2</sub> dosing beyond the valid range does not improve the productivity. The final chlorophyll contents for the cultures with different dosing conditions were expected to be the same, as the total amount of CO<sub>2</sub> mass transfer was kept identical. The contrast between the growth of the cultures fed with 5% CO<sub>2</sub> and the control cultures can also be seen from Figure 1. Unsurprisingly the chlorophyll content of the control cultures remained much lower than other cultures, and a similar phenomenon was also observed in the study of Ying *et al.* [7].

Arguably the most important finding is the comparison that can be drawn between the growth of the *D. salina* cultures supplied continuously and daily with gas. It appears that there was no significant difference between the two types of culture. The daily chlorophyll content of continuously dosed culture seemed slightly higher between day 4 and day 11, which indicates the *D. salina* grew a bit faster under continuous dosing (approximately 1 day in advance of the periodic dosed cultures). However, the cultures engaged with periodic dosing model were still competitive to the continuously dosed culture, as they achieved a similar level of final chlorophyll content although with one day of delay. Regarding the CO<sub>2</sub> capture efficiency, the cultures with different dosing conditions were compared (Table 1). As can be seen, by applying a periodic dosing strategy, CO<sub>2</sub> capture efficiency

achieved is about 10%–20%. It is expected that the capture efficiency could be further enhanced by improving the CO<sub>2</sub> mass transfer (e.g., by further reducing the microbubble size). In contrast, with continuous dosing, capture efficiency was only 0.25% of CO<sub>2</sub> supplied, which indicates that most of the CO<sub>2</sub> was wasted rather than been captured. With a view to using microalgae for CO<sub>2</sub> sequestration, this will mean not only wasted energy to dose CO<sub>2</sub>, but also any CO<sub>2</sub> that was prevented from entering the atmosphere by fixation in the algae culture, will be greatly exceeded by the amount of CO<sub>2</sub> passing straight through the culture into the atmosphere. Therefore, this result shows the potential for both economic and energy savings by adopting a periodic dosing strategy, as it appears that similar algal growth to a culture supplied continuously with CO<sub>2</sub> can be achieved with periodic dosing, but with minimal CO<sub>2</sub> waste and minimal energy consumption on dosing.



**Figure 1.** Daily chlorophyll content of the *D. salina* cultures with different dosing conditions. The cultures with 0.7 L·min<sup>-1</sup> of periodic dosing were conducted in parallel, as were the control cultures. Therefore, the daily chlorophyll content for these two dosing conditions shown in this graph is the average value, with the error bars given separately.


**Table 1.** Comparisons of CO<sub>2</sub> capture efficiency for different dosing conditions.

Dosing Type (5% CO <sub>2</sub> )	Flow Rate (1 atm, 25 °C)	Dosing Time (in Total)	CO <sub>2</sub> Total		
			Absorption	Input	η
Periodic	0.7 L·min <sup>-1</sup>	10 min·d <sup>-1</sup> × 8 d	0.89 g	5.03 g	18%
Periodic	0.3 L·min <sup>-1</sup>	36 min·d <sup>-1</sup> × 8 d	0.86 g	7.75 g	11%
Continuous	0.3 L·min <sup>-1</sup>	24 h·d <sup>-1</sup> × 8 d	0.81 g	311.1 g	0.26%

NB: CO<sub>2</sub> input<sub>total</sub> = (CO<sub>2</sub>% × Q × t<sub>d</sub> × t<sub>c</sub> × P)/(R × T); CO<sub>2</sub> absorption = Δ[CO<sub>2</sub>]<sub>uptake</sub> Equation (16) × V<sub>L</sub>;  
 η = CO<sub>2</sub> absorption/CO<sub>2</sub> input.

Additionally, the pH control achieved using periodic dosing was also observed during these experiments (Figure 2). The pH in the culture supplied periodically with gas was maintained in the target region of 7.5–9.5 without the use of expensive buffers. This also indicates the periodic dosing model for dosing time and dosing interval estimation are accurate, so that the pH was controlled in an expected range. These results agree with the previous studies by Ying *et al.* [7] who conducted a similar experiment culturing *D. salina* in Airlift Loop Bioreactors (ALBs) proved  $30 \text{ min} \cdot \text{d}^{-1}$  of gas (5%  $\text{CO}_2$ , 95%  $\text{N}_2$ ).

Finally, the predicted final concentrations of chlorophyll content for different periodic dosing conditions Equation (18) were compared with the experimental results, shown in Table 2. The errors between theoretical values and experimental values were about 2%–3%, which indicates the accuracy of Equation (18).



**Figure 2.** Daily pH values for *D. salina* cultures supplied periodically with 5%  $\text{CO}_2$ . There are two pH values for each day, a higher one and a lower one, representing the pH values before and after  $\text{CO}_2$  dosing, respectively.

**Table 2.** Comparisons between estimated final concentrations of chlorophyll and real values for *D. salina* cultures with different dosing conditions.


Logarithmic Growth Time Period (d)	Dosing Condition	Estimated $[\text{Chl}]_t$ (mg/L)	Real $[\text{Chl}]_t$ (mg/L)	Error
8 d	$0.7 \text{ L} \cdot \text{min}^{-1}$ , $10 \text{ min} \cdot \text{d}^{-1}$	26.90	26.09	3%
8 d	$0.3 \text{ L} \cdot \text{min}^{-1}$ , $36 \text{ min} \cdot \text{d}^{-1}$	26.07	26.48	2%

### 3. Experimental Section

#### 3.1. Experimental Setup

The experimental setup is illustrated in Figure 3. Samples of the *D. salina* were pre-cultured in shake flask (100 mL culture in 250 mL flasks) in a  $25 \pm 2 \text{ }^\circ\text{C}$  growth room. The growth medium

composition is identical to the one described in Ying *et al.* [7], shown in Table 3. For the start of the main culture, 50 mL of this *D. salina* was added to 2.5 L of fresh culture medium in ALB. The dosing time ( $t_d$ ) and dosing interval ( $t_i$ ) were estimated based on Equations (5) and (15) and applied for the cultures engaged with periodic dosing strategy (No. 2, 4 and 5). pH was measured for each culture daily (for the culture dosed periodically, pH was measured twice per day, before and after dosing). 15 mL of sample for each culture was taken after gas dosing, followed by topping up the culture with 15 mL of fresh medium (For flask cultures, a sterilized glass stick was inserted into the culture for a proper stirring. Samples were taken after that.). The chlorophyll content for each sample was determined by measuring each sample's optical density for wavelengths of 645 and 663 nm using the same method to that used in Zimmerman *et al.* [16] and Ying *et al.* [7].



**Figure 3.** Setup for *D. salina* cultures. Six bioreactors containing *D. salina* cultures (50 mL of inocula to 2.5 L culture medium) were used for this experiment. Two of the six (No. 1 and 6) were flask cultures with no CO<sub>2</sub> enriched gas being bubbled through the culture. Like the other four cultures, these control cultures were kept in identical ALBs to ensure that the illumination through these cultures was the same as those being supplied with gas. The remaining four reactors were supplied with CO<sub>2</sub> enriched gas (5% CO<sub>2</sub>, 95% N<sub>2</sub>). Among them, No. 3 was dosed continuously with fine-bubbles, while No. 2, 4 and 5 were connected to a fluidic oscillator, dosed with microbubbles using periodic dosing strategy. No. 2 and 5 were conducted under same dosing condition for error analysis. The detailed dosing conditions for each reactor are listed in Table 4. The dosing time selected for each condition was estimated based on Equation (5). The detailed calculation is shown in Section 3.2. The temperature for each culture was maintained at ambient temperature around 24 °C. Two fluorescent lamps ( $90 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ ) were situated behind all reactors for illumination. Growth medium for *D. salina* cultures is shown in Table 3.

**Table 3.** *D. salina* culture medium.

Composition of Growth Medium Per Litre
0.5 M·NaCl; 0.185 mM·H <sub>3</sub> BO <sub>3</sub> ; 10 mM·KCl; 0.007 mM MnSO <sub>4</sub> ; 20 mM·MgCl <sub>2</sub> ; $0.8 \times 10^{-3}$ mM·ZnCl <sub>2</sub> ; 10 mM CaCl <sub>2</sub> ; $0.2 \times 10^{-4}$ mM CoCl <sub>2</sub> ; 24 mM MgSO <sub>4</sub> ; $0.2 \times 10^{-6}$ mM·CuCl <sub>2</sub> ; 5 mM NaNO <sub>3</sub> ; 24 mM·Na <sub>2</sub> SO <sub>4</sub> ; 0.0119 mM·NaHCO <sub>3</sub> ; 0.1 mM NaH <sub>2</sub> PO <sub>4</sub> ; 0.0015 mM·FeEDTA

**Table 4.** The dosing conditions for each culture. The average size for the microbubbles and fine bubbles were measured by using high speed camera [17].

Reactor	Dosing Conditions				Represent
	Bubbles	Fluidic Oscillator	Dosing Flowrate	Dosing Time Equation (5)	
No. 1	No bubbles	Not Engaged	0	0	Flask culture
No. 2	Microbubble (d <sub>32</sub> : 388 μm)	Engaged	0.7 L·min <sup>-1</sup>	10 min·d <sup>-1</sup>	Periodic dosing
No. 3	Fine-bubble (d <sub>32</sub> : 719 μm)	Not Engaged	0.3 L·min <sup>-1</sup>	24 h·d <sup>-1</sup>	Continuous dosing
No. 4	Microbubble (d <sub>32</sub> : 388 μm)	Engaged	0.3 L·min <sup>-1</sup>	36 min·d <sup>-1</sup>	Periodic dosing
No. 5	Microbubble (d <sub>32</sub> : 388 μm)	Engaged	0.7 L·min <sup>-1</sup>	10 min·d <sup>-1</sup>	Duplication of No. 2
No. 6	No bubbles	Not Engaged	0	0	Duplication of No. 1

### 3.2. Estimation of the Dosing Time and Dosing Interval

In Table 4, the dosing time for 0.3 and 0.7 L·min<sup>-1</sup> were calculated based on Equation (5). The detailed calculation was shown as follows.

Since the selected pH range for *D. S* culture was 7.5–9.5, and the concentration of NaHCO<sub>3</sub> in the culture medium was 0.0119 mol·L<sup>-1</sup>, [C<sub>T</sub>] at pH = 7.5 and pH = 9.5 can be calculated based on Equation (7), which gives: [C<sub>T</sub>]<sub>pH=7.5</sub> = 0.0128 mol·L<sup>-1</sup>; [C<sub>T</sub>]<sub>pH=9.5</sub> = 0.0106 mol·L<sup>-1</sup>.

The K<sub>LA</sub> for CO<sub>2</sub> mass transfer in the real algal culture was previously determined to be 0.044 min<sup>-1</sup> for 0.3 L·min<sup>-1</sup> of bubbling flow rate and 0.174 min<sup>-1</sup> for 0.7 L·min<sup>-1</sup> [9]. Finally, t<sub>a</sub> was calculated to be about 36 min for 0.3 L·min<sup>-1</sup> of bubbling flow rate and 10 min for 0.7 L·min<sup>-1</sup>.

For dosing interval estimation, it can be calculated by Equation (15).

Based on the assumption that Chl<sub>0</sub> = 5 mg·L<sup>-1</sup> and t<sub>c</sub> = 8 days, t<sub>i</sub> is calculated to be 1.5 days. Conservatively, 1 day of dosing interval was used in the experiment so that DO can be removed by CO<sub>2</sub> dosing in time.

## 4. Conclusions

A periodic CO<sub>2</sub> dosing strategy for *D. salina* culture is proposed, with a model of periodic CO<sub>2</sub> dosing including dosing time calculation, dosing interval estimation and final chlorophyll yield prediction established. The cultures applying periodic CO<sub>2</sub> dosing strategy provide a similar productivity to the culture with continuous dosing, but with a greater CO<sub>2</sub> sequestration efficiency. Due to the time limitation, only two different gas dosing flow rates were tested in this study. More flow rates can be tested to explore the relation between dosing flow rate and CO<sub>2</sub> capture efficiency in the future studies.

## Acknowledgments

William B. Zimmerman would like to thank the Royal Society for a Brain Mercer Innovation Award. We acknowledge support for microbubble dynamics from EPSRC EP/I019790/1. D. James Gilmour would like to acknowledge support from Carbon Trust.

## Author Contributions

Kezhen Ying, D. James Gilmour and William B. Zimmerman conceived the hypothesis and analyzed the data. Kezhen Ying D. James Gilmour designed/performed the experiments and wrote the paper. Kezhen Ying D. James Gilmour and William B. Zimmerman contributed materials/analysis tools and commended on the paper draft.

## Abbreviations

$\Delta[\text{Chl}]$	changes in the concentration of chlorophyll content, $\text{mg}\cdot\text{L}^{-1}$
$\Delta[\text{CO}_2]_{\text{dosed}}$	the difference between the concentration of dissolved $\text{CO}_2$ before and after dosing, $\text{mol}\cdot\text{L}^{-1}$
$\Delta[\text{CO}_2]_{\text{uptake}}$	changes in the concentration of $\text{CO}_2(\text{aq})$ due to the $\text{CO}_2$ consumption by algal growth, $\text{mol}\cdot\text{L}^{-1}$
$\Delta[\text{Na}^+]$	the concentration of sodium ions in the liquid which comes from $\text{NaHCO}_3$ , $\text{mol}\cdot\text{L}^{-1}$
$\text{CO}_2\%$	the concentration of $\text{CO}_2$ in the gas stream, % (v/v)
$d_{32}$	the Sauter mean diameter of the bubbles, $\mu\text{m}$
$\eta$	the $\text{CO}_2$ capture efficiency, %
$K_{La}$	volumetric mass transfer coefficient ( $\text{min}^{-1}$ ), where “ $K_L$ ” is the $\text{CO}_2$ gas-liquid mass transfer coefficient ( $\text{m}\cdot\text{min}^{-1}$ ); “ $a$ ” means the gas-liquid interfacial area ( $\text{m}^{-1}$ )
$P$	standard atmospheric pressure, 101.325 Kpa
$\text{pH}$	the pH value of the liquid
$R$	universal gas constant, $8.314 \text{ J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$
$T$	the temperature, K
$t_c$	culture time period, d
$t_d$	$\text{CO}_2$ dosing time, min
$t_i$	$\text{CO}_2$ dosing interval, min
$v'_{\text{Chl}}$	average productivity of chlorophyll content, $\text{mg}\cdot\text{L}^{-1}\cdot\text{d}^{-1}$
$v_{\text{CO}_2 \text{ uptake}}$	instantaneous algal $\text{CO}_2$ uptake rate, $\text{mol}\cdot\text{L}^{-1}\cdot\text{min}^{-1}$
$v'_{\text{CO}_2 \text{ uptake}}$	average algal $\text{CO}_2$ uptake rate, $\text{mol}\cdot\text{L}^{-1}\cdot\text{min}^{-1}$
$Q$	the gas dosing flowrate, $\text{L}\cdot\text{min}^{-1}$
$v'_{\text{MTR}}$	$\text{CO}_2$ average mass transfer rate, $\text{mol}\cdot\text{L}^{-1}\cdot\text{min}^{-1}$
$V_L$	the volume of the liquid, L
$[C_T]$	total concentration of inorganic carbon in the liquid, $\text{mol}\cdot\text{L}^{-1}$
$[C_T]_{\text{pH=A}}$	total concentration of inorganic carbon in the liquid at $\text{pH=A}$ , $\text{mol}\cdot\text{L}^{-1}$
$[C_T]_{\text{pH=B}}$	total concentration of inorganic carbon in the liquid at $\text{pH=B}$ , $\text{mol}\cdot\text{L}^{-1}$
$[\text{Chl}]$	concentration of chlorophyll content in the culture, $\text{mg}\cdot\text{L}^{-1}$
$[\text{Chl}]_0$	initial concentration of chlorophyll content at the beginning of log growth phase, $\text{mg}\cdot\text{L}^{-1}$
$[\text{CO}_2]$	dissolved carbon dioxide concentration in the liquid, $\text{mol}\cdot\text{L}^{-1}$
$[\text{CO}_2]^*$	dissolved carbon dioxide equilibrium concentration in the liquid, $\text{mol}\cdot\text{L}^{-1}$
$[\text{CO}_2]_0$	initial dissolved carbon dioxide concentration in the liquid, $\text{mol}\cdot\text{L}^{-1}$
$[\text{CO}_2]_{\text{pH=A}}$	dissolved carbon dioxide concentration in the liquid at $\text{pH=A}$ , $\text{mol}\cdot\text{L}^{-1}$
$[\text{CO}_2]_{\text{pH=B}}$	dissolved carbon dioxide concentration in the liquid at $\text{pH=B}$ , $\text{mol}\cdot\text{L}^{-1}$

[CO<sub>3</sub><sup>2-</sup>] the concentration of carbonate, mol·L<sup>-1</sup>  
[HCO<sub>3</sub><sup>-</sup>] the concentration of bicarbonate, mol·L<sup>-1</sup>

## Conflicts of Interest

The authors declare no conflict of interest.

## References

1. Wang, B.; Li, Y.; Wu, N.; Lan, C.Q. CO<sub>2</sub> bio-mitigation using microalgae. *Appl. Microbiol. Biotechnol.* **2008**, *79*, 707–718.
2. Pulz, O. Photobioreactors: Production systems for phototrophic microorganisms. *Appl. Microbiol. Biotechnol.* **2001**, *57*, 287–293.
3. Satyanarayana, K.G.; Mariano, A.B.; Vargas, J.V.C. A review on microalgae, a versatile source for sustainable energy and materials. *Int. J. Energy Res.* **2011**, *35*, 291–311.
4. Zimmerman, W.B.; Tesar, V.; Butler, S.; Bandulasena, H.C.H. Microbubble generation. *Recent Pat. Eng.* **2008**, *2*, 1–8.
5. Molina, E.; Fernandez, J.; Acien, F.G.; Chisti, Y. Tubular photobioreactor design for algal cultures. *J. Biotechnol.* **2001**, *92*, 113–131.
6. Carvalho, A.P.; Malcata, F.X. Transfer of carbon dioxide within cultures of microalgae: Plain bubbling versus hollow-fiber modules. *Biotechnol. Prog.* **2001**, *17*, 265–272.
7. Ying, K.; Gilmour, D.J.; Shi, Y.; Zimmerman, W.B. Growth enhancement of *Dunaliella salina* by microbubble induced airlift loop bioreactor (ALB)—The relation between mass transfer and growth rate. *J. Biomater. Nanobiotechnol.* **2013**, *4*, doi:10.4236/jbnt.2013.42A001.
8. Ying, K.; Gilmour, D.J.; Zimmerman W.B. Effects of CO<sub>2</sub> and pH on growth of the microalga *dunaliella salina*. *J. Microb. Biochem. Technol.* **2014**, *6*, 167–173.
9. Ying, K.; Mahmood, K.H.A.; James, H.; Gilmour, D.J.; Zimmerman, W.B. Enhanced mass transfer in microbubble driven airlift bioreactor for Microalgal culture. *J. Eng.* **2013**, *5*, 735–743.
10. Livansky, K. Losses of CO<sub>2</sub> in outdoor mass algal cultures: Determination of the mass transfer coefficient KL by means of measured pH course in NaHCO<sub>3</sub> solution. *Algol. Stud.* **1990**, *58*, 87–97.
11. Camacho, R.F. Prediction of dissolved oxygen and carbon dioxide concentration profiles in tubular photobioreactors for microalgal culture. *Biotechnol. Bioeng.* **1999**, *62*, 71–84.
12. Scragg, A.H. *Bioreactors in Biotechnology: A Practical Approach*; Ellis Horwood: London, UK, 1991; pp. 47–48.
13. Stemler, A. Forms of dissolved carbon dioxide required for photosystem II activity in chloroplast membranes. *Plant Physiol.* **1980**, *65*, 1160–1165.
14. Richmond, A. *Handbook of Microalgal Culture: Biotechnology and Applied Phycology*; Blackwell Science: Oxford, UK, 2008.
15. Borowitzka, M.A.; Borowitzka, L.J. *Dunaliella*. In *Micro-algal Biotechnology*; Borowitzka, M.A., Borowitzka, L.J., Eds.; Cambridge University Press: Cambridge, UK, 1988.

16. Zimmerman, W.B.; Zandi, M.; Bandulasena, H.C.H.; Tesar, V.; Gilmour, D.J.; Ying, K. Design of an airlift loop bioreactor and pilot scales studies with fluidic oscillator induced microbubbles for growth of a microalgae *Dunaliella salina*. *Appl. Energy* **2011**, *88*, 3357–3369.
17. Brittle, S.; Desai, P.; Ng, W.C.; Dunbar, A.; Howell, R.; Zimmerman, W.B. Optimal system design for minimising microbubble size through frequency control. *Chem. Eng. Res. Des.*, submitted for publication, 2015.

© 2015 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).