

Review

# The Paleomineralogy of the Hadean Eon Revisited

Shaunna M. Morrison <sup>1,\*</sup>, Simone E. Runyon <sup>1,2</sup> and Robert M. Hazen <sup>1</sup> 

<sup>1</sup> Geophysical Laboratory, Carnegie Institution for Science, 5251 Broad Branch Road NW, Washington, DC 20015, USA; rhazen@carnegiescience.edu

<sup>2</sup> Department of Geology and Geophysics, University of Wyoming, Laramie, WY 82071, USA; srunyon@uwyo.edu

\* Correspondence: smorrison@carnegiescience.edu

Received: 31 October 2018; Accepted: 7 December 2018; Published: 17 December 2018



**Abstract:** A preliminary list of plausible near-surface minerals present during Earth's Hadean Eon (>4.0 Ga) should be expanded to include: (1) phases that might have formed by precipitation of organic crystals prior to the rise of predation by cellular life; (2) minerals associated with large bolide impacts, especially through the generation of hydrothermal systems in circumferential fracture zones; and (3) local formation of minerals with relatively oxidized transition metals through abiological redox processes, such as photo-oxidation. Additional mineral diversity arises from the occurrence of some mineral species that form more than one 'natural kind', each with distinct chemical and morphological characteristics that arise by different paragenetic processes. Rare minerals, for example those containing essential B, Mo, or P, are not necessary for the origins of life. Rather, many common minerals incorporate those and other elements as trace and minor constituents. A rich variety of chemically reactive sites were thus available at the exposed surfaces of common Hadean rock-forming minerals.

**Keywords:** mineral evolution; photo-oxidation; impacts; paleomineralogy; Hadean Eon

## 1. Introduction

The paleomineralogy of Earth's earliest half-billion years, though poorly preserved in the known rock record, is important for understanding key evolutionary episodes, including the rate of planetary differentiation, the initiation of plate tectonics, the rate and nature of continental crust formation, and the origins of life. Accordingly, Hazen [1] proposed a preliminary list of 420 plausible mineral species that were likely present during Earth's Hadean Eon. These minerals are known to arise through purely physical and chemical processes—mechanisms such as fractional crystallization, regional metamorphism, serpentinization, evaporite crystallization, and hydrothermal alteration that are thought to have been active during Earth's first 500 million years, prior to the origin and global distribution of life.

Knowledge of the diversity and distribution of Hadean minerals is especially important for models of life's origins. Numerous mineral species and mineral groups—including oxides, silicates, carbonates, phosphates, sulfides, borates, and molybdates—have been invoked in models of life's origins [2–17]. The varied roles proposed for these minerals include catalysis for synthesis of essential biomolecules; protection of those molecules through surface interactions; molecular selection and concentration of specific subsets of molecules, including chiral selection, from a dilute prebiotic milieu; templating and formation of biopolymers; formation of lipid membranes; and surface redox chemistry that promoted life's earliest metabolism (e.g., [18–21]). It has been assumed that mineral species proposed in these origins-of-life scenarios must have been present in the near-surface environment of prebiotic Earth for the scenario to be valid.

In this contribution, we first re-examine the list of plausible Hadean minerals proposed by Hazen [1], expanding the list as a consequence of three important insights. The first group of plausible prebiotic mineral species on Earth—some of which are not known in Earth's near-surface environment today—include a variety of crystalline organic compounds, including amino acids, nucleobases, hydrocarbons, co-crystals, clathrates, and other species that could have occurred but have long since been consumed by ubiquitous cellular life [22,23].

Second, the profound potential influence of large bolide impacts on mineralization, though previously recognized as a possibility [24,25], has been elaborated by the work of Glikson and Pirajno (2018). In particular, we must consider additional mineral species formed in local hydrothermal systems related to impacts.

The third possible expansion of the list of Hadean minerals results from increasing recognition of abiological mechanisms for creating significant redox gradients at local, regional, and global scales. The prebiotic near-surface redox state of Earth was controlled primarily by the coexistence of both ferrous and ferric iron—i.e., conditions close to the hematite-magnetite oxygen buffer. However, significant excursions from that buffered state by several plausible natural processes may have led to the formation of minerals at much higher or lower oxygen fugacities.

Given these additional modes of mineral formation—parageneses not fully considered among the 420 species tabulated in the preliminary list of Hazen [1]—we catalog scores of additional species that might have occurred on Earth prior to life's origins. In addition, we suggest that the nature of mineral diversity, itself, needs to be reassessed based on the emerging concept of “natural kind clustering” applied to mineralogy [26]. Such common mineral species as calcite, quartz, hydroxylapatite, and pyrite can form through varied physical, chemical, and biological mechanisms—processes that result in distinct chemical and morphological characteristics of individual samples. In the context of planetary evolution, each different mode of formation of a mineral species should be treated as a distinct ‘natural kind’, with its own contextual story to tell.

Finally, we note that the absence of certain rare minerals in the Hadean Eon, for example species incorporating essential boron, molybdenum, or phosphorus, need not be an impediment to origins-of-life scenarios that invoke exotic mineral catalysts (e.g., [15]). These and many other rare elements are ubiquitous as trace and minor elements in rock-forming minerals. Therefore, numerous types of chemically reactive sites have been exposed at the surfaces of Earth's commonest minerals throughout geological history.

## 2. Prebiotic Organic Materials

Carbon is a relatively minor constituent of Earth's crust—less than 0.1 weight percent, most of which today is stored in extensive platforms of carbonate minerals [27–29]. However, recent speculations on the mineralogy of Saturn's large, hydrocarbon-rich moon Titan [22], suggest that organic crystals can be diverse and abundant, especially in an environment not subjected to predation by cellular life [23].

Most of the more than 50 ‘organic minerals’ that have been identified on Earth today [30,31] are biologically derived—crystals formed by the subaerial alteration of excrement (notably in cave guano deposits), from decaying vegetation and animal matter, or derived from coal and oil shale (e.g., [30,32,33]), including crystals that form near coal fires from condensing hot vapors [34–39]. Benner et al. [40] emphasize that many of these compounds are highly oxidized oxalates and carboxylates, which have low value as nutrients in the modern oxygen-rich environment and thus may persist in the presence of cellular life. None of these extant organic phases were included in the preliminary list of Hadean minerals.

By contrast, many organic crystals that might have occurred on prebiotic Earth, either by endogenous synthesis or exogenous delivery, represent food that today would be consumed quickly. Early in Earth's history, when carbon-rich chondrites and comets transported abundant organic molecules to the sterile surface [41–46] and subsurface organic synthesis occurred in carbon-rich

aqueous fluids [47–52], numerous now extinct organic crystals may have existed. Table 1 lists representative potential organic compounds that might have occurred as crystalline minerals on prebiotic Earth.

**Table 1.** Examples of representative crystalline organic compounds that might have occurred during Earth’s Hadean Eon (>4.03 Ga) as a consequence of prebiotic endogenous production or exogenous delivery of soluble or volatile organic molecules, followed by molecular selection, concentration, and crystallization.

Molecular Class	Name
Amino acids	glycine
	alanine (D and L)
	β-alanine
	isovaline
Nucleobases	aspartate (D and L)
	adenine
	guanine <sup>1,*</sup>
	uracil
Polycyclic Aromatic Hydrocarbons	fluorene <sup>2,*</sup>
	anthracene <sup>3,*</sup>
	pentacene <sup>4,*</sup>
	coronene <sup>5,*</sup>
	pyrene
	chrysene
Other Organics	tetracene
	ferrocyanide
	2-aminooxazole

\* Compounds also known as organic minerals. <sup>1</sup> Guanine; <sup>2</sup> kratochvilite; <sup>3</sup> ravatite; <sup>4</sup> idrialite; <sup>5</sup> carpathite.

While details of Hadean organic materials will likely remain speculative, concentration of soluble molecular species in shallow water environments, followed by evaporation and precipitation, could have led to crystals of the most abundant amino acids, including glycine, alanine, and aspartate [46]; nucleobases such as adenine, guanine, and uracil [45]; carbohydrates [53,54]; varied hydrocarbons including alkanes, alkenes, and polycyclic aromatics; and other key biomolecules (Table 1). Of note is the possibility that some organic molecules precipitated as chirally pure ‘right-handed’ and ‘left-handed’ crystalline forms [55–58]. Furthermore, given the widespread influence of volcanic activity and associated vents, a variety of organic crystals, including polycyclic aromatic and aliphatic hydrocarbons, may have condensed from a hot gas phase as they do near coal fires today.

Maynard-Caseley et al. [22] also explored the fascinating diversity of organic ‘co-crystals’ with two or more small molecular building blocks. Their proposed examples focused on phases such as 1:1 acetylene:methanol, 2:1 acetylene:acetonitrile, and 3:1 benzene:ethane, consistent with 92 K surface environment of Titan. The possibility of higher-temperature molecular co-crystals, especially disordered sub-crystalline arrangements of varied lipid or polycyclic aromatic hydrocarbon molecules, should be considered. Organic-enclosing water- or silica-based clathrate compounds at low temperatures [59–61], including clathrates intercalated with clay minerals [62], represent additional potential Hadean species.

Note that prebiotic organic compounds have been invoked in chemical scenarios of life’s origins [40,63]. Benner et al. [40] consider reactive carbohydrates and related nitrogen-bearing compounds, while Ritson and Sutherland [64] focus on RNA assembly in the presence of crystalline ferrocyanide—phases that should be added to lists of plausible Hadean minerals. In addition, Sutherland and colleagues have studied 2-aminooxazole, a plausible prebiotic organic compound that can serve as an intermediate molecule in RNA synthesis [65,66].

An additional intriguing possibility is that a complex suite of organic microcrystals still exists in carbonaceous chondrite meteorites, which may have experienced extreme dehydration and thus

incorporate a rich variety of organic and inorganic “salts” [67]. Standard analyses would not easily detect these submicroscopic crystalline zones, so the opportunity for new discoveries of preserved ancient organic crystals may exist.

### 3. Impact Mineralization

Major impacts of asteroids and comets have the potential to affect the mineral diversity of a terrestrial planet or moon in at least four ways. On the one hand, such impacts may have temporally reduced mineral diversity, at least at a local scale, by disrupting the crust and melting and/or vaporizing minerals near the impact site. Hadean Earth may have experienced tens of thousands of impacts of objects greater than 1 km in diameter, with hundreds of impacts greater than 10 kilometers in diameter [68]. Such events represented significant ‘reset buttons’ for local mineralogy.

On the other hand, large impacts likely enhanced mineral diversity in three significant ways. (1) The presence of high-pressure shock minerals, such as the ‘lonsdaleite’ allotrope of carbon and the dense coesite and stishovite forms of SiO<sub>2</sub>, are commonly associated marker minerals for impact events. (2) Large impacts have the potential to excavate and scatter rocks from deep terrains, thus bringing to the surface high-pressure-temperature metamorphic minerals that would otherwise not be seen. (3) Impacts can create significant hydrothermal systems in circumferential fracture zones that persist for upwards of 100,000 years, thus creating local mineralizing environments (e.g., [25]).

The first two of these mechanisms were recognized and incorporated by Hazen [1] in a list of plausible Hadean minerals. However, the possibility of extensive hydrothermal zones, though previously explored by several authors [25,69,70], was less fully developed. New contributions, summarized by Glikson and Pirajno (2018), elaborate on this idea and point to several likely Hadean minerals that should be added to the list (Table 2).

The significance of impact-produced fracture zones lies in their potential geographic and temporal extent. Deep circumferential hydrothermal systems up to hundreds of kilometers in diameter could have persisted for hundreds of thousands of years [25,69,70]. These large-scale systems of fluid–rock interactions likely pre-dated significant hydrothermal systems associated with subduction and arc volcanism. The mineralogical consequences of such Hadean impact-related processes remain speculative, but the occurrence of significant local hydrothermal alteration, including production of zeolites and clay minerals, seems likely [71,72]. In addition, the selection and concentration of relatively rare chemical elements by hydrothermal fluids would have led to the emplacement of ore bodies, including those featuring abundant chalcogenide and sulfosalt minerals. More recent examples of impact-associated ore deposits, though probably not exact analogies of the Hadean context, have been documented by the Ni-Cu mineralization of the Sudbury Igneous Complex in Canada [73], Pb-Zn-Ag-Ba ores of the Siljjan Ring Complex in Sweden [74], and several Australian mines [68,72].

The question of what new minerals might have arisen as a consequence of these localized hydrothermal systems must remain speculative. However, a major impact on mafic or ultramafic lithologies, especially when coupled with the input of a bolide concentrated in iron, nickel, and other siderophile elements, as well as sulfur and other chalcophile elements, points to a potentially richer assembly of primary and secondary minerals than suggested by Hazen (2013). Notable additions to the preliminary list of 420 species could include minerals of first-row transition metals, including Ti, V, Cr, Mn, Fe, Co, Ni, Cu, and Zn, as well as As, Se, Mo, Ag, Sb, Au, Pb, Bi, and platinum group elements—elements commonly associated with sulfide deposits. Accordingly, Table 2 lists an additional 33 plausible Hadean mineral species, all of which are less common than those proposed by Hazen (2013), but which are frequently associated with hydrothermal ore bodies today.

**Table 2.** Plausible Hadean Eon (>4.03 Ga) minerals: primary sulfides, chalcogenides, and sulfosalts that might be associated with hydrothermal fracture zones near large bolide impacts. The following minerals were not included in the preliminary list of Hazen (2013).

Group	Species	IMA Formula	
Sulfides, Chalcogenides	argentopentlandite	Ag(Fe,Ni) <sub>8</sub> S <sub>8</sub>	
	argentopyrite	AgFe <sub>2</sub> S <sub>3</sub>	
	bismuthinite	Bi <sub>2</sub> S <sub>3</sub>	
	breithauptite	NiSb	
	clausthalite	PbSe	
	dyscrasite	Ag <sub>3+x</sub> Sb <sub>1-x</sub> (x ≈ 0.2)	
	geffroyite	(Cu,Fe,Ag) <sub>9</sub> Se <sub>8</sub>	
	maucherite	Ni <sub>11</sub> As <sub>8</sub>	
	naumannite	Ag <sub>2</sub> Se	
	siegenite	CoNi <sub>2</sub> S <sub>4</sub>	
	stibnite	Sb <sub>2</sub> S <sub>3</sub>	
	Sulfosalts	aikinite	CuPbBiS <sub>3</sub>
		andorite	AgPbSb <sub>3</sub> S <sub>6</sub>
		berthierite	FeSb <sub>2</sub> S <sub>4</sub>
		boulangierite	Pb <sub>5</sub> Sb <sub>4</sub> S <sub>11</sub>
bournonite		CuPbSbS <sub>3</sub>	
chalcostibite		CuSbS <sub>2</sub>	
cosalite		Pb <sub>2</sub> Bi <sub>2</sub> S <sub>5</sub>	
emlectite		CuBiS <sub>2</sub>	
jamesonite		Pb <sub>4</sub> FeSb <sub>6</sub> S <sub>14</sub>	
lindströmite		Pb <sub>3</sub> Cu <sub>3</sub> Bi <sub>7</sub> S <sub>15</sub>	
matildite		AgBiS <sub>2</sub>	
proustite		Ag <sub>3</sub> AsS <sub>3</sub>	
pyrargyrite		Ag <sub>3</sub> SbS <sub>3</sub>	
ramdohrite		Ag <sub>3</sub> Pb <sub>6</sub> Sb <sub>11</sub> S <sub>24</sub>	
samsonite		Ag <sub>4</sub> MnSb <sub>2</sub> S <sub>6</sub>	
seligmannite		CuPbAsS <sub>3</sub>	
smithite		AgAsS <sub>2</sub>	
sulvanite		Cu <sub>3</sub> VS <sub>4</sub>	
tealite	PbSnS <sub>2</sub>		
tetrahedrite	Cu <sub>6</sub> [Cu <sub>4</sub> (Fe,Zn) <sub>2</sub> ]Sb <sub>4</sub> S <sub>13</sub>		
ullmanite	NiSbS		
zinkenite	Pb <sub>9</sub> Sb <sub>22</sub> S <sub>42</sub>		

Among the 33 additional possible Hadean mineral species in Table 2 are relatively common sulfides, chalcogenides, and sulfosalts—primary hydrothermal minerals that incorporate elements that might have been mobilized in large impact zones. An additional 25 oxysalt and halide minerals, most of them secondary, appear in Table 3. Of special interest in terms of plausible Hadean hydrothermal minerals are molybdates, which have been proposed by Benner and colleagues [16] to play special chemical roles in catalytic synthesis of carbohydrate precursors of ribonucleosides and nucleotides. In the previous list, molybdenite (MoS<sub>2</sub>) was the only Hadean molybdenum-bearing mineral. Most molybdates are known principally as secondary minerals in the oxidized zones of ore bodies, and thus were excluded from consideration. However, here we consider the possibility of powellite (CaMoO<sub>4</sub>), a mineral known to occur in association with basalt, tachite, and granite pegmatite [75].

**Table 3.** Plausible Hadean Eon (>4.03 Ga) minerals: mostly secondary minerals, including those formed as a result of significant local redox excursions, for example those associated with photo-oxidation. The following minerals were not included in the preliminary list of Hazen (2013).

Group	Species	IMA Formula	
Oxides, Hydroxides, Halides	akaganeite	$\text{Fe}^{3+}\text{O}(\text{OH},\text{Cl})$	
	arsenolite	$\text{As}_2\text{O}_3$	
	cotunnite	$\text{PbCl}_2$	
	cuprite	$\text{Cu}_2\text{O}$	
	hibbingite	$\text{Fe}^{2+}_2(\text{OH})_3\text{Cl}$	
	lawrencite	$\text{FeCl}_2$	
	zirkelite	$(\text{Ti},\text{Ca},\text{Zr})\text{O}_{2-x}$	
	Silicates	allanite-(La)	$\text{CaLa}(\text{Al}_2\text{Fe}^{2+})[\text{Si}_2\text{O}_7][\text{SiO}_4]\text{O}(\text{OH})$
		allanite-(Y)	$\text{CaY}(\text{Al}_2\text{Fe}^{2+})[\text{Si}_2\text{O}_7][\text{SiO}_4]\text{O}(\text{OH})$
	Carbonates	azurite	$\text{Cu}_3(\text{CO}_3)_2(\text{OH})_2$
malachite		$\text{Cu}_2(\text{CO}_3)(\text{OH})_2$	
pyroaurite		$\text{Mg}_6\text{Fe}^{3+}_2(\text{CO}_3)(\text{OH})_{16}\cdot 4\text{H}_2\text{O}$	
Sulphates, Arsenates, etc.	brochantite	$\text{Cu}_4(\text{SO}_4)(\text{OH})_6$	
	chalcantite	$\text{Cu}(\text{SO}_4)\cdot 5\text{H}_2\text{O}$	
	copiapite	$\text{Fe}^{2+}\text{Fe}^{3+}_4(\text{SO}_4)_6(\text{OH})_2\cdot 20\text{H}_2\text{O}$	
	erythrite	$\text{Co}_3(\text{AsO}_4)_2\cdot 8\text{H}_2\text{O}$	
	jarosite	$\text{KFe}^{3+}_3(\text{SO}_4)_2(\text{OH})_6$	
	magnesiocopiapite	$\text{MgFe}^{3+}_4(\text{SO}_4)_6(\text{OH})_2\cdot 20\text{H}_2\text{O}$	
	morenosite	$\text{Ni}(\text{SO}_4)\cdot 7\text{H}_2\text{O}$	
	nickelhexahydrite	$\text{Ni}(\text{SO}_4)\cdot 6\text{H}_2\text{O}$	
	römerite	$\text{Fe}^{2+}\text{Fe}^{3+}_2(\text{SO}_4)_4\cdot 14\text{H}_2\text{O}$	
	scorodite	$\text{Fe}^{3+}(\text{AsO}_4)\cdot 2\text{H}_2\text{O}$	
	Molybdate, Tungstate, Phosphate	powellite	$\text{CaMoO}_4$
scheelite		$\text{CaWO}_4$	
xenotime-(Y)		$\text{YPO}_4$	

#### 4. Prebiotic Redox Gradients on Earth

Planets create redox gradients at varied scales from local to global. An assumption underlying virtually all hypotheses of life's origins and evolution is that ancient life exploited those gradients as a source of chemical potential energy. Only much later did life, especially cells engaged in oxygenic photosynthesis, begin to create and sustain significant redox gradients of their own. The hypothesis that thousands of oxidized mineral species represent a global-scale mineralogical biosignature (e.g., [76]) is based on the assumption that significant near-surface redox gradients did not exist before the ~2.5 Ga 'Great Oxidation Event'. However, this assumption might be challenged on the grounds that planets can display purely physico-chemical processes that create redox gradients at various scales, thus potentially generating extremely oxidized or reduced assemblages of minerals as electrons are shifted from a more oxidized to a more reduced environment.

Explorations of the nature and extent of prebiotic redox gradients on Hadean Earth deserve a detailed quantitative analysis with estimates of the magnitude and duration of such redox excursions. Here we present a qualitative summary and suggest a number of electron transfer mechanisms by which such redox gradients may have arisen before the origins of life. For convenience, such mechanisms can be divided into global-, regional-, and local-scale phenomena.

##### 4.1. Global-Scale Redox Gradients

At the largest-scale of planetary differentiation, the immiscibility of silicate- and metal-rich melts, and the consequent gravitational segregation of a metal-rich core, creates a significant redox gradient at the core-mantle boundary and a permanent oxidation of Earth's outer layers relative to the core. This mechanism, by which electrons are sequestered in the native iron-nickel alloys of the deepest interior, suggest that Earth's mantle approached its current oxidation state by 3.5 to 4.0 Ga. Geochemical evidence from both isotopes and trace elements (e.g., [77–82]), coupled with the pressure

effect on oxygen fugacity (e.g., [83,84]), suggest that biology did not play a role in this planetary-scale interior redox distribution.

Additional large-scale development of redox gradients can occur when upper atmosphere photo-dissociation of water and subsequent hydrogen escape to space leads to a gradual increase in atmospheric oxygen fugacity. The ~0.15 mole fraction of molecular oxygen in the martian atmosphere is a consequence of this planetary-scale process [85]—a physicochemical effect that creates a redox gradient independent of life [86]. Global changes in atmospheric redox might also occur through “impact erosion” [87]. Nevertheless, most models of Earth’s Hadean atmosphere suggest that it was significantly reduced compared to today’s atmosphere [88–91], with a surface environment sustained close to the hematite-magnetite buffer (i.e.,  $f_{\text{O}_2} \sim 10^{-7.2}$ ; e.g., [92,93]). These global-scale redox gradients were incorporated in the analysis of Hazen [1] and do not lead to any new predicted prebiotic mineral species.

#### 4.2. Regional-Scale Redox Gradients

Regional-scale (>1 km) movements of fluids and crystals can generate redox gradients in a variety of ways [94–96]. For example, redox gradients arise from crystal fractionation and settling of denser  $\text{Fe}^{2+}$ -rich olivine during the formation of dunite [97–102]. This process creates a redox gradient at the scale of a magma chamber. Similarly, the crystallization of magnetite within a magma body results in redox changes of a melt with the increased rate of total Fe and particularly  $\text{Fe}^{3+}$  removal from a magma (e.g., [103]). This redox change due to magnetite crystallization is commonly accompanied by abrupt sulfur saturation and can potentially lead to the formation of an immiscible sulfide melt (e.g., [104,105]).

Many examples of immiscibility within natural magmatic systems are known, and compositional differences resulting from unmixing can be extreme (e.g., [106]), potentially creating redox gradients. Immiscibility between felsic and mafic silicate liquids has been documented in layered mafic intrusions, anorthosite complexes, mid-ocean ridge magma chambers, and granitoids (e.g., [107]).

Even in situations in which one immiscible phase is volumetrically minor compared to the other, it may be extremely effective in extracting and transporting large amounts of trace and rare elements, such as the case for immiscible sulfide melts concentrating PGEs and other chalcophile elements such as Cu and Ni from a larger silicate melt body [106,108–110]. The more reduced sulfide melts, segregated from more oxidized silicate melts, are commonly called upon as an important ore-forming process in magmatic Ni-Cu  $\pm$  PGE deposits [110–116].

Similarly, carbonate-silicate liquid immiscibility has been invoked in the formation of carbonatite magma (e.g., [117–120]), though the extent to which such immiscibility leads to redox gradients is not fully resolved. Na, Ca, P, and F partition into carbonate melt, while Mg, Mn, Fe, Al, Si, and Ti concentrate in the silicate melts. Some experimental work has shown that oxidized sulfur ( $\text{SO}_4^{2-}$ ) and uranium ( $\text{U}^{6+}$ ) preferentially partition into carbonate melt (along with Ca, P, Nb, Ta, and Pb), whereas reduced sulfur and uranium partition into silicate melt [120]. However, other analyses point to the derivation of more reduced carbonate from more oxidized silicate melts [109,121–126].

Another controversy relates to the possibility that anorthosite and other mafic/ultramafic silicate melts may separate from immiscible Fe-Ti oxide melts [127–132]. Arsenide melt immiscibility is a special case that occurs when mafic or ultramafic magmas become saturated in As [133–137].

These regional processes not only create redox gradients as fluids and/or crystals separate, but they also have the potential to produce new mineral species in zones of relatively high or low oxidation states. On a potentially larger scale, recent experimental work has suggested that immiscibility between  $\text{H}_2\text{O}$  and  $\text{H}_2$  may occur in the mantle and may provide a mechanism for the rapid oxidation of the upper mantle immediately following core formation (e.g., [138]).

### 4.3. Local-Scale Redox Gradients

At least five non-biological, near-surface mechanisms, all prevalent during the Hadean Eon, would have produced significant redox excursions at the scale of millimeters to meters: (1) fine-scale redox gradients resulting from magma segregation; (2) meteorite falls; (3) lightning strikes, notably near active volcanoes; (4) seafloor venting of sulfide-rich hydrothermal fluids; and (5) photo-oxidation of exposed minerals. Meteorites, lightning, and hydrothermal vents would likely have created locally reduced environments, whereas photo-oxidation has the potential to promote excursions to significantly more oxidized mineral-forming environments.

#### 4.3.1. Local Immiscibility

Just as immiscibility (or magma unmixing) may form regional-scale redox gradients, separation of fluids may be a powerful mechanism for forming local redox gradients. Recent experimental studies (e.g., [139,140]) have created localized redox gradients at the scale of hundreds of microns to millimeters. For example, redox gradients are proposed to form due to ferrous iron liberation during clinopyroxene and orthopyroxene dissolution in a more mafic melt, coupled with the diffusion of that ferrous iron into a more felsic melt. Local redox gradients are enhanced by the charge imbalance resulting from diffusion of charged ions across a magma mixing interface [139]. These localized redox gradients may be large (e.g., almost two orders of magnitude  $fO_2$ ), develop rapidly, and remain in this dynamic equilibrium on a time scale of days to potentially weeks [140]. Felsic magmas were unlikely to be common in the Hadean Eon. However, the occurrence of ancient detrital zircon grains in the Jack Hills sediments of Western Australia supports the existence of granitic crust on earth as early as 4.4 Ga [141–143]. Whether these features formed from reworking through interaction with preexisting Hadean crust or from impact melting (e.g., [144,145]), the possibility exists for development of local-scale redox gradients forming from magma mixing events in the Hadean Eon.

#### 4.3.2. Meteorites

Relatively reducing environments would have occurred in the vicinity of falls of certain classes of meteorites, including irons, stony irons, and enstatite chondrites. Enstatite chondrites, in particular, incorporate a suite of extremely reduced mineral phases, including silicides, phosphides, nitrides, and carbides. Such phases in a shallow aqueous environment create environments of extreme chemical disequilibrium that are unlikely to be caused by other near-surface phenomena. Pasek and colleagues [13,146–148] have focused special attention on prebiotic aqueous reactions between various organic species and schreibersite  $[(Fe,Ni)_3P]$ , a common meteorite phosphide mineral.

#### 4.3.3. Lightning Strikes

Lightning strikes, though highly localized, are responsible for significant mineralogical novelty as violent pulses of electrons and associated heating to  $>2000$  K can cause dielectric breakdown of rock-forming minerals. The associated conditions are extremely reducing, approximating the  $SiO_2$ -Si buffer [149,150]. The most familiar mineralogical consequences of lightning are fulgurites, which form when cloud-to-ground lightning alters soils and rocks. Among the mineralogical novelties found in some fulgurites are such reduced phases as native silica and a variety of iron silicides [149,151], reduced phosphorus compounds [152,153], and fullerene-type allotropes of carbon [154]. Of special note in the context of prebiotic chemistry are the effects of volcanic lightning [155], which would have been more prevalent in the Hadean Eon.

#### 4.3.4. Hydrothermal Vents

Ocean floor hydrothermal vents through mafic and ultramafic lithologies represent a net electron flow from mantle to crust, with measured potentials up to 600 millivolt at an active vent [156]. Furthermore, Ang et al. [157] observed direct thermoelectric conversion in natural chalcopyrite

(CuFeS<sub>2</sub>) from a sulfide-rich ‘black smoker’ hydrothermal vent system in the southern Lau Basin at the Australian-Pacific plate boundary. The resulting sulfide-rich metal deposits incorporate a variety of transition metal minerals [158–160] and represent a significant worldwide economic resource [159,161].

#### 4.3.5. Photo-Oxidation

The examples above describe local-scale reducing conditions caused by immiscible fluids/magmas, meteorites, lightning, or hydrothermal vents. None of these processes is likely to result in significant additions to the lists of plausible Hadean minerals. By contrast, photo-oxidation by exposure to ultraviolet radiation at Earth’s unshielded ancient surface, though poorly studied in the context of mineral evolution, might have led to local production of more oxidized minerals—notably minerals that form at oxygen fugacities significantly greater than that of the hematite-magnetite buffer.

In some instances, photo-oxidation of a mineral that incorporates a reduced transition metal can lead to a reduced product plus a new mineral with a more oxidized transition metal. For example, Kim et al. [162] report on the photo-oxidation of siderite to form hydrogen and ferric iron oxides—a reaction similar to that of UV irradiation of aqueous Fe<sup>2+</sup> ions [163–165].

Special attention has been focused on possible photocatalytic roles of the surfaces of rutile and anatase polymorphs of TiO<sub>2</sub> [166–169]. When exposed to UV radiation, these minerals have been shown to photo-oxidize water, as well as a variety of organic species. Titanium dioxide is also capable of oxidizing nitrogen compounds, including di-nitrogen gas, to produce nitrates and nitrites [170,171]—compounds that could lead to local concentrations of minerals such as niter [K(NO<sub>3</sub>)] and nitratine [Na(NO<sub>3</sub>)]. In this regard, the surface of Mars today provides a possible analog for early Earth, as photo-oxidation has evidently led to the production of peroxides and perchlorates in martian soils [172–175]. In spite of these studies, much work remains to document the mineralogical consequences of photo-oxidation.

#### 4.4. ‘Natural Kinds’ of Hadean Minerals

All prior mineral evolution analyses, including discussions of the diversity of Earth’s Hadean mineralogy, were considered in the context of the formal definition of mineral species established by the International Mineralogical Association or “IMA” [176,177]. The IMA methodology assigns each mineral ‘species’ a unique combination of idealized end-member composition and crystal structure—a protocol that has led to the approval of more than 5400 mineral species (for a current list see: [rruff.info/ima](http://rruff.info/ima); [178]).

As useful as this unambiguous classification system continues to be, it fails to adequately capture the true diversity of mineral ‘natural kinds’—the demonstrably distinct occurrences of minerals. In the context of planetary evolution, particularly when surveying the mineral diversity of a specific period of Earth history, at least three types of planetary materials are imperfectly catalogued by the present system of mineral classification [26]: (1) some distinct kinds of minerals have been lumped according to the IMA classification; (2) some natural kinds of minerals have been split into multiple species by the IMA classification; and (3) noncrystalline materials are not included. Consider examples of each of these situations.

##### 4.4.1. Minerals That Have Been Lumped Together

Some mineral species occur in multiple distinctive ‘natural kinds’. For example, the species ‘diamond’, which is defined by the IMA as pure carbon in the diamond crystal structure, forms in at least five distinct natural kinds [26,179,180]. The first mineral in the cosmos was nano-crystalline diamond that condensed from hot, carbon-rich plasma in the cooling atmospheric envelopes of energetic stars [76,181]. Those vapor-deposited diamonds still fall to Earth in presolar grains and as constituents of chondrite meteorites [182]. Those nanodiamonds differ from high-pressure diamonds that form in Earth’s mantle, including “Type I” diamonds in carbon-saturated aqueous fluids and “Type II” diamonds in iron-nickel melts [180,183]. These forms in turn differ from diamonds formed by

impact processes, formerly known as “lonsdaleite” [184,185], and the enigmatic black, porous sintered diamond masses known as carbonado [186,187]. Thus, from a planetary evolution perspective the single mineral species ‘diamond’ is better treated as a suite of distinct natural kinds, each with its unique combination of morphology, trace elements, and other properties. Similar arguments can be made for numerous common minerals, including pyrite (FeS<sub>2</sub>), quartz (SiO<sub>2</sub>), calcite (CaCO<sub>3</sub>), and hydroxylapatite [Ca<sub>5</sub>(PO<sub>4</sub>)<sub>3</sub>OH].

This situation has relevance to the enumeration of Hadean minerals. We need to be concerned not simply with the number of IMA mineral species, but also the number of ‘natural kinds’ of minerals. In terms of life’s origins, different kinds of pyrite, quartz, calcite, hydroxylapatite, and clay may have played very different chemical roles. The enumeration of mineral natural kinds is in its infancy, but these efforts will inform and modify future descriptions of Hadean mineralogy.

#### 4.4.2. Mineral Kinds That Have Been Split

The convention that official mineral species must correspond to endmember chemical compositions has led to splitting of some natural kinds into multiple categories. A dramatic example is provided by the tourmaline group of minerals, represented by the general formula  $XY_3Z_6(T_6O_{18})V_3W$ , where sites *X*, *Y*, *Z*, *T*, *V*, and *W* each can hold multiple different elements ([188,189]; ruff.info/ima). As a result of natural compositional variations, this classification leads to more than 30 IMA approved species in the tourmaline group, with several other potential species pending. Given chemical zonation in individual mineral grains, it is not uncommon for a single crystal to hold two or more different tourmaline species (e.g., [188,190]), even if all tourmaline grains were formed in the same petrogenetic event. On the other hand, it is also possible for individual mineral grains to represent two different natural kinds—for example, a tourmaline grain with an igneous core and hydrothermally-deposited rim. A division of tourmaline into natural kinds might thus provide a more parsimonious description of the supergroup and would facilitate a more accurate understanding of boron mineral diversity and distribution [189].

Similar arguments can be made for many mineral groups, including the apatite group [191,192], the garnet supergroup [193], the amphibole group [194], the mica group [195], the oxide spinel group [196,197], and pyrite [198] and other iron sulfides, all of which were likely present in the Hadean Eon.

#### 4.4.3. Non-Crystalline Materials

The IMA convention on mineral classification considers only crystalline materials. This protocol precludes significant volumes of condensed matter in near-surface environments of Earth, Mars, and other terrestrial worlds. Thus, for example, volcanic glass (e.g., ‘obsidian’), significant components of coal, composite materials, organic mixtures, and common constituents of soil are not easily classified under present mineral classification protocols [199].

This situation is especially significant in the context of Mars. Analyses of soils by NASA’s Mars Science Laboratory reveal more than 50 weight percent amorphous materials in some soil samples [200]. If Mars today mimics aspects of Earth’s Hadean mineralogy (e.g., [61]), we may be ignoring a significant component of early Earth’s near-surface environment. We conclude that expansion of mineral classification to include materials lacking a strictly periodic atomic structure will facilitate the characterization of the actual materials that make up planets.

## 5. Conclusions: Minerals and Origins-of-Life Chemistry

An important motivation for investigating the diversity and distribution of Hadean minerals is the presumed role of specific minerals in life’s geochemical origins. Recent experiments, notably by Steven Benner and colleagues, have documented significant plausible prebiotic catalytic roles for minerals of relatively rare elements, such as boron, molybdenum, and phosphorus [11,15,16]. However, it is not obvious that such minerals as powellite, colemanite, or lüneburgite would have formed during Earth’s

first 500 million years. Even if these types of minerals were present at a few widely dispersed localities, the total volumes of these phases would have been trivial in the global context. As a consequence, a number of elegant chemical solutions to key steps in life's origins have been challenged by the implausibility of invoking rare mineral species (e.g., [1,201]).

A solution to this impasse is to be found in the ubiquity of trace and minor elements in common rock-forming minerals. For example, the average crustal abundance of boron is ~10 ppm [202], with concentrations in the feldspar group of aluminosilicates reported as high as 5000 ppm as boron substitutes for aluminum [203]. Feldspar group minerals are the most abundant phases in Earth's crust, accounting for as much as 60% of the volume of near-surface rocks [202]. As a consequence, immense numbers of borate-like surface sites are available to promote chemical reactions, even if no borate minerals *per se* are present.

Similar arguments can be presented for potentially reactive molybdate, phosphate, or any other surface sites, including sites where two or more atoms are juxtaposed in specific geometries. Indeed, for most relatively rare chemical elements the vast majority of atoms are to be found as trace or minor elements in common minerals, rather than in minerals with those elements as essential constituents. For example, Hazen et al. [204] calculate that all known reserves of cobalt minerals in major ore deposits represent less than one part in 50,000,000 of the cobalt atoms present at ~50 ppm in the vast expanses of basalt that pave the ocean floor [205].

We conclude that, in spite of the relative mineralogical parsimony of Earth's first 500 million years, virtually any type of chemically reactive surface site, containing any of the more than 70 essential mineral-forming elements, would have been widely available throughout the Hadean Eon.

**Author Contributions:** S.M.M. provided discussion on martian and amorphous materials, organic minerals, natural kind clustering, and photo-oxidation. S.E.R. provided discussion on hydrothermal systems and redox gradients. R.M.H. provided information on organic minerals and crystals, impact mineralization, redox gradients, natural kind clustering, and amorphous materials.

**Funding:** This publication is a contribution to the Deep Carbon Observatory. Studies of mineral evolution and mineral ecology are supported by the Deep Carbon Observatory, the Alfred P. Sloan Foundation, the John Templeton Foundation, the NASA Astrobiology Institute, a private foundation, and the Carnegie Institution for Science.

**Acknowledgments:** We are grateful to Steven Benner, Elisa Biondi, Robert Downs, Paul Falkowski, Chao Liu, Steve Mojzsis, Vikas Nanda, Matthew Pasek, Craig Schiffrin, and Nathan Yee for helpful discussions and suggestions.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Hazen, R.M. Paleomineralogy of the Hadean eon: A preliminary species list. *Am. J. Sci.* **2013**, *313*, 807–843. [[CrossRef](#)]
2. Bernal, J.D. *The Physical Basis of Life*; Routledge and Kegan Paul: Abingdon, UK, 1951; ISBN 594837739.
3. Goldschmidt, V.M. Geochemical aspects of the origin of complex organic molecules on the Earth, as precursors to organic life. *New Biol.* **1952**, *12*, 97–105.
4. Cairns-Smith, A.G. *Genetic Takeover and the Mineral Origins of Life*; Cambridge University Press: Cambridge, UK, 1982; ISBN 978-0521346825.
5. Acevedo, O.L.; Orgel, L.E. Template-directed oligonucleotide ligation on hydroxylapatite. *Nature* **1986**, *321*, 790–792. [[CrossRef](#)] [[PubMed](#)]
6. Wächtershäuser, G. Before enzymes and templates: Theory of surface metabolism. *Microbiol. Rev.* **1988**, *52*, 452–484. [[PubMed](#)]
7. Lahav, N. Minerals and the origin of life-hypotheses and experiments in heterogeneous chemistry. *Heterog. Chem. Rev.* **1994**, *1*, 159–179.
8. Ferris, J.P.; Hill, A.R., Jr.; Liu, R.; Orgel, L.E. Synthesis of long prebiotic oligomers on mineral surfaces. *Nature* **1996**, *381*, 59–61. [[CrossRef](#)] [[PubMed](#)]

9. Orgel, L.E. Polymerization on the rocks: Theoretical introduction. *Orig. Life Evol. Biosph.* **1998**, *28*, 227–234. [[CrossRef](#)]
10. Cody, G.D. Transition metal sulfides and the origins of metabolism. *Annu. Rev. Earth Planet. Sci.* **2004**, *32*, 569–599. [[CrossRef](#)]
11. Ricardo, A.; Carrigan, M.A.; Olcott, A.N.; Benner, S.A. Borate minerals stabilize ribose. *Science* **2004**, *303*, 196. [[CrossRef](#)]
12. Hazen, R.M. Presidential Address to the Mineralogical Society of America, Salt Lake City, October 18, 2005: Mineral surfaces and the prebiotic selection and organization of biomolecules. *Am. Mineral.* **2006**, *91*, 1715–1729. [[CrossRef](#)]
13. Pasek, M.A.; Dworkin, J.P.; Lauretta, D.S. A radical pathway for organic phosphorylation during schreibersite corrosion with implications for the origin of life. *Geochim. Cosmochim. Acta* **2007**, *71*, 1721–1736. [[CrossRef](#)]
14. Cleaves, H.J., II; Scott, A.M.; Hill, F.C.; Leszczynski, J.; Sahai, N.; Hazen, R. Mineral–organic interfacial processes: Potential roles in the origins of life. *Chem. Soc. Rev.* **2012**, *41*, 5502–5525. [[CrossRef](#)] [[PubMed](#)]
15. Kim, H.; Furukawa, Y.; Kakegawa, T.; Bitai, A.; Scorei, R.; Benner, S.A. Evaporite Borate-Containing Mineral Ensembles Make Phosphate Available and Regiospecifically Phosphorylate Ribonucleosides: Borate as a Multifaceted Problem Solver in Prebiotic Chemistry. *Angew. Chem. Int. Ed.* **2016**, *55*, 15816–15820. [[CrossRef](#)] [[PubMed](#)]
16. Ziegler, E.W.; Kim, H.-J.; Benner, S.A. Molybdenum (VI)-Catalyzed Rearrangement of Prebiotic Carbohydrates in Formamide, a Candidate Prebiotic Solvent. *Astrobiology* **2018**, *18*, 1159–1170. [[CrossRef](#)] [[PubMed](#)]
17. Schoonen, M.; Smirnov, A. Staging life in an early warm ‘seltzer’ ocean. *Elements* **2016**, *12*, 395–400. [[CrossRef](#)]
18. Lahav, N. *Biogenesis: Theories of Life’s Origin*; Oxford University Press on Demand: Oxford, UK, 1999; ISBN 0195117557.
19. Hazen, R.M. *Genesis: The Scientific Quest for Life’s Origin*; National Academies Press: Washington, DC, USA, 2005; ISBN 0309094321.
20. Sahai, N.; Kaddour, H.; Dalai, P.; Wang, Z.; Bass, G.; Gao, M. Mineral Surface Chemistry and Nanoparticle-aggregation Control Membrane Self-Assembly. *Sci. Rep.* **2017**, *7*, 43418. [[CrossRef](#)] [[PubMed](#)]
21. Hanczyc, M.M.; Fujikawa, S.M.; Szostak, J.W. Experimental Models of Primitive Cellular Compartments: Encapsulation, Growth, and Division. *Science* **2003**, *302*, 618–622. [[CrossRef](#)] [[PubMed](#)]
22. Maynard-Casely, H.E.; Cable, M.L.; Malaska, M.J.; Vu, T.H.; Choukroun, M.; Hodyss, R. Prospects for mineralogy on Titan. *Am. Mineral.* **2018**, *103*, 343–349. [[CrossRef](#)]
23. Hazen, R.M. Titan mineralogy: A window on organic mineral evolution. *Am. Mineral.* **2018**, *103*, 341–342. [[CrossRef](#)]
24. Glikson, A. Eugene Shoemaker and the impact paradigm in Earth and planetary science. *Celest. Mech. Dyn. Astron.* **1997**, *69*, 1–7. [[CrossRef](#)]
25. Pirajno, F. *Hydrothermal Processes and Mineral Systems*; Springer: Amsterdam, The Netherlands, 2009; ISBN 978-1-4020-8613-7.
26. Hazen, R.M. An evolutionary system of mineralogy: Proposal for a classification based on natural kind clustering. *Am. Mineral.* **2019**, in press. [[CrossRef](#)]
27. Reeder, R.J.; Barber, D.J. *Carbonates: Mineralogy and Chemistry*; Mineralogical Society of America: Washington, DC, USA, 1983; Volume 11, ISBN 978-0939950157.
28. Falkowski, P.; Scholes, R.J.; Boyle, E.E.A.; Canadell, J.; Canfield, D.; Elser, J.; Gruber, N.; Hibbard, K.; Höglberg, P.; Linder, S. The global carbon cycle: A test of our knowledge of earth as a system. *Science* **2000**, *290*, 291–296. [[CrossRef](#)] [[PubMed](#)]
29. Hirschmann, M.M.; Dasgupta, R. The H/C ratios of Earth’s near-surface and deep reservoirs, and consequences for deep Earth volatile cycles. *Chem. Geol.* **2009**, *262*, 4–16. [[CrossRef](#)]
30. Perry, R.S.; Mcloughlin, N.; Lynne, B.Y.; Sephton, M.A.; Oliver, J.D.; Perry, C.C.; Campbell, K.; Engel, M.H.; Farmer, J.D.; Brasier, M.D. Defining biominerals and organominerals: Direct and indirect indicators of life. *Sediment. Geol.* **2007**, *201*, 157–179. [[CrossRef](#)]
31. Hazen, R.M.; Downs, R.T.; Jones, A.P.; Kah, L. Carbon Mineralogy and Crystal Chemistry. In *Reviews in Mineralogy and Geochemistry*; Hazen, R.M., Jones, A., Baross, J., Eds.; Mineralogical Society of America: Chantilly, VA, USA, 2013; Volume 75, pp. 7–46. ISBN 1529-6466r978-0-939950-90-4.

32. Nasdala, L.; Pekov, I.V. Ravatite, C<sub>14</sub>HIO, a new organic mineral species from Ravat, Tadzhikistan. *Eur. J. Miner.* **1993**, *5*, 699–705. [[CrossRef](#)]
33. Hummer, D.R.; Noll, B.C.; Hazen, R.M.; Downs, R.T. Crystal structure of abelsonite, the only known crystalline geoporphyrin. *Am. Mineral.* **2017**, *102*, 1129–1132.
34. Oftedal, I. Minerals from the burning Coal Seam at Mt. Pyramide, Spitsbergen. *Result. av Nor. Statsunderstottede Spitsbergenekspeditioner* **1922**, *1*, 9–14.
35. Rost, R. Supplements to the mineralogy of the burning (coal) heaps in the region of Kladno. *Rozpravy II. Tridy Ces. Akad.* **1942**, *52*, 1–4.
36. Belakovskiy, D.I. A natural mineralization process resulting from in situ self-ignition and burning of coal. In Proceedings of the 17th General Meeting of the International Mineralogical Association, Toronto, ON, Canada, 9–14 August 1998.
37. Chesnokov, B.V.; Shcherbakova, Y.P.; Nishanbayev, T.P. Minerals from Burned Dumps of Chelyabinsk Coal Basin. *Miass. Russ. Acad. Sci. Ural. Div. Inst. Mineral.* **2008**, 139.
38. Witzke, T.; de Wit, F.; Kolitsch, U.; Blass, G.; Stracher, G.B.; Prakash, A.; Sokol, E.V. Mineralogy of the burning Anna I coal mine dump, Alsdorf, Germany. In *Coal and Peat Fires—A Global Perspective*; Elsevier: Amsterdam, The Netherlands, 2015; Volume 3, pp. 203–240.
39. Pekov, I.V.; Chukanov, N.V.; Belakovskiy, D.I.; Lykova, I.S.; Yapaskurt, V.O.; Zubkova, N.V.; Shcherbakova, E.P.; Britvin, S. Tinnunculite, IMA 2015-021a. *Mineral. Mag.* **2016**, *80*, 199–205.
40. Benner, S.A.; Kim, H.-J.; Kim, M.-J.; Ricardo, A. Planetary organic chemistry and the origins of biomolecules. *Cold Spring Harb. Perspect. Biol.* **2010**, *2*, a003467. [[CrossRef](#)] [[PubMed](#)]
41. Chyba, C.; Sagan, C. Endogenous production, exogenous delivery and impact-shock synthesis of organic molecules: An inventory for the origins of life. *Nature* **1992**, *355*, 125–132. [[CrossRef](#)] [[PubMed](#)]
42. Ehrenfreund, P.; Charnley, S.B. Organic molecules in the interstellar medium, comets, and meteorites: A voyage from dark clouds to the early Earth. *Annu. Rev. Astron. Astrophys.* **2000**, *38*, 427–483. [[CrossRef](#)]
43. Sephton, M.A. Organic compounds in carbonaceous meteorites. *Nat. Prod. Rep.* **2002**, *19*, 292–311. [[CrossRef](#)] [[PubMed](#)]
44. Pizzarello, S.; Cooper, G.W.; Flynn, G.J. The nature and distribution of the organic material in carbonaceous chondrites and interplanetary dust particles. *Meteorit. Early Sol. Syst. II* **2006**, *1*, 625–651.
45. Burton, A.S.; Stern, J.C.; Elsila, J.E.; Glavin, D.P.; Dworkin, J.P. Understanding prebiotic chemistry through the analysis of extraterrestrial amino acids and nucleobases in meteorites. *Chem. Soc. Rev.* **2012**, *41*, 5459–5472. [[CrossRef](#)]
46. Koga, T.; Naraoka, H. A new family of extraterrestrial amino acids in the Murchison meteorite. *Sci. Rep.* **2017**, *7*, 636. [[CrossRef](#)]
47. Glasby, G.P. Abiogenic origin of hydrocarbons: An historical overview. *Resour. Geol.* **2006**, *56*, 83–96. [[CrossRef](#)]
48. Fu, Q.; Lollar, B.S.; Horita, J.; Lacrampe-Couloume, G.; Seyfried, W.E., Jr. Abiotic formation of hydrocarbons under hydrothermal conditions: Constraints from chemical and isotope data. *Geochim. Cosmochim. Acta* **2007**, *71*, 1982–1998. [[CrossRef](#)]
49. Kolesnikov, A.; Kutcherov, V.G.; Goncharov, A.F. Methane-derived hydrocarbons produced under upper-mantle conditions. *Nat. Geosci.* **2009**, *2*, 566–570. [[CrossRef](#)]
50. McCollom, T.M. Laboratory simulations of abiotic hydrocarbon formation in Earth's deep subsurface. *Rev. Mineral. Geochem.* **2013**, *75*, 467–494. [[CrossRef](#)]
51. Sephton, M.A.; Hazen, R.M. On the origins of deep hydrocarbons. *Rev. Mineral. Geochem.* **2013**, *75*, 449–465. [[CrossRef](#)]
52. Huang, F.; Daniel, I.; Cardon, H.; Montagnac, G.; Sverjensky, D.A. Immiscible hydrocarbon fluids in the deep carbon cycle. *Nat. Commun.* **2017**, *8*, 15798. [[CrossRef](#)] [[PubMed](#)]
53. Cooper, G.; Kimmich, N.; Belisle, W.; Sarinana, J.; Brabham, K.; Garrel, L. Carbonaceous meteorites as a source of sugar-related organic compounds for the early Earth. *Nature* **2001**, *414*, 879–883. [[CrossRef](#)] [[PubMed](#)]
54. Cooper, G.; Rios, A.C. Enantiomer excesses of rare and common sugar derivatives in carbonaceous meteorites. *Proc. Natl. Acad. Sci. USA* **2016**, *113*, E3322–E3331. [[CrossRef](#)] [[PubMed](#)]
55. Springsteen, G.; Joyce, G.F. Selective derivatization and sequestration of ribose from a prebiotic mix. *J. Am. Chem. Soc.* **2004**, *126*, 9578–9583. [[CrossRef](#)] [[PubMed](#)]

56. Hein, J.E.; Tse, E.; Blackmond, D.G. A route to enantiopure RNA precursors from nearly racemic starting materials. *Nat. Chem.* **2011**, *3*, 704–706. [[CrossRef](#)]
57. Blackmond, D.G. Asymmetric autocatalysis and its implications for the origin of homochirality. *Proc. Natl. Acad. Sci.* **2004**, *101*, 5732–5736. [[CrossRef](#)]
58. Blackmond, D.G. The origin of biological homochirality. *Cold Spring Harb. Perspect. Biol.* **2010**, *2*, a002147. [[CrossRef](#)]
59. Kvenvolden, K.A. A review of the geochemistry of methane in natural gas hydrate. *Org. Geochem.* **1995**, *23*, 997–1008. [[CrossRef](#)]
60. Buffett, B.A. Clathrate hydrates. *Annu. Rev. Earth Planet. Sci.* **2000**, *28*, 477–507. [[CrossRef](#)]
61. Hazen, R.M.; Sverjensky, D.A.; Azzolini, D.; Bish, D.L.; Elmore, S.C.; Hinnov, L.; Milliken, R.E. Clay mineral evolution. *Am. Mineral.* **2013**, *98*, 2007–2029. [[CrossRef](#)]
62. Guggenheim, S.; van Groos, A.F.K. New gas-hydrate phase: Synthesis and stability of clay–methane hydrate intercalate. *Geology* **2003**, *31*, 653–656. [[CrossRef](#)]
63. Benner, S.A.; Devine, K.G.; Matveeva, L.N.; Powell, D.H. The missing organic molecules on Mars. *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 2425–2430. [[CrossRef](#)] [[PubMed](#)]
64. Ritson, D.J.; Sutherland, J.D. Synthesis of aldehydic ribonucleotide and amino acid precursors by photoredox chemistry. *Angew. Chem. Int. Ed.* **2013**, *52*, 5845–5847. [[CrossRef](#)]
65. Powner, M.W.; Gerland, B.; Sutherland, J.D. Synthesis of activated pyrimidine ribonucleotides in prebiotically plausible conditions. *Nature* **2009**, *459*, 239–242. [[CrossRef](#)] [[PubMed](#)]
66. Szostak, J.W. The origin of life on Earth and the design of alternative life forms. *Mol. Front. J.* **2017**, *1*, 121–131. [[CrossRef](#)]
67. Kebukawa, Y.; Cody, G.D. A kinetic study of the formation of organic solids from formaldehyde: Implications for the origin of extraterrestrial organic solids in primitive Solar System objects. *Icarus* **2015**, *248*, 412–423. [[CrossRef](#)]
68. Glikson, A.Y.; Pirajno, F. *Asteroids Impacts, Crustal Evolution and Related Mineral Systems with Special Reference to Australia*; Springer: Berlin, Germany, 2018; ISBN 331974545X.
69. Versh, E.; Kirsimäe, K.; Jõelet, A. Development of potential ecological niches in impact-induced hydrothermal systems: The small-to-medium size impacts. *Planet. Space Sci.* **2006**, *54*, 1567–1574. [[CrossRef](#)]
70. Abramov, O.; Kring, D.A. Impact-induced hydrothermal activity on early Mars. *J. Geophys. Res. Planets* **2005**, *110*. [[CrossRef](#)]
71. Allen, C.C.; Gooding, J.L.; Keil, K. Hydrothermally altered impact melt rock and breccia: Contributions to the soil of Mars. *J. Geophys. Res. Solid Earth* **1982**, *87*, 10083–10101. [[CrossRef](#)]
72. Pirajno, F. Hydrothermal processes associated with meteorite impact structures: Evidence from three Australian examples and implications for economic resources. *Aust. J. Earth Sci.* **2005**, *52*, 587–605. [[CrossRef](#)]
73. Naldrett, A.J. From impact to riches: Evolution of geological understanding. *GSA TODAY* **2003**, *5*, 4–9. [[CrossRef](#)]
74. Johansson, Å. Geochemical studies on the Boda Pb-Zn deposit in the Siljan astrobleme, central Sweden. *Geol. Föreningen i Stock. Förhandlingar* **1984**, *106*, 15–25. [[CrossRef](#)]
75. Anthony, J.W.; Bideaux, R.A.; Bladh, K.W.; Nichols, M.C. Borates, Carbonates, Sulfates. In *Handbook of Mineralogy*; Mineralogical Society of America: Chantilly, VA, USA, 2003; Volume V, ISBN 0-9622097-0-8.
76. Hazen, R.M.; Papineau, D.; Bleeker, W.; Downs, R.T.; Ferry, J.M.; McCoy, T.J.; Sverjensky, D.A.; Yang, H. Mineral evolution. *Am. Mineral.* **2008**, *93*, 1693–1720. [[CrossRef](#)]
77. Delano, J.W. Redox history of the Earth's interior since ~3900 Ma: Implications for prebiotic molecules. *Orig. Life Evol. Biosph.* **2001**, *31*, 311–341. [[CrossRef](#)]
78. Canil, D. Vanadium in peridotites, mantle redox and tectonic environments: Archean to present. *Earth Planet. Sci. Lett.* **2002**, *195*, 75–90. [[CrossRef](#)]
79. Li, Z.-X.A.; Lee, C.-T.A. The constancy of upper mantle fO<sub>2</sub> through time inferred from V/Sc ratios in basalts. *Earth Planet. Sci. Lett.* **2004**, *228*, 483–493.
80. Frost, D.J.; McCammon, C.A. The redox state of Earth's mantle. *Annu. Rev. Earth Planet. Sci.* **2008**, *36*, 389–420. [[CrossRef](#)]
81. Trail, D.; Watson, E.B.; Tailby, N.D. The oxidation state of Hadean magmas and implications for early Earth's atmosphere. *Nature* **2011**, *480*, 79–82. [[CrossRef](#)]

82. Righter, K.; Ghiorso, M.S. Redox systematics of a magma ocean with variable pressure-temperature gradients and composition. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 11955–11960. [[CrossRef](#)] [[PubMed](#)]
83. Armstrong, K.; Frost, D.J.; McCammon, C.M.; Rubie, D.; Boffa-Ballaran, T. Oxidation States of Fe in Silicate Melts as a Function of Pressure and Implications for Redox Evolution of the Early Mantle. *Proc. Lunar Planet. Sci. Conf.* **2016**, *47*, 2580.
84. Zhang, H.L.; Hirschmann, M.M.; Cottrell, E.; Withers, A.C. Effect of pressure on Fe<sup>3+</sup>/ΣFe ratio in a mafic magma and consequences for magma ocean redox gradients. *Geochim. Cosmochim. Acta* **2017**, *204*, 83–103. [[CrossRef](#)]
85. Mahaffy, P.R.; Webster, C.R.; Atreya, S.K.; Franz, H.; Wong, M.; Conrad, P.G.; Harpold, D.; Jones, J.J.; Leshin, L.A.; Manning, H. Abundance and isotopic composition of gases in the Martian atmosphere from the Curiosity rover. *Science* **2013**, *341*, 263–266. [[CrossRef](#)]
86. Nair, H.; Allen, M.; Anbar, A.D.; Yung, Y.L.; Clancy, R.T. A photochemical model of the Martian atmosphere. *Icarus* **1994**, *111*, 124–150. [[CrossRef](#)] [[PubMed](#)]
87. Sakuraba, H.; Kurokawa, H.; Genda, H. Impact degassing and atmospheric erosion on Venus, Earth, and Mars during the late accretion. *Icarus* **2019**, *317*, 48–58. [[CrossRef](#)]
88. Lyons, T.W.; Reinhard, C.T.; Planavsky, N.J. The rise of oxygen in Earth's early ocean and atmosphere. *Nature* **2014**, *506*, 307–315. [[CrossRef](#)]
89. Canfield, D.E. *Oxygen: A Four Billion Year History*; Princeton University Press: Princeton, NJ, USA, 2014; ISBN 1400849888.
90. Genda, H.; Brasser, R.; Mojzsis, S.J. The terrestrial late veneer from core disruption of a lunar-sized impactor. *Earth Planet. Sci. Lett.* **2017**, *480*, 25–32. [[CrossRef](#)]
91. Genda, H.; Iizuka, T.; Sasaki, T.; Ueno, Y.; Ikoma, M. Ejection of iron-bearing giant-impact fragments and the dynamical and geochemical influence of the fragment re-accretion. *Earth Planet. Sci. Lett.* **2017**, *470*, 87–95. [[CrossRef](#)]
92. Hazen, R.M.; Ewing, R.C.; Sverjensky, D.A. Evolution of uranium and thorium minerals. *Am. Mineral.* **2009**, *94*, 1293–1311. [[CrossRef](#)]
93. Hao, J.; Sverjensky, D.A.; Hazen, R.M. A model for late Archean chemical weathering and world average river water. *Earth Planet. Sci. Lett.* **2017**, *457*, 191–203. [[CrossRef](#)]
94. Roedder, E. Silicate liquid immiscibility in magmas and in the system K<sub>2</sub>O-FeO-Al<sub>2</sub>O<sub>3</sub>-SiO<sub>2</sub>: An example of serendipity. *Geochim. Cosmochim. Acta* **1978**, *42*, 1597–1617. [[CrossRef](#)]
95. Philpotts, A.R. Compositions of immiscible liquids in volcanic rocks. *Contrib. Mineral. Petrol.* **1982**, *80*, 201–218. [[CrossRef](#)]
96. Charlier, B.; Grove, T.L. Experiments on liquid immiscibility along tholeiitic liquid lines of descent. *Contrib. Mineral. Petrol.* **2012**, *164*, 27–44. [[CrossRef](#)]
97. Wager, L.R.; Brown, G.M. *Layered Igneous Rocks*; W. H. Freeman and Company: New York, NY, USA, 1968; ISBN 978-0716702368.
98. Gray, N.H.; Crain, I.K. Crystal settling in sills: A model for suspension settling. *Can. J. Earth Sci.* **1969**, *6*, 1211–1216. [[CrossRef](#)]
99. Fujii, T. Crystal settling in a sill. *Lithos* **1974**, *7*, 133–137. [[CrossRef](#)]
100. Martin, D.; Nokes, R. Crystal settling in a vigorously converting magma chamber. *Nature* **1988**, *332*, 534–536. [[CrossRef](#)]
101. Weinstein, S.A.; Yuen, D.A.; Olson, P.L. Evolution of crystal-settling in magma-chamber convection. *Earth Planet. Sci. Lett.* **1988**, *87*, 237–248. [[CrossRef](#)]
102. Hoshida, T.; Obata, M.; Akatsuka, T. Crystal settling and crystal growth of olivine in magmatic differentiation—The Murotomisaki Gabbroic Complex, Shikoku, Japan. *J. Mineral. Petrol. Sci.* **2006**, *101*, 223–239. [[CrossRef](#)]
103. Sun, W.; Arculus, R.J.; Kamenetsky, V.S.; Binns, R.A. Release of gold-bearing fluids in convergent margin magmas prompted by magnetite crystallization. *Nature* **2004**, *431*, 975–978. [[CrossRef](#)]
104. Maier, W.D. Platinum-group element (PGE) deposits and occurrences: Mineralization styles, genetic concepts, and exploration criteria. *J. Afr. Earth Sci.* **2005**, *41*, 165–191. [[CrossRef](#)]
105. Collins, S.J.; Maclennan, J.; Pyle, D.M.; Barnes, S.-J.; Upton, B.G.J. Two phases of sulphide saturation in Réunion magmas: Evidence from cumulates. *Earth Planet. Sci. Lett.* **2012**, *337*, 104–113. [[CrossRef](#)]

106. Savelyev, D.P.; Kamenetsky, V.S.; Danyushevsky, L.V.; Botcharnikov, R.E.; Kamenetsky, M.B.; Park, J.-W.; Portnyagin, M.V.; Olin, P.; Krashenninikov, S.P.; Hauff, F. Immiscible sulfide melts in primitive oceanic magmas: Evidence and implications from picrite lavas (Eastern Kamchatka, Russia). *Am. Mineral.* **2018**, *103*, 886–898. [[CrossRef](#)]
107. Lester, G.W.; Clark, A.H.; Kyser, T.K.; Naslund, H.R. Experiments on liquid immiscibility in silicate melts with H<sub>2</sub>O, P, S, F and Cl: Implications for natural magmas. *Contrib. Mineral. Petrol.* **2013**, *166*, 329–349. [[CrossRef](#)]
108. Roedder, E. Fluid inclusion evidence for immiscibility in magmatic differentiation. *Geochim. Cosmochim. Acta* **1992**, *56*, 5–20. [[CrossRef](#)]
109. Veksler, I.V.; Dorfman, A.M.; Dulski, P.; Kamenetsky, V.S.; Danyushevsky, L.V.; Jeffries, T.; Dingwell, D.B. Partitioning of elements between silicate melt and immiscible fluoride, chloride, carbonate, phosphate and sulfate melts, with implications to the origin of natrocarbonatite. *Geochim. Cosmochim. Acta* **2012**, *79*, 20–40. [[CrossRef](#)]
110. Naldrett, A.J. *Magmatic Sulfide Deposits: Geology, Geochemistry and Exploration*; Springer Science & Business Media: Berlin, Germany, 2004; ISBN 3662084449.
111. Naldrett, A.J. Ores associated with flood basalts. In *Reviews in Economic Geology: Ore Deposition Associated with Magmas*; Whitney, J.A., Naldrett, A.J., Eds.; Society of Economic Geologists: Littleton, CO, USA, 1989; Volume 4, pp. 103–118. ISBN 0961307404.
112. Naldrett, A.J. A model for the Ni-Cu-PGE ores of the Noril'sk region and its application to other areas of flood basalt. *Econ. Geol.* **1992**, *87*, 1945–1962. [[CrossRef](#)]
113. Naldrett, A.J. World-class Ni-Cu-PGE deposits: Key factors in their genesis. *Miner. Depos.* **1999**, *34*, 227–240. [[CrossRef](#)]
114. Lightfoot, P.C.; Hawkesworth, C.J. Flood basalts and magmatic Ni, Cu, and PGE sulphide mineralization: Comparative geochemistry of the Noril'sk (Siberian Traps) and West Greenland sequences. *Geophys. Monogr. Geophys. Union* **1997**, *100*, 357–380.
115. Leshner, C.M.; Keays, R.R. Komatiite-associated Ni-Cu-PGE deposits. *Can. Inst. Min. Metall. Pet. Spec.* **2002**, *54*, 579–617.
116. Czamanske, G.K.; Zen'ko, T.E.; Fedorenko, V.A.; Calk, L.C.; Budahn, J.R.; Bullock, J.H.; Fries, T.L.; King, B.-S.W.; Siems, D.F. Petrographic and Geochemical Characterization of Ore-bearing Intrusions of the Noril'sk type, Siberia; With Discussion of Their Origin. *Resour. Geol. Spec. Issue* **1995**, *18*, 1–48.
117. Kjarsgaard, B.A.; Hamilton, D.L. The immiscibility of silicate and carbonate liquids. *S. Afr. J. Geol.* **1993**, *96*, 139–142.
118. Solovova, I.P.; Girmis, A.V.; Kogarko, L.N.; Kononkova, N.N.; Stoppa, F.; Rosatelli, G. Compositions of magmas and carbonate–silicate liquid immiscibility in the Vulture alkaline igneous complex, Italy. *Lithos* **2005**, *85*, 113–128. [[CrossRef](#)]
119. Guzmics, T.; Mitchell, R.H.; Szabó, C.; Berkesi, M.; Milke, R.; Ratter, K. Liquid immiscibility between silicate, carbonate and sulfide melts in melt inclusions hosted in co-precipitated minerals from Kerimasi volcano (Tanzania): Evolution of carbonated nephelinitic magma. *Contrib. Mineral. Petrol.* **2012**, *164*, 101–122. [[CrossRef](#)]
120. Guzmics, T.; Zajacz, Z.; Mitchell, R.H.; Szabó, C.; Wälle, M. The role of liquid–liquid immiscibility and crystal fractionation in the genesis of carbonatite magmas: Insights from Kerimasi melt inclusions. *Contrib. Mineral. Petrol.* **2015**, *169*, 17. [[CrossRef](#)]
121. Lee, W.; Wyllie, P.J. Liquid immiscibility between nephelinite and carbonatite from 1.0 to 2.5 GPa compared with mantle melt compositions. *Contrib. Mineral. Petrol.* **1997**, *127*, 1–16. [[CrossRef](#)]
122. Brooker, R.A. The effect of CO<sub>2</sub> saturation on immiscibility between silicate and carbonate liquids: An experimental study. *J. Petrol.* **1998**, *39*, 1905–1915. [[CrossRef](#)]
123. Veksler, I.V.; Dorfman, A.M.; Danyushevsky, L.V.; Jakobsen, J.K.; Dingwell, D.B. Immiscible silicate liquid partition coefficients: Implications for crystal–melt element partitioning and basalt petrogenesis. *Contrib. Mineral. Petrol.* **2006**, *152*, 685–702. [[CrossRef](#)]
124. Thomsen, T.B.; Schmidt, M.W. Melting of carbonated pelites at 2.5–5.0 GPa, silicate–carbonatite liquid immiscibility, and potassium–carbon metasomatism of the mantle. *Earth Planet. Sci. Lett.* **2008**, *267*, 17–31. [[CrossRef](#)]

125. Brooker, R.A.; Kjarsgaard, B.A. Silicate–carbonate liquid immiscibility and phase relations in the system  $\text{SiO}_2\text{--Na}_2\text{O--Al}_2\text{O}_3\text{--CaO--CO}_2$  at 1–2.5 GPa with applications to carbonatite genesis. *J. Petrol.* **2010**, *52*, 1281–1305. [[CrossRef](#)]
126. Martin, L.H.J.; Schmidt, M.W.; Mattsson, H.B.; Guenther, D. Element partitioning between immiscible carbonatite and silicate melts for dry and  $\text{H}_2\text{O}$ -bearing systems at 1–3 GPa. *J. Petrol.* **2013**, *54*, 2301–2338. [[CrossRef](#)]
127. Lister, G.F. The composition and origin of selected iron-titanium deposits. *Econ. Geol.* **1966**, *61*, 275–310. [[CrossRef](#)]
128. Philpotts, A.R. Origin of certain iron-titanium oxide and apatite rocks. *Econ. Geol.* **1967**, *62*, 303–315. [[CrossRef](#)]
129. Namur, O.; Charlier, B.; Holness, M.B. Dual origin of Fe–Ti–P gabbros by immiscibility and fractional crystallization of evolved tholeiitic basalts in the Sept Iles layered intrusion. *Lithos* **2012**, *154*, 100–114. [[CrossRef](#)]
130. Chen, W.T.; Zhou, M.-F.; Zhao, T.-P. Differentiation of nelsonitic magmas in the formation of the ~ 1.74 Ga Damiao Fe–Ti–P ore deposit, North China. *Contrib. Mineral. Petrol.* **2013**, *165*, 1341–1362. [[CrossRef](#)]
131. Lindsley, D.H.; Epler, N. Do Fe-Ti-oxide magmas exist? Probably not! *Am. Mineral.* **2017**, *102*, 2157–2169. [[CrossRef](#)]
132. Charlier, B.; Namur, O.; Bolle, O.; Latypov, R.; Duchesne, J.-C. Fe–Ti–V–P ore deposits associated with Proterozoic massif-type anorthosites and related rocks. *Earth-Sci. Rev.* **2015**, *141*, 56–81. [[CrossRef](#)]
133. Oen, I.S. A peculiar type of Cr-Ni-mineralization; cordierite-chromite-niccolite ores of Malaga, Spain, and their possible origin by liquid unmixing. *Econ. Geol.* **1973**, *68*, 831–842. [[CrossRef](#)]
134. Makovicky, E.; Karup-Møller, S.; Makovicky, M.; Rose-Hansen, J. Experimental studies on the phase systems Fe-Ni-Pd-S and Fe-Pt-Pd-As-S applied to PGE deposits. *Mineral. Petrol.* **1990**, *42*, 307–319. [[CrossRef](#)]
135. Fleet, M.E.; Chryssoulis, S.L.; Stone, W.E.; Weisener, C.G. Partitioning of platinum-group elements and Au in the Fe–Ni–Cu–S system: Experiments on the fractional crystallization of sulfide melt. *Contrib. Mineral. Petrol.* **1993**, *115*, 36–44. [[CrossRef](#)]
136. Gervilla, F.; Leblanc, M.; Torres-Ruiz, J.; Fenoll Hach-Ali, P. Immiscibility between arsenide and sulfide melts; a mechanism for concentration of noble metals. *Can. Mineral.* **1996**, *34*, 485–502.
137. Gervilla, F.; Gutiérrez-Narbona, R.; Fenoll-Hach-Alí, P. The origin of different types of magmatic mineralizations from small-volume melts in the lherzolite massifs of the Serranía de Ronda. *Boletín la Soc. Esp. Mineral.* **2002**, *25*, 79–96.
138. Bali, E.; Audétat, A.; Keppler, H. Water and hydrogen are immiscible in Earth’s mantle. *Nature* **2013**, *495*, 220–222. [[CrossRef](#)] [[PubMed](#)]
139. Ruprecht, P.; Fiege, A.; Simon, A. Local redox gradients induced during magma mixing. In Proceedings of the XIV Congreso Geológico Chileno, La Serena, Chile, 4–8 October 2015; pp. 643–646.
140. Fiege, A.; Ruprecht, P.; Simon, A. A magma mixing redox trap that moderates mass transfer of sulphur and metals. *GEOCHEMICAL Perspect. Lett.* **2017**, *3*, 190–198. [[CrossRef](#)]
141. Wilde, S.A.; Valley, J.W.; Peck, W.H.; Graham, C.M. Evidence from detrital zircons for the existence of continental crust and oceans on the Earth 4.4 Gyr ago. *Nature* **2001**, *409*, 175–178. [[CrossRef](#)] [[PubMed](#)]
142. Harrison, T.M.; Schmitt, A.K.; McCulloch, M.T.; Lovera, O.M. Early ( $\geq 4.5$  Ga) formation of terrestrial crust: Lu–Hf,  $\delta^{18}\text{O}$ , and Ti thermometry results for Hadean zircons. *Earth Planet. Sci. Lett.* **2008**, *268*, 476–486. [[CrossRef](#)]
143. Valley, J.W.; Cavosie, A.J.; Ushikubo, T.; Reinhard, D.A.; Lawrence, D.F.; Larson, D.J.; Clifton, P.H.; Kelly, T.F.; Wilde, S.A.; Moser, D.E. Hadean age for a post-magma-ocean zircon confirmed by atom-probe tomography. *Nat. Geosci.* **2014**, *7*, 219–223. [[CrossRef](#)]
144. O’neil, J.; Carlson, R.W. Building Archean cratons from Hadean mafic crust. *Science* **2017**, *355*, 1199–1202. [[CrossRef](#)]
145. Johnson, T.E.; Gardiner, N.J.; Miljković, K.; Spencer, C.J.; Kirkland, C.L.; Bland, P.A.; Smithies, H. An impact melt origin for Earth’s oldest known evolved rocks. *Nat. Geosci.* **2018**, *11*, 795–799. [[CrossRef](#)]
146. Pasek, M.A.; Lauretta, D.S. Aqueous corrosion of phosphide minerals from iron meteorites: A highly reactive source of prebiotic phosphorus on the surface of the early Earth. *Astrobiology* **2005**, *5*, 515–535. [[CrossRef](#)]
147. Pasek, M.A. Rethinking early Earth phosphorus geochemistry. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 853–858. [[CrossRef](#)] [[PubMed](#)]

148. Gull, M.; Mojica, M.A.; Fernández, F.M.; Gaul, D.A.; Orlando, T.M.; Liotta, C.L.; Pasek, M.A. Nucleoside phosphorylation by the mineral schreibersite. *Sci. Rep.* **2015**, *5*, 17198. [[CrossRef](#)] [[PubMed](#)]
149. Essene, E.J.; Fisher, D.C. Lightning strike fusion: Extreme reduction and metal-silicate liquid immiscibility. *Science* **1986**, *234*, 189–193. [[CrossRef](#)] [[PubMed](#)]
150. Parnell, J.; Thackrey, S.; Muirhead, D.K.; Wright, A.J. Transient high-temperature processing of silicates in fulgurites as analogues for meteorite and impact melts. In Proceedings of the Lunar and Planetary Science Conference, League City, TX, USA, 10–14 March 2008; Volume 39, p. 1286.
151. Pasek, M.A.; Block, K.; Pasek, V. Fulgurite morphology: A classification scheme and clues to formation. *Contrib. Mineral. Petrol.* **2012**, *164*, 477–492. [[CrossRef](#)]
152. Pasek, M.; Block, K. Lightning-induced reduction of phosphorus oxidation state. *Nat. Geosci.* **2009**, *2*, 553–556. [[CrossRef](#)]
153. Glindemann, D.; Edwards, M.; Schrems, O. Phosphine and methylphosphine production by simulated lightning—A study for the volatile phosphorus cycle and cloud formation in the earth atmosphere. *Atmos. Environ.* **2004**, *38*, 6867–6874. [[CrossRef](#)]
154. Daly, T.K.; Buseck, P.R.; Williams, P.; Lewis, C.F. Fullerenes from a fulgurite. *Science* **1993**, *259*, 1599–1601. [[CrossRef](#)] [[PubMed](#)]
155. Wadsworth, F.B.; Vasseur, J.; Llewellyn, E.W.; Genareau, K.; Cimarelli, C.; Dingwell, D.B. Size limits for rounding of volcanic ash particles heated by lightning. *J. Geophys. Res. Solid Earth* **2017**, *122*, 1977–1989. [[CrossRef](#)] [[PubMed](#)]
156. Yamamoto, M.; Nakamura, R.; Kasaya, T.; Kumagai, H.; Suzuki, K.; Takai, K. Spontaneous and Widespread Electricity Generation in Natural Deep-Sea Hydrothermal Fields. *Angew. Chem. Int. Ed.* **2017**, *56*, 5725–5728. [[CrossRef](#)]
157. Ang, R.; Khan, A.U.; Tsujii, N.; Takai, K.; Nakamura, R.; Mori, T. Thermoelectricity Generation and Electron–Magnon Scattering in a Natural Chalcopyrite Mineral from a Deep-Sea Hydrothermal Vent. *Angew. Chem. Int. Ed.* **2015**, *54*, 12909–12913. [[CrossRef](#)]
158. Rona, P.A. The changing vision of marine minerals. *Ore Geol. Rev.* **2008**, *33*, 618–666. [[CrossRef](#)]
159. Lusty, P.A.J.; Murton, B.J. Deep-ocean mineral deposits: Metal resources and windows into Earth processes. *Elem. Int. Mag. Mineral. Geochem. Petrol.* **2018**, *14*, 301–306. [[CrossRef](#)]
160. Petersen, S.; Lehrmann, B.; Murton, B.J. Modern seafloor hydrothermal systems: New perspectives on ancient ore-forming processes. *Elem. Int. Mag. Mineral. Geochem. Petrol.* **2018**, *14*, 307–312. [[CrossRef](#)]
161. Hein, J.R.; Mizell, K.; Koschinsky, A.; Conrad, T.A. Deep-ocean mineral deposits as a source of critical metals for high-and green-technology applications: Comparison with land-based resources. *Ore Geol. Rev.* **2013**, *51*, 1–14. [[CrossRef](#)]
162. Kim, J.D.; Yee, N.; Nanda, V.; Falkowski, P.G. Anoxic photochemical oxidation of siderite generates molecular hydrogen and iron oxides. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 10073–10077. [[CrossRef](#)] [[PubMed](#)]
163. Braterman, P.S.; Cairns-Smith, A.G.; Sloper, R.W. Photo-oxidation of hydrated Fe<sup>2+</sup>—Significance for banded iron formations. *Nature* **1983**, *303*, 163–164. [[CrossRef](#)]
164. Borowska, Z.; Mauzerall, D. Formation of hydrogen on irradiation of aqueous ferrous ion with UV light at neutral pH. *Orig. Life Evol. Biosph.* **1986**, *16*, 194–195. [[CrossRef](#)]
165. Borowska, Z.K.; Mauzerall, D.C. Efficient near ultraviolet light induced formation of hydrogen by ferrous hydroxide. *Orig. Life Evol. Biosph.* **1987**, *17*, 251–259. [[CrossRef](#)]
166. Matthews, R.W. Photooxidation of organic impurities in water using thin films of titanium dioxide. *J. Phys. Chem.* **1987**, *91*, 3328–3333. [[CrossRef](#)]
167. Chen, X.D.; Wang, Z.; Liao, Z.F.; Mai, Y.L.; Zhang, M.Q. Roles of anatase and rutile TiO<sub>2</sub> nanoparticles in photooxidation of polyurethane. *Polym. Test.* **2007**, *26*, 202–208. [[CrossRef](#)]
168. Valdes, A.; Qu, Z.-W.; Kroes, G.-J.; Rossmesl, J.; Nørskov, J.K. Oxidation and photo-oxidation of water on TiO<sub>2</sub> surface. *J. Phys. Chem. C* **2008**, *112*, 9872–9879. [[CrossRef](#)]
169. Valdés, Á.; Kroes, G.-J. Cluster study of the photo-oxidation of water on rutile titanium dioxide (TiO<sub>2</sub>). *J. Phys. Chem. C* **2010**, *114*, 1701–1708. [[CrossRef](#)]
170. Waki, K.; Zhao, J.; Horikoshi, S.; Watanabe, N.; Hidaka, H. Photooxidation mechanism of nitrogen-containing compounds at TiO<sub>2</sub>/H<sub>2</sub>O interfaces: An experimental and theoretical examination of hydrazine derivatives. *Chemosphere* **2000**, *41*, 337–343. [[CrossRef](#)]

171. Kebede, M.A.; Varner, M.E.; Scharko, N.K.; Gerber, R.B.; Raff, J.D. Photooxidation of Ammonia on TiO<sub>2</sub> as a Source of NO and NO<sub>2</sub> under Atmospheric Conditions. *J. Am. Chem. Soc.* **2013**, *135*, 8606–8615. [[CrossRef](#)] [[PubMed](#)]
172. Miller, G.; Kempley, R.; Awadh, G.; Richman, K. Photo-oxidation of chloride to perchlorate in the presence of titanium dioxide and nitrate. In *Abstracts of Papers of the American Chemical Society*; American Chemical Society: Washington, DC, USA, 2004; Volume 228, p. U92.
173. Schuttlefield, J.D.; Sambur, J.B.; Gelwicks, M.; Eggleston, C.M.; Parkinson, B.A. Photooxidation of chloride by oxide minerals: Implications for perchlorate on Mars. *J. Am. Chem. Soc.* **2011**, *133*, 17521–17523. [[CrossRef](#)] [[PubMed](#)]
174. Kim, Y.S.; Wo, K.P.; Maity, S.; Atreya, S.K.; Kaiser, R.I. Radiation-induced formation of chlorine oxides and their potential role in the origin of Martian perchlorates. *J. Am. Chem. Soc.* **2013**, *135*, 4910–4913. [[CrossRef](#)] [[PubMed](#)]
175. Nie, N.X.; Dauphas, N.; Greenwood, R.C. Iron and oxygen isotope fractionation during iron UV photo-oxidation: Implications for early Earth and Mars. *Earth Planet. Sci. Lett.* **2017**, *458*, 179–191. [[CrossRef](#)]
176. Mills, S.J.; Hatert, F.; Nickel, E.H.; Ferraris, G. The standardisation of mineral group hierarchies: Application to recent nomenclature proposals. *Eur. J. Mineral.* **2009**, *21*, 1073–1080. [[CrossRef](#)]
177. Schertl, H.-P.; Mills, S.J.; Maresch, W.V. A Compendium of IMA-Approved Nomenclature. In Proceedings of the XXII General Meeting of the International Mineralogical Association, Melbourne, Australia, 13–17 August 2018.
178. Lafuente, B.; Downs, R.T.; Yang, H.; Stone, N. *The Power of Databases: The RRUFF Project*; Walter de Gruyter GmbH: Berlin, Germany, 2015; ISBN 9783110417043.
179. Davis, G. *Diamond*; Adam Hilger Ltd.: Bristol, UK, 1984.
180. Shirey, S.B.; Cartigny, P.; Frost, D.J.; Keshav, S.; Nestola, F.; Nimis, P.; Pearson, D.G.; Sobolev, N.V.; Walter, M.J. Diamonds and the geology of mantle carbon. *Rev. Mineral. Geochem.* **2013**, *75*, 355–421. [[CrossRef](#)]
181. Greaves, J.S.; Scaife, A.M.M.; Frayer, D.T.; Green, D.A.; Mason, B.S.; Smith, A.M.S. Anomalous microwave emission from spinning nanodiamonds around stars. *Nat. Astron.* **2018**, *2*, 662–667. [[CrossRef](#)]
182. Ott, U. Nanodiamonds in meteorites: Properties and astrophysical context. *J. Achiev. Mater. Manuf. Eng.* **2009**, *37*, 779–784.
183. Smith, E.M.; Shirey, S.B.; Nestola, F.; Bullock, E.S.; Wang, J.; Richardson, S.H.; Wang, W. Large gem diamonds from metallic liquid in Earth's deep mantle. *Science* **2016**, *354*, 1403–1405. [[CrossRef](#)] [[PubMed](#)]
184. Vishnevsky, S.; Raitala, J. Impact diamonds as indicators of shock metamorphism in strongly-reworked Pre-Cambrian impactites. In *Impacts and the Early Earth*; Springer: Berlin, Germany, 2000; pp. 229–247.
185. Németh, P.; Garvie, L.A.J.; Aoki, T.; Dubrovinskaia, N.; Dubrovinsky, L.; Buseck, P.R. Lonsdaleite is faulted and twinned cubic diamond and does not exist as a discrete material. *Nat. Commun.* **2014**, *5*, 5447. [[CrossRef](#)] [[PubMed](#)]
186. Heaney, P.J.; Vicenzi, E.P.; De, S. Strange diamonds: The mysterious origins of carbonado and framesite. *Elements* **2005**, *1*, 85–89. [[CrossRef](#)]
187. Garai, J.; Haggerty, S.E.; Rekhi, S.; Chance, M. Infrared absorption investigations confirm the extraterrestrial origin of carbonado diamonds. *Astrophys. J. Lett.* **2006**, *653*, L153. [[CrossRef](#)]
188. Henry, D.J.; Novák, M.; Hawthorne, F.C.; Ertl, A.; Dutrow, B.L.; Uher, P.; Pezzotta, F. Nomenclature of the tourmaline-supergroup minerals. *Am. Mineral.* **2011**, *96*, 895–913. [[CrossRef](#)]
189. Grew, E.S.; Hystad, G.; Hazen, R.M.; Krivovichev, S.V.; Gorelova, L.A. How many boron minerals occur in Earth's upper crust? *Am. Mineral.* **2017**, *102*, 1573–1587. [[CrossRef](#)]
190. Grew, E.S.; Dymek, R.F.; De Hoog, J.C.M.; Harley, S.L.; Boak, J.; Hazen, R.M.; Yates, M.G. Boron isotopes in tourmaline from the ca. 3.7–3.8 Ga Isua supracrustal belt, Greenland: Sources for boron in Eoarchean continental crust and seawater. *Geochim. Cosmochim. Acta* **2015**, *163*, 156–177. [[CrossRef](#)]
191. Pasero, M.; Kampf, A.R.; Ferraris, C.; Pekov, I.V.; Rakovan, J.; White, T.J. Nomenclature of the apatite supergroup minerals. *Eur. J. Mineral.* **2010**, *22*, 163–179. [[CrossRef](#)]
192. Harlov, D.E.; Rakovan, J.F. Apatite: A mineral for all seasons. *Elements* **2015**, *11*, 165–200.
193. Grew, E.S.; Locock, A.J.; Mills, S.J.; Galuskina, I.O.; Galuskin, E.V.; Hälenius, U. Nomenclature of the garnet supergroup. *Am. Mineral.* **2013**, *98*, 785–811. [[CrossRef](#)]
194. Hawthorne, F.C.; Oberti, R.; Harlow, G.E.; Maresch, W.V.; Martin, R.F.; Schumacher, J.C.; Welch, M.D. Nomenclature of the amphibole supergroup. *Am. Mineral.* **2012**, *97*, 2031–2048. [[CrossRef](#)]

195. Rieder, M.; Cavazzini, G.; D'yakonov, Y.S.; Frank-Kamenetskii, V.A.; Gottardi, G.; Guggenheim, S.; Koval', P.V.; Mueller, G.; Neiva, A.M.R.; Radoslovich, E.W. Nomenclature of the micas. *Clays Clay Miner.* **1998**, *46*, 586–595. [[CrossRef](#)]
196. Lindsley, D.H. *Oxide Minerals: Petrologic and Magnetic Significance*; Mineral Society of America: Chantilly, VA, USA, 1991; Volume 25.
197. Biagioni, C.; Pasero, M. The systematics of the spinel-type minerals: An overview. *Am. Mineral.* **2014**, *99*, 1254–1264. [[CrossRef](#)]
198. Rickard, D. *Pyrite: A Natural History of Fool's Gold*; Oxford University Press: Oxford, UK, 2015; ISBN 0190203684.
199. Rogers, A.F. A review of the amorphous minerals. *J. Geol.* **1917**, *25*, 515–541. [[CrossRef](#)]
200. Morrison, S.M.; Downs, R.T.; Blake, D.F.; Vaniman, D.T.; Ming, D.W.; Hazen, R.M.; Treiman, A.H.; Achilles, C.N.; Yen, A.S.; Morris, R.V.; et al. Crystal chemistry of Martian minerals from Bradbury Landing through Naukluft Plateau, Gale crater, Mars. *Am. Mineral.* **2018**, *103*, 857–871. [[CrossRef](#)]
201. Grew, E.S.; Bada, J.L.; Hazen, R.M. Borate minerals and origin of the RNA world. *Orig. Life Evol. Biosph.* **2011**, *41*, 307–316. [[CrossRef](#)]
202. Rudnick, R.L.; Gao, S. Composition of the continental crust. In *The Crust*; Rudnick, R.L., Holland, H.D., Turekian, K.K., Eds.; Elsevier: Amsterdam, The Netherlands, 2005; Volume 3, pp. 1–64. ISBN 008044847X.
203. Deer, W.A.; Howie, R.A.; Zussman, J. *Rock-forming Minerals: Feldspars*; Geological Society of London: London, UK, 2001; Volume 4A.
204. Hazen, R.M.; Hystad, G.; Golden, J.J.; Hummer, D.R.; Liu, C.; Downs, R.T.; Morrison, S.M.; Ralph, J.; Grew, E.S. Cobalt mineral ecology. *Am. Mineral.* **2017**, *102*, 108–116. [[CrossRef](#)]
205. Herzberg, C.; Vidito, C.; Starkey, N.A. Nickel–cobalt contents of olivine record origins of mantle peridotite and related rocks. *Am. Mineral.* **2016**, *101*, 1952–1966. [[CrossRef](#)]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).