

Short Communication

A Novel Adenovirus in Chinstrap Penguins (*Pygoscelis antarctica*) in Antarctica

Sook-Young Lee ¹, Jeong-Hoon Kim ², Yon Mi Park ¹, Ok Sarah Shin ³, Hankyeom Kim ⁴, Han-Gu Choi ² and Jin-Won Song ^{1,*}

¹ Department of Microbiology, College of Medicine, Institute for Viral Diseases, Korea University, Seoul 136-705, Korea; E-Mails: sylee163@gmail.com (S.-Y.L.); suerte8422@hotmail.com (Y.M.P.)

² Division of Life Sciences, Korea Polar Research Institute, Incheon 406-840, Korea; E-Mails: jhkim94@kopri.re.kr (J.-H.K.); hchoi82@kopri.re.kr (H.-G.C.)

³ Department of Biomedical Science, College of Medicine, Korea University, Seoul 136-705, Korea; E-Mail: oshin@korea.ac.kr

⁴ Department of Pathology, College of Medicine, Korea University, Guro Hospital, Seoul 152-703, Korea; E-Mail: sswords@naver.com

* Author to whom correspondence should be addressed; E-Mail: jwsong@korea.ac.kr; Tel.: +82-2-2286-1011; Fax: +82-2-927-1036.

Received: 19 March 2014; in revised form: 26 April 2014 / Accepted: 28 April 2014 /

Published: 7 May 2014

Abstract: Adenoviruses (family *Adenoviridae*) infect various organ systems and cause diseases in a wide range of host species. In this study, we examined multiple tissues from Chinstrap penguins (*Pygoscelis antarctica*), collected in Antarctica during 2009 and 2010, for the presence of novel adenoviruses by PCR. Analysis of a 855-bp region of the hexon gene of a newly identified adenovirus, designated Chinstrap penguin adenovirus 1 (CSPAdV-1), showed nucleotide (amino acid) sequence identity of 71.8% (65.5%) with South Polar skua 1 (SPSAdV-1), 71% (70%) with raptor adenovirus 1 (RAdV-1), 71.4% (67.6%) with turkey adenovirus 3 (TAdV-3) and 61% (61.6%) with frog adenovirus 1 (FrAdV-1). Based on the genetic and phylogenetic analyses, CSPAdV-1 was classified as a member of the genus, *Siadenovirus*. Virus isolation attempts from kidney homogenates in the MDTC-RP19 (ATCC® CRL-8135™) cell line were unsuccessful. In conclusion, this study provides the first evidence of new adenovirus species in Antarctic penguins.

Keywords: adenovirus; *Siadenovirus*; Chinstrap penguin; Antarctica

1. Introduction

Adenoviruses are linear, double-stranded DNA viruses, with a genome ranging from 26 to 45-kbp and an icosahedral capsid [1]. Adenoviruses, which can infect the respiratory and gastrointestinal tracts, eyes and other organs, cause gastroenteritis and respiratory disease in many species [2]. The family, *Adenoviridae*, comprises five genera: *Mastadenovirus*, *Aviadenovirus*, *Atadenovirus*, *Siadenovirus* and *Ichtadenovirus* [3]. *Mastadenoviruses* infect a wide range of mammalian species, including man, monkey, dog, cattle, swine, mouse and bat [4–10]. *Aviadenovirus* have been identified in birds [11,12], and *Atadenovirus* have been isolated from reptiles, birds and mammals [13–15]. *Siadenovirus* has been detected in birds, frog and a tortoise [16–19], and *Ichtadenovirus* has been detected in fish [20].

Previously, adenoviruses had been isolated from various vertebrate species on all continents, except Antarctica. Recently, the South Polar skua adenovirus 1 (SPSAdV-1), the single known member of the species, *Skua siadenovirus A*, was discovered in dead South Polar skua (*Catharacta maccormicki*) collected near the King Sejong Station in Antarctica [21]. However, there are limited surveillance studies of adenovirus infection in Antarctic wild birds. The South Polar skua can be observed in Antarctica only during the breeding season and feed on the chicks of penguins. Moreover, the South Polar skua shares breeding grounds with the Chinstrap penguin (*Pygoscelis antarctica*), an endemic species [22]. To ascertain if Chinstrap penguins in Antarctica are also infected with adenoviruses, we conducted an exploratory study on samples collected during 2009 and 2010.

2. Results and Discussion

Two of the 10 Chinstrap penguin carcasses (designated CSP09-1 and CSP09-2) were collected in the summer of 2009 and eight (CSP10-1 to CSP10-8) in early 2010. Fifty-six tissues, comprising lung, liver, kidney, heart, intestine, brain, colon, lymph node, spleen, trachea and wounded-bill, were tested. An approximately 1240-bp genomic fragment, including parts of two adjacent genes, that of pVI and hexon, was amplified from 28 tissues (lung, liver, kidney, heart, intestine and/or trachea) from eight Chinstrap penguins (CSP09-1, CSP10-1, CSP10-2, CSP10-3, CSP10-5, CSP10-6, CSP10-7 and CSP10-8), by nested PCR (Table 1). The Chinstrap penguin adenoviruses (CSPAdV-1) from eight Chinstrap penguins (from CSP09-1 to CSP10-8) were designated as CSPAdVno1 to CSPAdVno8, respectively. Adenovirus gene sequences, detected in PCR-positive tissues of individual Chinstrap penguins, were almost identical, suggesting widespread systemic infection.

Pair-wise alignment by ClustalW showed a six-amino acid difference between CSPAdVno1 originating from a Chinstrap penguin collected in 2009 and CSPAdVno2, no4 and no8. Twelve amino acid differences were found between CSPAdVno1 and CSPAdVno3, no5, no6 and no7 (Figure 1). Particularly, CSPAdVno1, no2, no4 and CSPAdVno8 showed a deletion at position 242. As a result of sequence comparison with other types within the genus, *Siadenovirus*, the insertion of amino acid residue S at position 243 was found in the amino acid sequences of some CSPAdV-1 variants. Further


study on the full genome sequence of CSPAdV-1 will be required to support the significance of sequence differences.

Table 1. Detection of Chinstrap penguin adenovirus in various tissues by PCR, with GenBank accession numbers.

Samples No.	Tissues tested	Positives	Designation	Accession No.
CSP09-1	Lu, Li, K, Col, LN, Sp, Br	Lu	CSPAdVno1	KC593379
CSP09-2	W-B	-	-	-
CSP10-1		Lu, Li, K, Ht, Int	CSPAdVno2	KC593380
CSP10-2		Li, K	CSPAdVno3	KC593381
CSP10-3		Lu, Li, K, Int	CSPAdVno4	KC593382
CSP10-4		-	-	-
CSP10-5	Lu, Li, K, Ht, Int, Tr	Lu, K, Int, Tr	CSPAdVno5	KC593383
CPS10-6		Lu, Li, K, Ht, Int, Tr	CSPAdVno6	KC593384
CSP10-7		Ht, Tr	CSPAdVno7	KC593385
CSP10-8		Lu, K, Int, Tr	CSPAdVno8	KC593386

Abbreviations: Lu, lung; Li, liver; K, kidney; Ht, heart; Int, intestine; Tr, trachea; Col, colon; Sp, spleen; Br, brain; LN, lymph node; W-B, wounded-bill.

Figure 1. Differences of the amino acid sequence in the partial hexon gene of novel Chinstrap penguin adenoviruses. Within the 285-amino acid region, six amino acid changes were found between CSPAdVno1 and no2, no4 and no8 (residues 134, 154, 159, 169, 215 and 226), and 12-amino acid differences were found between CSPAdVno1 and no3, no5, no6 and no7 (residues 134, 152, 157, 159, 166, 169, 194, 203, 215, 226, 241 and 242). A deletion of an amino acid at position 242 was found in the hexon gene of CSPAdVno1, no2, no4 and no8 (gray rectangle, deletion of amino acid residue G; CAG, nucleotide sequences in 722–724 nt).



Within the genus, *Siadenovirus*, CSPAdVno1 was shown to have a nucleotide (amino acid) identity of 71.8% (65.5%) with SPSAdV-1, 71% (70%) with raptor adenovirus 1 (RAdV-1), 71.4% (67.6%) with turkey adenovirus 3 (TAdV-3), 69.4% (66.5%) with great tit adenovirus 1 (GTAdV-1) and 61% (61.6%) with frog adenovirus 1 (FrAdV-1). The nucleotide and amino acid sequence identity of each strain of CSPAdV-1 was more than 97% and 96%, respectively (Table 2). However, the nucleotide and amino acid sequences of CSPAdVno3, no5 and no7 and CSPAdVno2, no4 and no8 were identical (Figures 1 and 2). The nucleotide sequence identity of CSPAdV-1 was less (<33%) with other adenovirus genera, such as *Atadenovirus*, *Aviadenovirus* and *Mastadenovirus*. The G+C content of CSPAdVno1 and CSPAdVno2, no4 and no8 was found to be 37.44% and 37.2%, respectively. The partial hexon of CSPAdVno3, no5 and no7 had a G+C content of 37.42%.


Table 2. Nucleotide and amino acid sequence identity (%) between CSPAdVno1 and other CSPAdV-1 strains and representative *Siadenoviruses*.

Virus strain	Sequence identity (%)	
	nucleotide	amino acid
CSPAdVno2	98.7	97.9
CSPAdVno3	97.5	96.1
CSPAdVno4	98.7	97.9
CSPAdVno5	97.5	96.1
CSPAdVno6	97.8	96.1
CSPAdVno7	97.5	96.1
CSPAdVno8	98.7	97.9
SPSAdV-1	71.8	65.5
RAdV-1	71.0	70.0
TAdV-3	71.4	67.6
GTAdV-1	69.4	66.5
FrAdV-1	61.0	61.6

GenBank accession No.: CSPAdVno1 to CSPAdVno8 (KC593379 to KC593386), SPSAdV-1 (HM585353), RAdV-1 (EU715130), TAdV-3 (AC000016), GTAdV-1 (FJ849795), FrAdV-1 (AF224336).

For phylogenetic analysis, approximately 855-bp of the hexon gene, which contains structural loop regions that encode serotype-specific epitopes, were selected [23]. CSPAdV-1 showed the highest similarity with SPSAdV-1 and RAdV-1 and was classified into the genus, *Siadenovirus*, based on phylogenetic trees generated by the maximum-parsimony and neighbor-joining methods, implemented in MEGA5.1 (Molecular Evolutionary Genetics Analysis, 5.1) (Figure 2) and PAUP version 4.0b (Phylogenetic Analysis Using Parsimony, 4.0b) [24,25]. A novel adenovirus species is usually defined as one detected in a new host species and having more than a 5%–15% sequence difference at the nucleotide and amino acid levels compared with previously characterized adenovirus species [3]. Based on these criteria, we conclude that CSPAdV-1 (*Penguin siadenovirus A*) seems to merit the establishment of a new species for it.

Figure 2. Phylogenetic relationship between Chinstrap penguin adenovirus (CSPAdV-1) and other adenoviruses. The phylogenetic tree, based on nucleotide (a) and amino acid sequences (b) of the hexon protein contained loop regions, was generated by the neighbor-joining method. CSPAdVno1 to no8 (KC593379 to KC593386) were compared with adenoviruses of five genera: bat adenovirus 3 (BtAdV-3, GU226970), canine adenovirus 2 (CAdV-2, U77082), bat adenovirus 2 (BtAdV-2, NC015932), porcine adenovirus 5 (PAdV-5, AF289262), porcine adenovirus 3 (PAdV-3, AB126117), bovine adenovirus 10 (BAdV-10, AAF82136), murine adenovirus 1 (MAdV-1, NC000942), human adenovirus 11 (HAdV-11, AY163756), simian adenovirus 25 (SAdV-25, AF394196), human adenovirus 2 (HAdV-2, J01917), human adenovirus 12 (HAdV-12, X73487), human adenovirus 40 (HAdV-40, L19443), bovine adenovirus 3 (BAdV-3, AF030154), tree shrew adenovirus 1 (TSAdV-1, AF258784), duck adenovirus 1 (DAdV-1, Y09598), ovine adenovirus 7 (OAdV-7, U40839), fowl adenovirus 9 (FAdV-9, AF083975), fowl adenovirus 1 (FAdV-1, U46933), fowl adenovirus 4 (FAdV-4, NC015323), frog adenovirus 1 (FrAdV-1, AF224336), great tit adenovirus 1 (GTAdV-1, FJ849795), turkey adenovirus 3 (TAdV-3, AC000016), raptor adenovirus 1 (RAdV-1, EU715130) and South Polar skua adenovirus 1 (SPSAdV-1, HM585353). Scale bars indicate the number of nucleotide and amino acid substitutions per site. Bootstrap values are given at the respective nodes, as determined for 1000 iterations using the MEGA5.1 software [26].



Isolation of CSPAdV-1 was attempted by inoculating the MDTC-RP19 (ATCC® CRL-8135™) cell line [27] with kidney homogenates, but all such attempts were unsuccessful. Failure of the


isolation of RAdV-1 on the chicken embryonic liver cell or MDTC-RP19 cell line has been reported previously [28,29].

3. Experimental Section

3.1. Sample Collection

Chinstrap penguin carcasses were gathered at Narębski Point, located on the southeast coast of Barton Peninsula, King George Island, Antarctica ($62^{\circ}13'40''\text{S}$ – $62^{\circ}14'23''\text{S}$ and $58^{\circ}45'25''\text{W}$ – $58^{\circ}47'00''\text{W}$) during 2009 and early 2010 (Figure 3). All carcasses were identified, weighed and measured. Internal organs (lung, liver, kidney, heart, intestine, trachea, brain, lymph node and spleen) were dissected using sterile instruments and stored at -70°C until use for adenovirus identification.

Figure 3. Collection site of Chinstrap penguin carcasses. Dead Chinstrap penguins were collected at Narębski Point (Antarctic Specially Protected Area; ASPA No. 171, red circle) on King George Island, Antarctica.



3.2. DNA Extraction and PCR

Total genomic DNA was extracted from tissue samples, using the High Pure PCR Template preparation kit (Roche, Indianapolis, IN), according to the manufacturer's instructions. For screening of adenovirus infection, polymerase chain reaction (PCR) was used. PCR assay targeting capsid protein precursor pVI and the capsid protein hexon gene was performed using oligonucleotide primer pairs (outer: 5'-ACC (C/T)GG ATT AGC TGG TGA T-3', 5'-TAA TTT CTG TAT TCC TGT CCT-3'; inner: 5'-CCT GC(A/T) GAT CAA CTG GCT-3', 5'-GGA TCC CTA ACC ATT ATC GTA ATA-3'). The sequences of PCR primers were designated from the conserved region by the alignment of

adenovirus sequences within the genus, *Siadenovirus*. PCR conditions were performed as follows: 1 cycle of 95 °C for 5 min followed by 14 cycles of denaturation at 95 °C for 40 s, one degree step-down each of 1 cycle annealing from 50 °C to 37 °C for 40 s, extension at 72 °C for 1 min, then 25 cycles of denaturation at 95 °C for 40 s, annealing at 42 °C for 40 s, extension at 72 °C for 1 min and, finally, at 72 °C for 5 min in a Mastercycler (Eppendorf, Germany).

3.3. Sequencing and Sequence Analysis

All PCR products were sequenced with the Big Dye terminator v3.1 cycle sequencing kit (ABI) and ABI3730 Automated DNA Sequencer (ABI). Nucleotide sequences were analyzed by ClustalW in MegAlign of DNASTAR programs. Phylogenetic trees were generated by maximum-parsimony and neighbor-joining methods, implemented in MEGA5.1 and PAUP version 4.0b [24,25]. Topologies were evaluated by a bootstrap analysis of 1000 iterations by using MEGA5.1 software [26].

3.4. Isolation Attempts

For isolation attempts of CSPAdV-1, 5% (w/v) kidney homogenates of virus-infected penguins (CSP10-2 and CSP10-3) were inoculated onto MDTC-RP19 (ATCC® CRL-8135™), a lymphoblastoid cell line of turkey origin (*Meleagris gallopavo*) with high susceptibility to TAdV-3, a member of the genus, *Siadenovirus* [27]. Cultures were observed daily for the cytopathic effect, and the cells and supernatant were screened for the adenoviral hexon gene by PCR at each passage.

4. Conclusions

Previous studies on sub-Antarctic and Antarctic penguins have shown that avian species in this region may become infected with various viruses, including paramyxoviruses, Newcastle disease virus, infectious bursal disease virus and influenza viruses [30–38]. The identification of a novel adenovirus species in Chinstrap penguins suggests the possibility of other viruses, including additional previously unrecognized adenoviruses in Antarctic birds.

There is an increasing risk of infectious diseases being introduced into the Antarctic fauna, because of the increased numbers of people travelling to and within Antarctica [38,39]. Consequently, further studies in Antarctic birds may provide new insights into the emergence and dissemination of viral infectious diseases from Antarctica.

Acknowledgments

We thank Sung-Ho Kang, Min-Goo Lee, Kyeong Hoon Cho and Yeong Woong Kim for assistance in the sample collection, Seo Tae-Kun for help with phylogenetic analysis and Richard Yanagihara for editorial assistance. This work was supported by grants PD12010 (Polar Academic Program) and PE13030/PE14020 provided by the Korea Polar Research Institute, Korea.

Author Contributions

S.Y.L. and Y.M.P. performed primer design, DNA extraction, PCR and DNA sequencing reactions. S.Y.L. and O.S.S. performed phylogenetic analysis and S.Y.L. attempt to isolate virus. H.K.K.

performed histopathological study and ecological analysis. J.H.K. and H.G.C. provided penguin samples. J.W.S. conceived the project, and provided overall scientific oversight. All authors contributed to the preparation of the final manuscript.

Conflicts of Interest

The authors declare no conflict of interest.

References and Notes

1. Davison, A.J.; Benkő, M.; Harrach, B. Genetic content and evolution of adenoviruses. *J. Gen. Virol.* **2003**, *84*, 2895–2908.
2. David, M.K.; Peter, M.H.; Diane, E.G.; Robert, A.L.; Malcolm, A.M.; Bernard, R.; Stephen, E.S. Adenoviruses. In *Fields Virology*, 4th ed.; David, M.K., Peter, M.H., Eds.; Lippincott Williams & Wilkins: Philadelphia, PA, USA, 2001; Volume 2, pp. 2404–2408.
3. Harrach, B.; Benkő, M.; Both, G.W.; Brown, M.; Davison, A.J.; Echavarría, M.; Hess, M.; Jones, M.S.; Kajon, A.; Lehmkuhl, H.D.; *et al.* Family adenoviridae. In *Virus Taxonomy: IXth Report of the International Committee on Taxonomy of Viruses.*; King, A.M.Q., Lefkowitz, E., Adams, M.J., Carstens, E.B., Eds.; Elsevier: New York, NY, USA, 2011; Volume 9, pp. 125–141.
4. Kovács, G.M.; Davison, A.J.; Zakhartchouk, A.N.; Harrach, B. Analysis of the first complete genome sequence of an Old World monkey adenovirus reveals a lineage distinct from the six human adenovirus species. *J. Gen. Virol.* **2004**, *85*, 2799–2807.
5. Morrison, M.D.; Onions, D.E.; Nicolson, L. Complete DNA sequence of canine adenovirus type 1. *J. Gen. Virol.* **1997**, *78*, 873–878.
6. Rusvai, M.; Harrach, B.; Bánrévi, A.; Evans, P.S.; Benkő, M. Identification and sequence analysis of the core protein genes of bovine adenovirus 2. *Virus Res.* **2000**, *70*, 25–30.
7. Aggarwal, N.; Mittal, S.K. Sequence analysis of procine adenovirus type 3 E1 region, pIX and pIVa2 genes, and two novel open reading frames. *Intervirology* **2000**, *43*, 6–12.
8. Klempa, B.; Krüger, D.H.; Auste, B.; Stanko, M.; Krawczyk, A.; Nickel, K.F.; Uberta, K.; Stang, A. A novel cardiotropic murine adenovirus representing a distinct species of mastadenoviruses. *J. Virol.* **2009**, *83*, 5749–5759.
9. Maeda, K.; Hondo, E.; Terakawa, J.; Kiso, Y.; Nakaichi, N.; Endoh, D.; Sakai, K.; Morikawa, S.; Mizutani, T. Isolation of novel adenovirus from fruit bat (*Pteropus dasymallus yayeyamae*). *Emerg. Infect. Dis.* **2008**, *14*, 347–249.
10. Sonntag, M.; Mühldorfer, K.; Speck, S.; Wibbelt, G.; Kurth, A. New adenovirus in bats, Germany. *Emerg. Infect. Dis.* **2009**, *15*, 2052–2055.
11. Schrenzel, M.; Oaks, J.L.; Rotstein, D.; Maalouf, G.; Snook, E.; Sandfort, C.; Rideout, B. Characterization of a new species of adenovirus in falcons. *J. Clin. Microbiol.* **2005**, *43*, 3402–3412.
12. Chiocca, S.; Kurzbauer, R.; Schaffner, G.; Baker, A.; Mautner, V.; Cotton, M. The complete DNA sequence and genomic organization of the avian adenovirus CELO. *J. Virol.* **1996**, *70*, 2939–2949.

13. Dán, A.; Ruzsics, Z.; Russell, W.C.; Benkő, M.; Harrach, B. Analysis of the hexon gene sequence of bovine adenovirus type 4 provides further support for a new adenovirus genus (*Atadenovirus*). *J. Gen. Virol.* **1998**, *79*, 1453–1460.
14. Farkas, S.L.; Harrach, B.; Benkő, M. Completion of the genome analysis of snake adenovirus type 1, a representative of the reptilian lineage within the novel genus *Atadenovirus*. *Virus Res.* **2008**, *132*, 132–139.
15. Hess, M.; Blöcker, H.; Brandt, P. The complete nucleotide sequence of the egg drop syndrome virus: An intermediate between mastadenoviruses and aviadenoviruses. *Virology* **1997**, *238*, 145–156.
16. Davison, A.J.; Wright, K.M.; Harrach, B. DNA sequence of frog adenovirus. *J. Gen. Virol.* **2000**, *81*, 2431–2439.
17. Pitcovski, J.; Mualem, M.; Rei-Koren, Z.; Krispel, S.; Shmueli, E.; Peretz, Y.; Gutter, B.; Gallili, G.E.; Michael, A.; Goldberg, D. The complete DNA sequence and genome organization of the avian adenovirus, hemorrhagic enteritis virus. *Virology* **1998**, *249*, 307–315.
18. Kovács, E.R.; Benkő, M. Complete sequence of raptor adenovirus 1 confirms the characteristic genome organization of siadenoviruses. *Infect. Genet. Evol.* **2011**, *5*, 1058–1065.
19. Kovács, E.R.; Jánoska, M.; Harrach, B.; Benkő, M. Recognition and partial genome characterization by non-specific DNA amplification and PCR of a new siadenovirus species in a sample originating from *Parus major*, a great tit. *J. Virol. Methods* **2010**, *163*, 262–268.
20. Kovács, G.M.; LaPatra, S.E.; D'Halluin, J.C.; Benkő, M. Phylogenetic analysis of the hexon and protease genes of a fish adenovirus isolated from white sturgeon (*Acipenser transmontanus*) supports the proposal for a new adenovirus genus. *Virus Res.* **2003**, *98*, 27–34.
21. Park, Y.M.; Kim, J.H.; Gu, S.H.; Lee, S.Y.; Lee, M.G.; Kang, Y.K.; Kang, S.H.; Kim, H.J.; Song, J.W. Full genome analysis of a novel adenovirus from the South Polar skua (*Catharacta maccormicki*) in Antarctica. *Virology* **2012**, *422*, 144–150.
22. Yogui, G.T.; Sericano, J.L. Levels and pattern of polybrominated diphenyl ethers in eggs of Antarctic seabirds: Endemic versus migratory species. *Environ. Pollut.* **2009**, *157*, 975–980.
23. Raue, R.; Gerlach, H.; Müller, H. Phylogenetic analysis of the hexon loop 1 region of an adenovirus from psittacine birds supports the existence of a new psittacine adenovirus (PsAdV). *Arch. Virol.* **2005**, *150*, 1933–1943.
24. Tamura, K.; Peterson, D.; Peterson, N.; Stecher, G.; Nei, M.; Kumar, S. MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionally Distance, and Maximum Parsimony methods. *Mol. Biol. Evol.* **2011**, *28*, 2731–2739.
25. *PAUP*4.0: Phylogenetic Analysis Using Parsimony*, version 4.0; Sinauer Associates, Inc: Sunderland, MA, USA, 2002.
26. *MEGA: Molecular Evolutionary Genetics Analysis*. Available online: <http://www.megasoftware.net/> (accessed on 8 March 2013).
27. Nazerian, K.; Fadly, A.M. Propagation of virulent and avirulent turkey hemorrhagic enteritis virus in cell culture. *Avian. Dis.* **1982**, *26*, 816–827.

28. Zsivanovits, P.; Monks, D.J.; Forbes, N.A.; Ursu, K.; Raue, R.; Benkő, M. Presumptive identification of a novel adenovirus in a Harris hawk (*Parabuteo unicinctus*), a Bengal eagle owl (*Bubo bengalensis*), and a Verreaux's eagle owl (*Bubo lacteus*). *J. Avian Med. Sug.* **2006**, *20*, 105–112.
29. Kovács, E.R.; Benkő, M. Confirmation of a novel siadenovirus species detected in raptors: partial sequence and phylogenetic analysis. *Virus Res.* **2009**, *140*, 64–70.
30. Barbosa, A.; Palacios, M.J. Health of Antarctic birds: A review of their parasites, pathogens and diseases. *Polar Biol.* **2009**, *32*, 1095–1115.
31. Pearce, D.A.; Wilson, W.H. Viruses in Antarctic ecosystems. *Antarctic Sci.* **2003**, *15*, 319–331.
32. Alexander, D.J.; Manvell, R.J.; Collins, M.S.; Brockman, S.J.; Westbury, H.A.; Morgan, I.; Austin, F.J. Characterization of paramyxoviruses isolated from penguins in Antarctica and sub-Antarctica during 1976–1979. *Arch. Virol.* **1989**, *109*, 135–143.
33. Austin, F.J.; Webster, R.G. Evidence of ortho and paramyxoviruses in fauna from Antarctica. *J. Wild. Dis.* **1993**, *29*, 568–571.
34. Thomazelli, L.M.; Araujo, J.; Oliveira, D.B.; Sanfilippo, L.; Ferreira, C.S.; Brentano, L.; Pelizari, V.H.; Nakayama, C.; Duarte, R.; Hurtado, R.; *et al.* Newcastle disease virus in penguins from King George Island on the Antarctic region. *Vet. Microbiol.* **2010**, *146*, 155–160.
35. Wallensten, A.; Munster, V.J.; Osterhaus, A.D.M.E.; Waldenström, J.; Bonnedahl, J.; Broman, T.; Fouchier, R.A.M.; Olson, B. Mounting evidence for the presence of influenza A virus in the avifauna of the Antarctic region. *Antarctic Sci.* **2006**, *18*, 353–356.
36. Chang, C.M.; Lebarbenchon, C.; Gauthier-Clerc, M.; Bohec, C.L.; Beaune, D.; Maho, Y.L.; Werf, S. Molecular surveillance for avian influenza A virus in king penguins (*Aptenodytes patagonicus*). *Polar Biol.* **2009**, *32*, 663–665.
37. Abad, F.X.; Busquets, N.; Sanchez, A.; Ryan, P.G.; Majö, N.; Gonzalez-Solis, J. Serological and virological surveys of the influenza A viruses in Antarctic and sub-Antarctic penguins. *Antarctic Sci.* **2013**, *25*, 339–344.
38. Kerry, K.; Riddle, M.; Clarke, J. *Diseases of Antarctic wildlife*; 18; Australian Antarctic Division: Channel Highway, Kingston, Australia, August 1998; pp. 89–91.
39. Chown, S.L.; Lee, J.E.; Hughes, K.A.; Barrett, P.J.; Bergstrom, D.M.; Convey, P.; Cowan, D.A.; Crosbie, K.; Dyer, G.; Frenot, Y.; *et al.* Challenges to the future conservation of the Antarctica. *Science* **2012**, *337*, 158–159.